

# A universal $U(1)$ -RCC invariant of links and rationality conjecture.

L. Rozansky<sup>1</sup>

*Department of Mathematics, University of North Carolina*

*CB #3250, Phillips Hall*

*Chapel Hill, NC 27599*

*E-mail address: rozansky@math.unc.edu*

## Abstract

We define a graph algebra version of the stationary phase integration over the coadjoint orbits in the Reshetikhin formula for the colored Jones-HOMFLY polynomial. As a result, we obtain a ‘universal’  $U(1)$ -RCC invariant of links in rational homology spheres, which determines the  $U(1)$ -RCC invariants based on simple Lie algebras. We formulate a rationality conjecture about the structure of this invariant.

---

<sup>1</sup>This work was supported by NSF Grants DMS-0196235 and DMS-0196131

# Contents

|          |                                                                                                                                                     |           |
|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                                                                                                                 | <b>2</b>  |
| 1.1      | Motivation . . . . .                                                                                                                                | 2         |
| 1.2      | Results . . . . .                                                                                                                                   | 5         |
| 1.3      | Notations . . . . .                                                                                                                                 | 6         |
| <b>2</b> | <b>A Lie algebra based <math>U(1)</math>-RCC invariant</b>                                                                                          | <b>7</b>  |
| 2.1      | Kontsevich integral and colored Jones-HOMFLY polynomial . . . . .                                                                                   | 7         |
| 2.2      | Wheeling, Kirillov integral and Reshetikhin formula . . . . .                                                                                       | 10        |
| 2.3      | Invariant measure on a coadjoint orbit . . . . .                                                                                                    | 14        |
| 2.4      | $\mathfrak{g}$ -based $U(1)$ -RCC invariant as a stationary phase integral . . . . .                                                                | 18        |
| <b>3</b> | <b>Graph algebras <math>\mathcal{B}_\mathfrak{g}^C</math>, <math>\mathcal{D}_\mathfrak{g}^C</math> and <math>\mathcal{QD}_\mathfrak{g}^C</math></b> | <b>23</b> |
| 3.1      | Root and Cartan edges of algebra $\mathcal{B}_\mathfrak{g}^C$ . . . . .                                                                             | 23        |
| 3.2      | Injectons . . . . .                                                                                                                                 | 25        |
| 3.3      | Symmetric algebras of cohomologies of graphs . . . . .                                                                                              | 29        |
| 3.4      | Converting Cartan legs into cohomology . . . . .                                                                                                    | 33        |
| 3.5      | Edge-related denominators . . . . .                                                                                                                 | 37        |
| 3.6      | Gluing of legs . . . . .                                                                                                                            | 41        |
| 3.7      | Weight system . . . . .                                                                                                                             | 44        |
| <b>4</b> | <b>Calculus and differential geometry of graph algebras</b>                                                                                         | <b>45</b> |
| 4.1      | Functions and general tensor fields . . . . .                                                                                                       | 45        |
| 4.2      | Vector fields . . . . .                                                                                                                             | 47        |
| 4.3      | Diffeomorphisms . . . . .                                                                                                                           | 48        |
| 4.4      | Derivatives and matrix fields . . . . .                                                                                                             | 53        |
| 4.5      | Determinant and its properties . . . . .                                                                                                            | 55        |
| 4.6      | Integration measure . . . . .                                                                                                                       | 61        |
| 4.7      | A formal gaussian integral . . . . .                                                                                                                | 62        |
| 4.8      | Properties of the integral . . . . .                                                                                                                | 63        |

|          |                                                                                                |           |
|----------|------------------------------------------------------------------------------------------------|-----------|
| <b>5</b> | <b>A formal integration over a coadjoint orbit</b>                                             | <b>70</b> |
| 5.1      | Invariant integration measure on a coadjoint orbit . . . . .                                   | 70        |
| 5.2      | Properties of the coadjoint orbit integral . . . . .                                           | 76        |
| <b>6</b> | <b>A universal <math>U(1)</math>-RCC invariant</b>                                             | <b>82</b> |
| 6.1      | Definition and invariance . . . . .                                                            | 82        |
| 6.2      | Haircomb graphs, wheels and Kontsevich integral formula for the Alexander polynomial . . . . . | 88        |
| 6.3      | Alexander polynomial and the structure of the universal $U(1)$ -RCC invariant                  | 92        |
| 6.4      | The rationality conjecture . . . . .                                                           | 95        |
| <b>A</b> | <b>A superalgebra <math>u(1 1)</math> weight system and the Alexander polynomial of a link</b> | <b>97</b> |

# 1 Introduction

## 1.1 Motivation

A classical topology interpretation of ‘quantum’ invariants of knots, links and 3-manifolds still remains an open question. A discovery of the Alexander polynomial inside the colored Jones polynomial as well as a discovery of the Casson-Walker invariant inside the Witten-Reshetikhin-Turaev invariant of rational homology spheres suggests that quantum invariants are somehow packed up with classical invariants (both known and unknown). Thus the appropriate unpacking of quantum invariants may lead to a discovery of new classical invariants which may, in turn, provide a solution to long-standing puzzles of 3-dimensional topology.

The main method of unpacking the quantum invariants is splitting them into invariants of finite type. For rational homology spheres, this procedure seems to correspond to the isolation of the contribution of the trivial connection to the Witten-Reshetikhin-Turaev invariant in the semi-classical limit of  $q \rightarrow 1$ . It turns out that the best way of implementing the same procedure for the Jones polynomial of knots and links is *not* to study the trivial connection contribution in the limit  $q \rightarrow 1$ ,  $\alpha_1, \dots, \alpha_L = \text{const}$  (which would amount to simply expanding the colored Jones polynomial in powers of  $q - 1$ ), but rather to isolate

a contribution of the  $U(1)$ -reducible flat connections in the knot or link complement (we abbreviate this contribution as  $U(1)$ -RCC) to the colored Jones polynomial in the limit

$$q \longrightarrow 1, \quad q^{\alpha_1}, \dots, q^{\alpha_L} = \text{const}, \quad (1.1)$$

where  $\alpha_1, \dots, \alpha_N$  are colors (dimensions of  $SU(2)$  modules) assigned to the components of an  $L$ -component link. This approach yields the following formula [11] for the colored Jones polynomial of a knot  $\mathcal{K} \subset S^3$ :

$$J_\alpha(\mathcal{K}; q) = \frac{[\alpha]}{\Delta_A(\mathcal{K}; q^\alpha)} \left( 1 + \sum_{n=1}^{\infty} \frac{P_n(\mathcal{K}; q^\alpha)}{\Delta_A^{2n}(\mathcal{K}; q^\alpha)} (q-1)^n \right). \quad (1.2)$$

Here  $[\alpha] = (q^\alpha - q^{-\alpha})/(q - q^{-1})$ ,  $\Delta_A(\mathcal{K}; t)$  is the Alexander polynomial of  $\mathcal{K}$  normalized by the condition  $\Delta_A(\text{unknot}; t) = 1$  and

$$P_n(\mathcal{K}; t) \in \mathbb{Z}[t, t^{-1}] \quad (1.3)$$

are polynomial invariants of  $\mathcal{K}$  which, due to the manner of their appearance in eq.(1.2), may have a classical topology interpretation similar to that of the Alexander polynomial and Casson-Walker invariant. The formula (1.2) itself should be interpreted as an equation between the expansion of both sides in powers of  $q-1$  and  $\alpha$  (known as the Melvin-Morton expansion of the colored Jones polynomial).

Since the expansion of  $J_\alpha(\mathcal{K}; q)$  in powers of  $q-1$  can be deduced from the Kontsevich integral of  $\mathcal{K}$ , then equation (1.2) implies that Kontsevich integral has a special ‘rational’ structure which matches the structure of the *r.h.s.* of eq.(1.2). The corresponding conjecture about Kontsevich integral was formulated in [14] and A. Kriker proved it in [7] (a stronger version was proved by S. Garoufalidis and A. Kriker in [4]). The conjecture states that the coefficients at individual graphs in the Kontsevich integral of  $\mathcal{K}$  can be assembled into ‘universal’ polynomials which determine the polynomials  $P_n(\mathcal{K}; t)$  of expansions of the type (1.2) for colored Jones-HOMFLY polynomials based on any simple Lie algebra.

The case of links is less simple than that of knots. On one hand, S. Garoufalidis and A. Kriker [4] proved the rationality of Kontsevich integral for boundary links. On the other hand, the  $R$ -matrix calculation, which led to eq.(1.2) for knots, can be repeated for the opposite class of ‘sufficiently connected’ links (that is, links whose Alexander polynomial is not identically equal to 0). This  $R$ -matrix calculation yields a  $U(1)$ -RCC invariant of an oriented link  $\mathcal{L}$ , which has the form

$$\check{J}^{(r)}(\mathcal{L}; t_1, \dots, t_L; h) = \frac{1}{h} \frac{1}{\nabla_A(\mathcal{L}; t_1, \dots, t_L)} \left( 1 + \sum_{n=1}^{\infty} \frac{P_n(\mathcal{L}; t_1, \dots, t_L)}{\nabla_A^{2n}(\mathcal{L}; t_1, \dots, t_L)} h^n \right), \quad (1.4)$$

where  $\nabla_A(\mathcal{L}; t_1, \dots, t_L)$  is the Alexander ‘polynomial’ of  $\mathcal{L}$  in the normalization

$$\nabla_A(\text{unknot}; t) = \frac{1}{(t^{1/2} - t^{-1/2})},$$

which is appropriate for links, while

$$P_n(\mathcal{L}; t_1, \dots, t_L) \in \mathbb{Z}[t_1^{\pm 1/2}, \dots, t_L^{\pm 1/2}, 1/2] \quad (1.5)$$

are polynomial invariants of  $\mathcal{L}$ . Path integral arguments suggest that the expression

$$q^{\frac{1}{4} \sum_{i,j=1}^L l_{ij} \alpha_i \alpha_j} \check{J}^{(r)}(\mathcal{L}; q^\alpha; q-1) \quad (1.6)$$

represents a contribution of one of  $2^{L-1}$   $U(1)$ -reducible flat connections in  $S^3 \setminus \mathcal{L}$  with appropriate monodromies to the colored Jones polynomial  $J_\alpha(\mathcal{L}; q)$  in the semi-classical limit (1.1). In contrast to the knot case, the expression (1.6) does not determine the whole Jones polynomial through a simple relation similar to (1.2)<sup>2</sup>. Also, deriving the series (1.4) from the Kontsevich integral of  $\mathcal{L}$  requires a few extra steps outlined in [13]:

$$\begin{array}{ccccccc} \text{link } \mathcal{L} & \xrightarrow{1} & \begin{array}{c} \text{Kontsevich} \\ \text{integral} \\ \text{of } \mathcal{L} \text{ in } \mathcal{B} \end{array} & \xrightarrow{2} & \begin{array}{c} \text{Wheeled} \\ \text{Kontsevich} \\ \text{integral} \\ \text{of } \mathcal{L} \text{ in } \mathcal{B} \end{array} & \xrightarrow{3} & \begin{array}{c} \text{Reshetikhin} \\ \text{integrand} \\ \text{for } J_\alpha(\mathcal{L}; q) \end{array} & \xrightarrow{4} & \begin{array}{c} U(1)\text{-RCC} \\ \text{invariant} \\ \text{of } \mathcal{L} \end{array} \end{array} \quad (1.7)$$

Here  $\mathcal{B}$  is the space of (1,3)-valent graphs with cyclic ordering at 3-valent vertices, and 1-valent vertices labeled by (that is, assigned to) components of  $\mathcal{L}$ . Map 1 is a standard combination of Kontsevich integral (which maps  $\mathcal{L}$  into the space  $\mathcal{A}$ ) and a PBW symmetrization map (which identifies  $\mathcal{A}$  with  $\mathcal{B}$ ). Map 2 is the wheeling map defined in [2] (it maps  $\mathcal{B}$  into  $\mathcal{B}$ ). Map 3 is a link version of the application of the ‘wheeled’  $su(2)$  weight system. Instead of simply attaching the shifted highest weights of  $su(2)$  modules to the 1-valent vertices of the graphs (as we did in the case of knots), we also have to perform a Kirillov-type integration over their (co-)adjoint orbits in order to recover the colored Jones polynomial. In the limit (1.1), the stationary phase approximation may be applied to this integral. Map 4 picks up a contribution of a particular stationary phase point corresponding to all integrands belonging to the same Cartan subalgebra.

The goal of this paper is to remove the Lie algebra from the maps 3,4. In other words, we will describe the stationary phase integration in Reshetikhin integral purely in diagrammatic

---

<sup>2</sup>A knot relation (1.2) is rather an exception which, from the path integral point of view, is due to the fact that a  $U(1)$ -reducible connection is the only flat connection in the knot complement when the monodromy along the meridian of the knot is sufficiently small.

terms. This integration will become a map from the gaussian expressions in  $\mathcal{B}$  into a new space  $\mathcal{QD}$ . The image of  $\mathcal{L}$  in  $\mathcal{QD}$  will be a ‘universal’  $U(1)$ -RCC invariant in the following sense: for any simple Lie algebra  $\mathfrak{g}$  and for any assignment of its modules to components of  $\mathcal{L}$ , the  $\mathfrak{g}$ -based  $U(1)$ -RCC invariant can be obtained by applying an appropriate weight system to the image of  $\mathcal{L}$  in  $\mathcal{QD}$ , so that instead of the sequence (1.7) we may use a new one

$$\begin{array}{ccccccc}
 & & \text{Wheeled} & & \text{Universal} & & U(1)\text{-RCC invariant} \\
 & & \text{Kontsevich} & & U(1)\text{-RCC} & & \\
 \text{link } \mathcal{L} & \xrightarrow{1} & \text{Kontsevich} & \xrightarrow{2} & \text{integral} & \xrightarrow{5} & \text{related to} \\
 & & \text{integral} & & \text{invariant} & & \text{the } \mathfrak{g}\text{-Jones} \\
 & & \text{of } \mathcal{L} \text{ in } \mathcal{B} & & \text{of } \mathcal{L} \text{ in } \mathcal{QD} & & \text{polynomial of } \mathcal{L} \\
 & & & & & & \\
 & & & & & & \text{polynomial of } \mathcal{L}
 \end{array} \quad (1.8)$$

where map 5 is the diagrammatic stationary phase integration over the (co-)adjoint orbits and map 6 is the application of  $\mathfrak{g}$  weight system to an element of  $\mathcal{QD}$ .

## 1.2 Results

Here is the outline of the paper. In Section 2 we define the Lie algebra based  $U(1)$ -RCC invariant by performing the stationary phase integration in a familiar context of integration over the coadjoint orbits. In Sections 3–5 we translate this calculation into the pure language of graphs and then in Section 6 we prove the topological invariance of graph  $U(1)$ -RCC invariant.

We begin Section 2 by recalling the basic facts about a Kontsevich integral of a link in  $S^3$  and in a rational homology sphere. We look at the relation between Kontsevich integral and the Melvin-Morton expansion of the Jones-HOMFLY polynomial. We modify it by using Kirillov integrals instead of traces in representations and thus come to the Reshetikhin formula for the Melvin-Morton expansion. By rescaling the integration variables, we make the Reshetikhin integral suitable for the stationary phase approximation and define the  $\mathfrak{g}$ -based  $U(1)$ -RCC invariant as a contribution of a particular stationary phase point (when all the variables belong to the same Cartan subalgebra).

In Section 3 we introduce a new graph algebra in which the edges are split into their root and Cartan parts. Then we encode Cartan legs of a graph as elements of that graph’s cohomology. Thus Cartan legs are converted into a symmetric algebra of the graph cohomology space. We extend that symmetric algebra by allowing division by polynomials depending on individual edges. Finally, we formulate basic graph operations of disconnected union and leg gluing in terms of the cohomology.

In Section 4 we define the basic elements of differential geometry (such as functions,

tensors, vector fields and matrix fields) on our graph algebras. We also define a determinant of a matrix field and a formal gaussian integral and establish their properties.

In Section 5 we concentrate on the differential geometry of the graph algebra version of a coadjoint orbit. We find an invariant measure on the orbit and study the properties of related gaussian integrals.

In Section 6 we apply all this knowledge to define a universal  $U(1)$ -RCC invariant of an oriented link as an integral over the ‘graph’ coadjoint orbits. We describe the structure of this invariant and conjecture its rationality.

The main results of the paper are Theorems 6.5 and 6.6, which establish the topological invariance of the  $U(1)$ -RCC invariant, Theorem 6.13 which relates the universal  $U(1)$ -RCC invariant to its Lie algebra based analog, and the rationality conjecture 6.19, which suggests that the  $U(1)$ -RCC invariant is a source of interesting polynomial invariants of links, possibly with a nice topological interpretation. As a by-product, we establish Theorem 6.15 which expresses the Alexander polynomial of a link in terms of a tree and 1-loop parts of its Kontsevich integral. We give an alternative proof of this theorem in Appendix A. Finally, while considering the simplest example of the graph coadjoint orbit integral, we conjecture a graph version of the Duistermaat-Heckmann theorem (Conjecture 5.12).

### 1.3 Notations

Throughout the paper we will work with (1,3)-valent graphs. We assume that every graph is endowed with a cyclic order of edges at 3-valent vertices. If that order changes at one vertex, then we assume that the whole graph acquires a minus sign. We refer to the edges incident to 1-valent vertices as *legs* and we call the edges incident to two 3-valent vertices *internal*. A graph which consists of a single edge, is called a *strut*.

An edge of a graph is called a *bridge*, if its removal increases the number of connected components of the graph. A sequence of edges is called a *path*, if the beginning of the next edge coincides with the end of the previous edge. A path without self-intersections is called a *chain*. A closed chain is called a *cycle*. A *haircomb* is a graph consisting of an open chain and legs attached to it. A *wheel* is a graph consisting of a cycle and legs attached to it.

We call an element of a graph algebra *narrow* if it is a linear combination of connected graphs.

We use multi-index notations:  $\mathbf{x} = x_1, \dots, x_L$ , where in topological applications  $L$  is usually a number of components of a link  $\mathcal{L}$ .  $\mathbf{x}_{(i)}$  denotes the same sequence  $x_1, \dots, x_L$  but

with  $x_i$  removed. We also use the following notations for exponentials and scalar products

$$e^{\mathbf{x}} = e^{x_1}, \dots, e^{x_L}, \quad (\mathbf{x}, y) = (x_1, y), \dots, (x_L, y). \quad (1.9)$$

$\mathfrak{G}$  is a simple Lie group,  $\mathfrak{g}$  is its Lie algebra.  $\mathfrak{T} \subset \mathfrak{G}$  is a maximal torus and  $\mathfrak{h} \subset \mathfrak{g}$  is a corresponding Cartan subalgebra.  $\mathfrak{g}^*$  and  $\mathfrak{h}^*$  are dual spaces of  $\mathfrak{g}$  and  $\mathfrak{h}$ . Lie algebra splits into a sum  $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{r}$ , where  $\mathfrak{r}$  is the span of the root spaces. We assume that  $\mathfrak{g}$  has an invariant scalar product (normalized in such a way that the length of short roots is  $\sqrt{2}$ ), so we do not always distinguish  $\mathfrak{g}$  from  $\mathfrak{g}^*$ . We use vector notations  $\vec{x}$  for their elements.  $\Delta_{\mathfrak{g}} \subset \mathfrak{h}^*$  denotes the set of all roots of  $\mathfrak{g}$ ,  $\Delta_+ \subset \mathfrak{h}^*$  denotes the set of positive roots of  $\mathfrak{g}$ , and we denote  $\vec{\rho} = \sum_{\vec{\lambda} \in \Delta_+} \vec{\lambda}$ . Sometimes we will assume that  $\Delta_{\mathfrak{g}}, \Delta_+ \subset \mathfrak{h}$ , root being transferred from  $\mathfrak{h}^*$  to  $\mathfrak{h}$  with the help of the scalar product.  $V_{\vec{\alpha}}$  denotes a module of  $\mathfrak{g}$  with highest weight  $\vec{\alpha} - \vec{\rho}$ .

$\mathcal{L}$  denotes an  $L$ -component link either in  $S^3$  or in a rational homology sphere  $M$ . The numbers  $\ell_{ij}$  denote the linking numbers between its components.

For a ring  $R$ ,  $\mathcal{Q}(R)$  denotes the field of its fractions.

## 2 A Lie algebra based $U(1)$ -RCC invariant

### 2.1 Kontsevich integral and colored Jones-HOMFLY polynomial

Let us recall some basic facts about Kontsevich integral of links (we refer the reader to [3] and references therein). We will work with links in rational homology spheres, but we start with a link in  $S^3$ . Let  $\mathcal{L}$  be an  $L$ -component oriented *dotted Morse link*. In other words, we assume that  $\mathcal{L}$  is imbedded in  $\mathbb{R}^3$ , and a dot is marked on each component. Then Kontsevich integral  $Z_{\mathcal{A}}(\mathcal{L})$  of  $\mathcal{L}$  takes value in the space  $\mathcal{A}(\uparrow_1 \cdots \uparrow_L)$ . This is a (factored over the STU relations) space of (1,3)-valent graphs, whose 1-valent vertices are placed on  $L$  oriented segments, which correspond to the link components. The paper [3] uses the notation  $\mathcal{A}(\uparrow_{\mathfrak{L}})$ , where  $\mathfrak{L}$  is a finite set of labels, in our case  $X$  obviously denotes the set of link components, so

$$\mathfrak{L} = \{1, \dots, L\}; \quad (2.1)$$

we will abuse notations by using  $\mathcal{L}$  directly instead of  $\mathfrak{L}$ , thus writing  $\mathcal{A}(\uparrow_{\mathcal{L}})$  when we deal with Kontsevich integral of  $\mathcal{L}$ .



Note that  $Z_{\mathcal{A}}(\mathcal{L})$  is not an invariant of the link  $\mathcal{L}$ , because it depends on the positions of the dots on the link components. This dependence could be eliminated, had we defined Kontsevich integral as taking values in the space  $\mathcal{A}(\bigcirc_1, \dots, \bigcirc_L)$ , in which the 1-valent vertices of (1,3)-valent graphs are placed on oriented circles. However, we have to keep the dots, since we are about to pass from  $\mathcal{A}(\uparrow_{\mathcal{L}})$  to  $\mathcal{B}_{\mathcal{L}}$ .

The space  $\tilde{\mathcal{B}}_{\mathfrak{L}}$  is a space of (1,3)-valent graphs, whose 1-valent vertices (or, equivalently, legs) are assigned to the elements of  $\mathfrak{L}$ . The vertices which carry the same label (that is, element of  $\mathfrak{L}$ ), are considered equivalent and may be permuted when an isomorphism between two graphs is considered. The space  $\mathcal{B}_{\mathfrak{L}}$  is a quotient of  $\tilde{\mathcal{B}}_{\mathfrak{L}}$  over its subspace  $\tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}}$ , which is a span of the IHX relation.

The space  $\mathcal{B}_{\mathfrak{L}}$  is a bi-graded algebra. The multiplication of two graphs is defined as their disjoint union. The unit of this multiplication is the empty graph, and we denote it as  $\mathbf{1}$ . The two gradings that are preserved by the multiplication of  $\mathcal{B}_{\mathfrak{L}}$  are

$$\deg_1 D = \#(\text{edges}) - \#(3\text{-vertices}) = \frac{1}{2} \#(\text{all vertices}) = \#(1\text{-vertices}) + \chi(D), \quad (2.2)$$

$$\deg_2 D = \#(\text{edges}) - \#(\text{all vertices}) = \chi(D) = \deg_1 D - \#(1\text{-vertices}), \quad (2.3)$$

where  $\chi(D) = \#(\text{edges}) - \#(\text{all vertices})$  is the (*opposite*) Euler characteristic of  $D$ . The grading  $\deg_1$  is also compatible with the multiplication in the algebra  $\mathcal{A}(\uparrow_{\mathcal{L}})$ .

A transition from  $\mathcal{A}(\uparrow_{\mathfrak{L}})$  to  $\mathcal{B}_{\mathfrak{L}}$  is produced by the Poincare-Birkhoff-Witt symmetrization of the order in which 1-valent vertices appear on the oriented segments. The inverse map is denoted as

$$\hat{\chi}: \mathcal{B}_{\mathfrak{L}} \rightarrow \mathcal{A}(\uparrow_{\mathfrak{L}}). \quad (2.4)$$

We define  $Z_{\mathcal{B}}(\mathcal{L})$  as the Kontsevich integral of  $\mathcal{L}$  with value in  $\mathcal{B}_{\mathcal{L}}$ , which in line with (2.1) is, of course, a space of (1,3)-valent graphs, whose 1-valent vertices are labeled by components of  $\mathcal{L}$  or simply by the numbers  $1, \dots, L$ . In other words,

$$Z_{\mathcal{A}}(\mathcal{L}) = \hat{\chi}(Z_{\mathcal{B}}(\mathcal{L})). \quad (2.5)$$

If  $\mathcal{L}$  is in a rational homology sphere  $M$ , then we present it as a surgery on the framed  $\mathcal{L}'$  part of a combined link  $\mathcal{L} \cup \mathcal{L}' \in S^3$ . Then  $Z_{\mathcal{A}}(\mathcal{L}, M)$  is defined with the help of the LMO map [8] or its Århus version [3]. Namely, we first symmetrize all legs of  $Z_{\mathcal{A}}(\mathcal{L} \cup \mathcal{L}')$  which are attached to the components of  $\mathcal{L}'$ , and then we apply the LMO or Århus map to those legs. The result is  $Z_{\mathcal{A}}(\mathcal{L}, M) \in \mathcal{A}(\uparrow_{\mathcal{L}})$ . Then we pass from it to  $Z_{\mathcal{B}}(\mathcal{L}, M) \in \mathcal{B}_{\mathcal{L}}$  in exactly the same way as we did it for  $\mathcal{L} \in S^3$ . Therefore from now on we consider  $\mathcal{L}$  to be a link in

a rational homology sphere  $M$ , and since  $M$  will always be the same, we drop it from our notations.

By using the fact that  $Z_{\mathcal{B}}(\mathcal{L})$  (by definition) has coefficient 1 at the empty graph, we can define the logarithm of  $Z_{\mathcal{B}}(\mathcal{L})$

$$W(\mathcal{L}) = \log Z_{\mathcal{B}}(\mathcal{L}) \quad (2.6)$$

by the formula  $\log(1+x) = \sum_{n=1}^{\infty} (-1)^{n-1} x^n / n$ . We call an element of a graph algebra *narrow*, if it can be presented as a linear combination of connected graphs. An important property of Kontsevich integral is that  $W(\mathcal{L})$  is narrow.

As we have mentioned,  $W(\mathcal{L})$  depends on a presentation of  $\mathcal{L}$  as a dotted Morse link. However, the strut part of  $W(\mathcal{L})$  is topologically invariant and, in fact, is well-known to be

$$W_{\text{str}}(\mathcal{L}) = \frac{1}{2} \sum_{i,j=1}^L \ell_{ij} \quad i \text{ --- } j, \quad (2.7)$$

where  $\ell_{ij}$  are the linking numbers of  $\mathcal{L}$ .

Let  $\mathfrak{G}$  be a compact Lie group,  $\mathfrak{g}$  be its Lie algebra and  $\mathfrak{h} \subset \mathfrak{g}$  its Cartan subalgebra. We assume that  $\mathfrak{g}$  is equipped with a positive-definite Killing form, so we will not make a consistent effort to distinguish between  $\mathfrak{g}$ ,  $\mathfrak{h}$  and their dual spaces  $\mathfrak{g}^*$ ,  $\mathfrak{h}^*$ . Let  $\vec{\alpha} \in \mathfrak{h}^*$  be the highest weights (shifted by a half-sum of positive roots) of  $\mathfrak{G}$ -modules  $V_{\vec{\alpha}_1}, \dots, V_{\vec{\alpha}_L}$  assigned to the components of  $\mathcal{L}$ . We denote by  $J_{\vec{\alpha}}(\mathcal{L}; q)$  the colored Jones polynomial of  $\mathcal{L}$  which corresponds to this data. It is well-known that it has an expansion in powers of  $\hbar = \log q$

$$J_{\vec{\alpha}}(\mathcal{L}; q) = \left( \prod_{j=1}^L \dim V_{\vec{\alpha}_j} \right) \sum_{n=0}^{\infty} p_n(\mathcal{L}; \vec{\alpha}) \hbar^n, \quad (2.8)$$

where the link invariants  $p_n(\mathcal{L}; \vec{\alpha})$  are Weyl-invariant polynomials of  $\vec{\alpha}$ . This expansion can be derived from  $Z_{\mathcal{B}}(\mathcal{L})$ . The standard method is to step back to  $Z_{\mathcal{A}}(\mathcal{L})$  through eq.(2.5) and then apply the  $\mathfrak{g}$  weight system. The weight system is applied in two steps:

$$\mathcal{A}(\uparrow_{\mathcal{L}}) \xrightarrow{\mathcal{T}_{\mathfrak{g}}^{1,\hbar}} ((\mathbf{U}\mathfrak{g})^{\otimes L})^{\mathfrak{G}}[[\hbar]] \xrightarrow{\text{Tr}_{V_{\vec{\alpha}}}} \mathbb{C}[[\hbar]] \quad (2.9)$$

The map  $\mathcal{T}_{\mathfrak{g}}^{1,\hbar}$  is a combination of two maps. First, a graph  $D \in \mathcal{A}(\uparrow_{\mathcal{L}})$  is multiplied by  $\hbar^{\deg_1 D}$ . Then the map  $\mathcal{A}(\uparrow_{\mathcal{L}}) \xrightarrow{\mathcal{T}_{\mathfrak{g}}} ((\mathbf{U}\mathfrak{g})^{\otimes L})^{\mathfrak{G}}$  is applied. Let us recall the definition [1] of two similar maps (which we denote by the same symbol)

$$\mathcal{A}(\uparrow_{\mathcal{L}}) \xrightarrow{\mathcal{T}_{\mathfrak{g}}} ((\mathbf{U}\mathfrak{g})^{\otimes |\mathcal{L}|})^{\mathfrak{G}}, \quad \mathcal{B}_{\mathcal{L}} \xrightarrow{\mathcal{T}_{\mathfrak{g}}} ((S\mathfrak{g})^{\otimes |\mathcal{L}|})^{\mathfrak{G}}, \quad (2.10)$$

where  $|\mathfrak{L}|$  denotes the number of elements in  $\mathfrak{L}$ . Let  $f_{\mathfrak{g}} \in \bigwedge^3 \mathfrak{g}^*$  be the structure constant tensor of  $\mathfrak{g}$  and let  $h_{\mathfrak{g}} \in S^2 \mathfrak{g}^*$  be the positive definite Killing metric on  $\mathfrak{g}$  normalized in such a way that short roots have length  $\sqrt{2}$ . We denote by  $h_{\mathfrak{g}}^{-1} \in S^2 \mathfrak{g}$  its inverse matrix. Now let  $D$  be a (1,3)-valent graph with  $n_1(D)$  1-valent vertices,  $n_3(D)$  3-valent vertices and  $n_e(D)$  edges. Assume for a moment that all 1-valent vertices of  $D$  have distinct labels. Associate a space  $\bigwedge^3 \mathfrak{g}^*$  with each 3-vertex of  $D$  and a space  $S^2 \mathfrak{g}$  with each edge of  $D$ . Consider their tensor products  $(\bigwedge^3 \mathfrak{g}^*)^{\otimes n_3(D)}$  and  $(S^2 \mathfrak{g})^{\otimes n_e(D)}$ . There is a natural ‘index contraction’ map

$$(\bigwedge^3 \mathfrak{g}^*)^{\otimes n_3(D)} \otimes (S^2 \mathfrak{g})^{\otimes n_e(D)} \xrightarrow{\mathcal{C}_D} \mathfrak{g}^{\otimes n_1(D)}, \quad (2.11)$$

which pairs up the spaces  $\mathfrak{g}$  and  $\mathfrak{g}^*$  from incident edges and 3-valent vertices, so that the spaces  $\mathfrak{g}$  of edges which are incident to 1-valent vertices remain uncontracted. Then we define  $\tilde{\mathcal{T}}_{\mathfrak{g}}(D)$  as

$$f_{\mathfrak{g}}^{\otimes n_3(D)} \otimes (h_{\mathfrak{g}}^{-1})^{\otimes n_e(D)} \xrightarrow{\mathcal{C}_D} \tilde{\mathcal{T}}_{\mathfrak{g}}(D) \in \mathfrak{g}^{\otimes n_1(D)}. \quad (2.12)$$

Now, as we know, if the graph  $D$  is an element of  $\mathcal{A}(\uparrow \mathfrak{L})$ , then the STU relations project  $\tilde{\mathcal{T}}_{\mathfrak{g}}(D)$  naturally to  $\mathcal{T}_{\mathfrak{g}}(D) \in ((U\mathfrak{g})^{\otimes |\mathfrak{L}|})^{\mathfrak{g}}$ , whereas if  $D$  is an element of  $\mathcal{B}_{\mathfrak{L}}$ , then the equivalences between 1-vertices project  $\tilde{\mathcal{T}}_{\mathfrak{g}}(D)$  to  $\mathcal{T}_{\mathfrak{g}}(D) \in ((S\mathfrak{g})^{\otimes |\mathfrak{L}|})^{\mathfrak{g}}$ .

The map  $\text{Tr}_{V_{\vec{\alpha}}}$  of (2.9) evaluates the traces of the elements of  $U\mathfrak{g}$  in  $\mathfrak{g}$ -modules  $V_{\vec{\alpha}_1}, \dots, V_{\vec{\alpha}_L}$  assigned to the link components. Finally,

$$J_{\vec{\alpha}}(\mathcal{L}; q) = \text{Tr}_{V_{\vec{\alpha}}} \mathcal{T}_{\mathfrak{g}}^{1, \hbar}(Z_{\mathcal{A}}(\mathcal{L})). \quad (2.13)$$

**Remark 2.1** Since there is a natural inclusion  $\mathfrak{g} \hookrightarrow \mathfrak{g}_{\mathbb{C}}$ , then we may consider  $f_{\mathfrak{g}}$  and  $h_{\mathfrak{g}}^{-1}$  as elements of  $\bigwedge^3 \mathfrak{g}_{\mathbb{C}}^*$  and  $S^2 \mathfrak{g}_{\mathbb{C}}$ . Obviously, both maps (2.9) commute with complexification. The complexified version of (2.9) will be more convenient later when we split  $\mathfrak{g}$  into a Cartan subalgebra  $\mathfrak{h}$  and the space of roots  $\mathfrak{r}$ , because the adjoint action of  $\mathfrak{h}$  on  $\mathfrak{r}$  could be diagonalized over  $\mathbb{C}$ .

## 2.2 Wheeling, Kirillov integral and Reshetikhin formula

An alternative way of deriving the expansion (2.8) from Kontsevich integral is based on Kirillov’s integral formula for the trace map  $\text{Tr}_{V_{\vec{\alpha}}}$ . Let  $S\mathfrak{g} \xrightarrow{\beta_{\mathfrak{g}}} U\mathfrak{g}$  be the Poincare-Birkhoff-Witt symmetrization isomorphism and let  $D(j_{\mathfrak{g}}^{1/2}): S\mathfrak{g} \rightarrow S\mathfrak{g}$  be the Duflo isomorphism (see [2] and references therein). For an element  $\vec{\alpha} \in \mathfrak{h}^*$  let  $\mathcal{O}_{\vec{\alpha}} \subset \mathfrak{g}^*$  be its coadjoint orbit

equipped with the integration measure derived from the Kirillov-Kostant symplectic form. Then the following square is commutative

$$\begin{array}{ccc}
\mathbf{S}\mathfrak{g} & \xrightarrow{\beta_{\mathfrak{g}}} & \mathbf{U}\mathfrak{g} \\
\downarrow D(j_{\mathfrak{g}}^{1/2}) & & \downarrow \text{Tr}_{V_{\vec{\alpha}}} \\
\mathbf{S}\mathfrak{g} & \xrightarrow{\int_{\mathcal{O}_{\vec{\alpha}}}} & \mathbb{C}
\end{array} \tag{2.14}$$

Here the map  $\int_{\mathcal{O}_{\vec{\alpha}}} : \mathbf{S}\mathfrak{g} \rightarrow \mathbb{C}$  means the following: the space  $\mathbf{S}\mathfrak{g}$  is naturally isomorphic to the space of polynomial functions on  $\mathfrak{g}^*$  and these polynomials can be integrated over  $\mathcal{O}_{\vec{\alpha}}$ .

According to [2], Duflo isomorphism can be performed at the graph level as a wheeling map  $\hat{\Omega}$ . Here is the definition of  $\hat{\Omega}_j$  acting on the legs of the graphs of  $\mathcal{B}_{\mathcal{L}}$  assigned to the link component  $\mathcal{L}_j$ . We define the ‘modified Bernoulli numbers’  $b_{2n}$  by the expansion

$$\frac{1}{2} \log \left( \frac{\sinh(x/2)}{(x/2)} \right) = \sum_{n=1}^{\infty} b_{2n} x^{2n}. \tag{2.15}$$

Then we define  $\Omega \in \mathcal{B}$  as the exponential

$$\Omega = \exp \left( - \sum_{n=1}^{\infty} b_{2n} \omega_{2n} \right), \tag{2.16}$$

where  $\omega_{2n}$  is a circle with  $2n$  radial edges attached to it<sup>3</sup>. Following the notations of [3], we label the 1-valent vertices of the graphs of  $\mathcal{B}_{\mathcal{L}}$  with link components by assigning to them formal variables  $\mathbf{x}$ . We also assign the dual variable  $\partial_{x_j}$  to all the 1-valent vertices of  $\Omega$  thus producing  $\Omega_j = \Omega(\partial_{x_j})$ . Now for any  $F \in \mathcal{B}_{\mathcal{L}}$

$$\hat{\Omega}_j F = \Omega_j \mathfrak{b} F, \tag{2.17}$$

where the operation  $\mathfrak{b}$  was defined in [3] as the ‘application of derivatives’  $\partial_{x_j}$  of  $\Omega_j$  to the variables  $x_j$  sitting at some 1-valent vertices of  $F$ . This application of the derivatives means that the corresponding edges of  $\Omega_j$  and  $F$  are joined into single edges:

$$\begin{array}{c} \text{---} \bullet \text{---} \end{array} \mathfrak{b}_{\partial_{x_j}} \begin{array}{c} \text{---} \bullet \text{---} \\ \text{---} \bullet \text{---} \end{array} \begin{array}{c} | \\ | \end{array} = \begin{array}{c} \text{---} \bullet \text{---} \bullet \text{---} \\ | \quad | \end{array} \tag{2.18}$$

Denote by  $\hat{\Omega}_{\mathcal{L}}$  the composition of the wheeling maps  $\hat{\Omega}_j$  for all the link components of  $\mathcal{L}$ . This map allows us to extend a commutative square (2.14) to a commutative diagram (*cf.*

---

<sup>3</sup>In contrast to [2], we placed an extra minus sign in the exponent of the formula for  $\Omega$ . Thus our ‘wheeling’ is the inverse of the wheeling of [2].

the monster diagram of [2])

$$\begin{array}{ccccc}
& & \mathcal{A}(\uparrow_{\mathcal{L}}) & & \\
& \nearrow \hat{\chi} & & \nwarrow \mathcal{T}_{\mathfrak{g}} & \\
\mathcal{B}_{\mathcal{L}} & \xrightarrow{\mathcal{T}_{\mathfrak{g}}} & ((\mathbf{S}\mathfrak{g})^{\otimes L})^{\mathfrak{G}} & \xrightarrow{\beta_{\mathfrak{g}}} & ((\mathbf{U}\mathfrak{g})^{\otimes L})^{\mathfrak{G}} \\
\downarrow \hat{\Omega}_{\mathcal{L}} & & \downarrow D_{\mathcal{L}}(j_{\mathfrak{g}}^{1/2}) & & \downarrow \text{Tr}_{V_{\vec{\alpha}}} \\
\mathcal{B}_{\mathcal{L}} & \xrightarrow{\mathcal{T}_{\mathfrak{g}}} & ((\mathbf{S}\mathfrak{g})^{\otimes L})^{\mathfrak{G}} & \xrightarrow{\int_{\mathcal{O}_{\vec{\alpha}}}} & \mathbb{C}
\end{array} \tag{2.19}$$

Since the map (2.4) is the graph algebra counterpart of the PBW symmetrization map  $\beta_{\mathfrak{g}}$ , then the upper triangle of (2.19) is commutative. The maps  $\mathcal{T}_{\mathfrak{g}}: \mathcal{B}_{\mathcal{L}} \rightarrow ((\mathbf{S}\mathfrak{g})^{\otimes L})^{\mathfrak{G}}$  convert (1,3)-valent graphs into the elements of  $((\mathbf{S}\mathfrak{g})^{\otimes L})^{\mathfrak{G}}$  by placing the structure constants of  $\mathfrak{g}$  at 3-valent vertices and contracting indices along the internal edges, while  $D_{\mathcal{L}}(j_{\mathfrak{g}}^{1/2})$  is the Dulfo isomorphism applied to all components of the tensor product  $((\mathbf{S}\mathfrak{g})^{\otimes L})^{\mathfrak{G}}$ . We also used a shortcut notation  $\mathcal{O}_{\vec{\alpha}}$  for  $\mathcal{O}_{\vec{\alpha}_1} \times \cdots \times \mathcal{O}_{\vec{\alpha}_L}$ .

The  $\mathfrak{G}$ -invariance of the elements of  $((\mathbf{S}\mathfrak{g})^{\otimes L})^{\mathfrak{G}}$  means that we can omit one of the integrals in  $\int_{\mathcal{O}_{\vec{\alpha}}}$ , replacing it by the volume of the coadjoint orbit. Since  $\text{Vol } \mathcal{O}_{\vec{\alpha}} = \dim V_{\vec{\alpha}}$ , then in the context of the diagram (2.19)

$$\int_{\mathcal{O}_{\vec{\alpha}}} = \dim V_{\vec{\alpha}_k} \text{ev}_{k;\vec{\alpha}_k} \int_{\mathcal{O}_{\vec{\alpha}_{(k)}}}, \quad 1 \leq k \leq L, \tag{2.20}$$

where  $\text{ev}_{k;\vec{\alpha}}: ((\mathbf{S}\mathfrak{g})^{\otimes L})^{\mathfrak{G}} \rightarrow (\mathbf{S}\mathfrak{g})^{\otimes(L-1)}$  evaluates the elements of the  $k$ -th component of  $((\mathbf{S}\mathfrak{g})^{\otimes L})^{\mathfrak{G}}$  (considered as polynomials on  $\mathfrak{h}^*$ ) on  $\vec{\alpha}$ .

It is easy to see that the wheeling map  $\hat{\Omega}$  does not change the degree (2.2) of graphs, therefore we can replace  $\mathcal{T}_{\mathfrak{g}}$  by  $\mathcal{T}_{\mathfrak{g}}^{1,\hbar}$  and add  $[[\hbar]]$  to the spaces  $((\mathbf{S}\mathfrak{g})^{\otimes L})^{\mathfrak{G}}$ ,  $((\mathbf{U}\mathfrak{g})^{\otimes L})^{\mathfrak{G}}$  and  $\mathbb{C}$  in the diagram (2.19) without spoiling its commutativity.

A combination of equations (2.5) and (2.13) demonstrates that

$$J_{\vec{\alpha}}(\mathcal{L}; q) = \text{Tr}_{V_{\vec{\alpha}}} \circ \mathcal{T}_{\mathfrak{g}}^{1,\hbar} \circ \hat{\chi}(Z_B(\mathcal{L})). \tag{2.21}$$

At the same time, the  $[[\hbar]]$  version of the diagram (2.19) shows that

$$\text{Tr}_{V_{\vec{\alpha}}} \circ \mathcal{T}_{\mathfrak{g}}^{1,\hbar} \circ \hat{\chi} = \int_{\mathcal{O}_{\vec{\alpha}}} \mathcal{T}_{\mathfrak{g}}^{1,\hbar} \circ \hat{\Omega}_{\mathcal{L}}. \tag{2.22}$$

Thus we obtain the alternative version of eq.(2.13)

$$J_{\vec{\alpha}}(\mathcal{L}; q) = \int_{\mathcal{O}_{\vec{\alpha}}} \mathcal{T}_{\mathfrak{g}}^{1,\hbar} \circ \hat{\Omega}_{\mathcal{L}}(Z_B(\mathcal{L})). \tag{2.23}$$

Let us look more closely at the wheeled Kontsevich integral

$$Z_B^\Omega(\mathcal{L}) = \hat{\Omega}_\mathcal{L}(Z_B(\mathcal{L})). \quad (2.24)$$

According to [3], the operation  $\flat$  between two exponentials of connected graphs is itself an exponential of connected graphs, whose structure is described in Exercise 2.4 of [3] (one just has to expand both exponentials, apply  $\flat$  and then pick only connected graphs – they will make up the exponent of the resulting exponential). Therefore

$$Z_B^\Omega(\mathcal{L}) = \exp(W^\Omega(\mathcal{L})) \quad (2.25)$$

and  $W^\Omega(\mathcal{L}) \in \mathcal{B}_\mathcal{L}$  is narrow. The strut part of  $W^\Omega(\mathcal{L})$  is the same as that of  $W(\mathcal{L})$ :

$$W_{\text{str}}^\Omega(\mathcal{L}) = \frac{1}{2} \sum_{i,j=1}^L \ell_{ij} \quad i \text{ --- } j, \quad (2.26)$$

The maps  $\mathcal{T}_\mathfrak{g}$  and  $\mathcal{T}_\mathfrak{g}^{1,\hbar}$  are algebra homomorphisms, so

$$\mathcal{T}_\mathfrak{g}^{1,\hbar}(Z_B^\Omega(\mathcal{L})) = \exp(\mathcal{T}_\mathfrak{g}^{1,\hbar}(W^\Omega(\mathcal{L}))), \quad (2.27)$$

and we proved the following

**Theorem 2.2 (Reshetikhin's formula)** *The expansion of the colored Jones-HOMFLY polynomial of a link  $\mathcal{L} \subset S^3$  in powers of  $\hbar = \log q$  can be presented as Kirillov's integral over the coadjoint orbits of the (shifted) highest weights of  $\mathfrak{G}$ -modules assigned to the link components*

$$J_{\vec{\alpha}}(\mathcal{L}; q) = \int_{\mathcal{O}_{\vec{\alpha}}} \exp(\mathcal{T}_\mathfrak{g}^{1,\hbar}(W^\Omega(\mathcal{L}))). \quad (2.28)$$

Let us write eq.(2.28) more explicitly. First, we split  $W^\Omega(\mathcal{L})$  into a sum

$$W^\Omega(\mathcal{L}) = \sum_{m,n=0}^{\infty} W_{m,n}^\Omega(\mathcal{L}), \quad (2.29)$$

where  $W_{m,n}^\Omega(\mathcal{L})$  are linear combinations of graphs  $D$  such that  $\#(1\text{-vertices}) = m$  and  $\chi(D) = n - 1$  so that according to eq.(2.2),

$$\deg_1 W_{m,n}^\Omega(\mathcal{L}) = m + n - 1. \quad (2.30)$$

Then  $\mathcal{T}_\mathfrak{g}(W_{m,n}^\Omega(\mathcal{L}))$  is an element of  $((\mathbf{Sg})^{\otimes L})^\mathfrak{G}$  of total degree  $m$  and  $\mathcal{T}_\mathfrak{g}^{1,\hbar}(W_{m,n}^\Omega(\mathcal{L}))$  is proportional to  $\hbar^{m+n-1}$ . Since  $\mathbf{Sg}$  is canonically isomorphic to the algebra of polynomials on  $\mathfrak{g}^*$ ,

we can present  $\mathcal{T}_{\mathfrak{g}}(W_{m,n}^{\Omega}(\mathcal{L}))$  as a  $\mathfrak{G}$ -invariant homogeneous polynomial  $L_{m,n}(\mathcal{L}; \vec{\chi})$  of degree  $m$ , where  $\vec{\chi} \in \mathfrak{g}^*$ . Then in view of eq.(2.30),

$$\mathcal{T}_{\mathfrak{g}}^{1,\hbar} W^{\Omega}(\mathcal{L}) = \sum_{m,n=0}^{\infty} L_{m,n}(\mathcal{L}; \vec{\chi}) \hbar^{m+n-1}, \quad (2.31)$$

and we can rewrite eq.(2.28) as

$$J_{\vec{\alpha}}(\mathcal{L}; q) = \int_{\mathcal{O}_{\vec{\alpha}}} d\vec{\chi} \exp \left( \sum_{m,n=0}^{\infty} L_{m,n}(\mathcal{L}; \vec{\chi}) \hbar^{m+n-1} \right). \quad (2.32)$$

Since all graphs with a single leg are equal to zero in  $\mathcal{B}_{\mathcal{L}}$ , we conclude that

$$L_{m,1}(\mathcal{L}; \vec{\chi}) = 0. \quad (2.33)$$

Also, since a tree graph has at least 2 legs,  $L_{0,0}(\mathcal{L}; \vec{\chi}) = 0$ . We deduce from eq.(2.26) that

$$L_{2,0}(\mathcal{L}; \vec{\chi}) = \frac{1}{2} \sum_{i,j=1}^L \ell_{ij} (\vec{\chi}_i, \vec{\chi}_j), \quad (2.34)$$

Finally, since  $\mathcal{B}_{\mathcal{L}}$  does not contain a  $\chi(D) = 0$  graph without legs, then  $L_{m,1}(\mathcal{L}; \vec{\chi}) = 0$ . Since  $L_{m,0}(\mathcal{L}; \vec{\chi}) = L_{m,1}(\mathcal{L}; \vec{\chi}) = 0$  for  $m \leq 1$ , then the exponent of the *r.h.s.* of eq.(2.32) contains only strictly positive powers of  $\hbar$ . If we expand the exponential in Taylor series and then integrate the coefficients at each power of  $\hbar$ , then we will reproduce the Melvin-Morton expansion (2.8).

## 2.3 Invariant measure on a coadjoint orbit

Let us perform a substitution (2.20) and change the integration variables in eq.(2.32) from  $\vec{\chi}_j$  to  $\vec{X}_j = \hbar \vec{\chi}_j$ ,  $1 \leq j \leq L$ . Since the polynomials  $L_{m,n}(\mathcal{L}; \vec{\chi})$  are homogeneous of degree  $m$ , then eq.(2.32) becomes

$$J_{\vec{\alpha}}(\mathcal{L}; q) = \hbar^{-(L-1)|\Delta_+|} \dim V_{\vec{\alpha}_k} \int_{\mathcal{O}_{\hbar \vec{\alpha}(k)}} d\vec{X}_{(k)} \exp \left( \sum_{m,n=0}^{\infty} L_{m,n}(\mathcal{L}; \vec{X}) \hbar^{n-1}, \right) \quad (2.35)$$

where  $|\Delta_+| = (1/2) \dim \mathcal{O}_{\vec{\alpha}}$  is the number of positive roots of  $\mathfrak{g}$ . Now for the variables  $\vec{a} \in \mathfrak{h}^*$  let us consider a formal expression

$$J(\mathcal{L}; \vec{a}; \hbar, k) = \hbar^{-L|\Delta_+|} \frac{d_{\mathfrak{g}}(\vec{a}_k)}{d_{\mathfrak{g}}(\vec{\rho})} \int_{\mathcal{O}_{\vec{a}(k)}} d\vec{X}_{(k)} \exp \left( \frac{1}{\hbar} \sum_{m,n=0}^{\infty} L_{m,n}(\mathcal{L}; \vec{X}) \Big|_{\vec{X}_k = \vec{a}_k} \hbar^n \right), \quad (2.36)$$

where

$$d_{\mathfrak{g}}(\vec{a}) = \prod_{\vec{\lambda} \in \Delta_+} (\vec{a}, \vec{\lambda}), \quad (2.37)$$

so that  $d_{\mathfrak{g}}(\vec{\alpha})/d_{\mathfrak{g}}(\vec{\rho}) = \dim V_{\vec{\alpha}}$ . Expression (2.36) does not make sense in itself, because the coefficients at the powers of  $\hbar$  in the exponent are formal power series in  $\vec{\mathbf{X}}$ , and we can not make sense of the integral. We can only make sense of the expression  $J(\mathcal{L}; \hbar \vec{\alpha}; \hbar, k)$ , because in view of eq.(2.35) it can be defined as a power series in  $\hbar$  and

$$J(\mathcal{L}; \hbar \vec{\alpha}; \hbar, k) = J_{\vec{\alpha}}(\mathcal{L}; q). \quad (2.38)$$

Despite the lack of proper definition of the whole integral (2.36), its integrand looks suitable for the stationary phase approximation in the limit of

$$\hbar \longrightarrow 0, \quad (q \longrightarrow 1), \quad \vec{a}_j (= \hbar \vec{\alpha}_j) = \text{const}, \quad 1 \leq j \leq L. \quad (2.39)$$

Indeed, we may assume that a formal parameter  $\hbar$  is purely imaginary; actually, this is the case in the Quantum Field Theory approach to quantum invariants of 3d topology. The advantage of the stationary phase approximation for us is that within its calculations the ‘honest’ integrals are essentially replaced by simple formal combinatorial manipulations with the integrand. This combinatorics is called ‘Feynman rules’ in Quantum Field Theory.

The first step in the stationary phase approximation is to identify the stationary points of  $\sum_{m=2}^{\infty} L_{m,0}(\mathcal{L}; \vec{\mathbf{X}}) \Big|_{\vec{X}_k = \vec{a}_k}$  on  $\mathcal{O}_{\vec{\mathbf{a}}(k)}$ . Generally, this may be tricky, since  $\sum_{m=2}^{\infty} L_{m,0}(\mathcal{L}; \vec{\mathbf{X}})$  is only a formal power series: one might try to find the stationary points of its lowest terms and then correct them perturbatively by taking into account the higher order terms. Luckily, there exists a point

$$\vec{\mathbf{X}} = \vec{\mathbf{a}} \quad (2.40)$$

(that is, when all  $\vec{\mathbf{X}}$  belong to the same Cartan subalgebra) which is manifestly stationary (we will check this a bit later). We are going to calculate (or, more precisely, define) its contribution to the integral (2.36) by following the formal rules of the stationary phase approximation. We will call the resulting power series in  $\hbar$  ‘the  $\mathfrak{g}$ -based  $U(1)$ -RCC invariant’.

Let us find a convenient parametrization of  $\mathcal{O}_{\vec{\mathbf{a}}(k)}$  around the point (2.40). A map

$$g\mathfrak{T} \mapsto \text{Ad}_{g\mathfrak{T}} \vec{a} \quad (2.41)$$

establishes an isomorphism between a coadjoint orbit  $\mathcal{O}_{\vec{a}}$  and the quotient  $\mathfrak{G}/\mathfrak{T}$ . A composition of maps

$$\mathfrak{r} \xrightarrow{\text{exp}} \mathfrak{G} \longrightarrow \mathfrak{G}/\mathfrak{T}, \quad (2.42)$$



identifies the vicinity of the origin in the space of roots  $\mathfrak{r} \subset \mathfrak{g}$  with the vicinity of the identity at  $\mathfrak{G}/\mathfrak{T}$ . Thus a composition of the maps (2.41) and (2.42) makes  $\mathfrak{r}$  a coordinate space for a vicinity of  $\vec{a}$  in  $\mathcal{O}_{\vec{a}}$ :

$$\vec{X} = R_{\vec{a}}(\vec{x}), \quad \text{where} \quad R_{\vec{a}}(\vec{x}) = \text{Ad}_{e^{\vec{x}}} \vec{a} = \sum_{m=0}^{\infty} \frac{\text{ad}_{\vec{x}}^m \vec{a}}{m!}. \quad (2.43)$$

Let us find the measure  $\mu_{\vec{a}}(\vec{x})$  on  $\mathfrak{r}$  which corresponds to the Kirillov-Kostant measure on  $\mathcal{O}_{\vec{a}}$ . It is easy to find the measure at the origin:

$$\mu_{\vec{a}}(\vec{0}) = (-2\pi)^{-|\Delta_+|} d_{\mathfrak{g}}(\vec{a}). \quad (2.44)$$

(the minus sign compensates for the fact that the metric on  $\mathfrak{g}$  is negative-definite).

We are going to use the invariance of the Kirillov-Kostant measure under the adjoint action of the elements  $e^{\vec{y}}$  ( $\vec{y} \in \mathfrak{r}$ ) on  $\mathcal{O}_{\vec{a}}$  in order to find  $\mu_{\vec{a}}(\vec{x})$  at a general point  $\vec{x}$ . Let us rewrite this action in terms of  $\mathfrak{r}$  coordinates. We define a (non-linear) map  $F_{\vec{y}}: \mathfrak{r} \rightarrow \mathfrak{r}$  by a commutative diagram

$$\begin{array}{ccc} \mathfrak{r} & \xrightarrow{F_{\vec{y}}} & \mathfrak{r} \\ \downarrow R_{\vec{a}} & & \downarrow R_{\vec{a}} \\ \mathcal{O}_{\vec{a}} & \xrightarrow{\text{Ad}_{e^{\vec{y}}}} & \mathcal{O}_{\vec{a}} \end{array} \quad (2.45)$$

where the map  $R_{\vec{a}}: \mathfrak{r} \rightarrow \mathcal{O}_{\vec{a}}$  is defined by eq.(2.43). This diagram implies a defining relation for  $F_{\vec{y}}(\vec{x})$  and its companion  $\tilde{F}_{\vec{y}}(\vec{x})$

$$e^{\vec{y}} e^{\vec{x}} = e^{F_{\vec{y}}(\vec{x})} e^{\tilde{F}_{\vec{y}}(\vec{x})}, \quad \tilde{F}_{\vec{y}}(\vec{x}) \in \mathfrak{h}, \quad \vec{y}, \vec{x}, F_{\vec{y}}(\vec{x}) \in \mathfrak{r}. \quad (2.46)$$

Since the integration measure  $\mu_{\vec{a}}(\vec{x})$  has to be  $F_{\vec{y}}$ -invariant, it must be of the ‘left-invariant’ form

$$\mu_{\vec{a}}(\vec{x}) = \mu_{\vec{a}}(\vec{0}) / \det F'_{\vec{x}}(\vec{0}), \quad (2.47)$$

where  $F'_{\vec{y}}(\vec{x}) = \partial F_{\vec{y}}(\vec{x}) / \partial \vec{x}$  is the derivative of the map  $F_{\vec{y}}(\vec{x})$ . Since  $T_{\vec{x}}\mathfrak{r} \cong \mathfrak{r}$ , we assume that  $F'_{\vec{y}}(\vec{x}): \mathfrak{r} \rightarrow \mathfrak{r}$ .

**Theorem 2.3** *The measure (2.47) is F-invariant.*

*Quick proof of Theorem 2.3.* The F-invariant measure exists: it is a pull-back of the Kirillov-Kostant measure by the map  $R_{\vec{a}}$  of (2.43). Also it is unique (up to a constant factor), since it must satisfy eq.(2.47). Therefore, the measure (2.47) is F-invariant.  $\square$

This proof is based on the uniqueness of the conjugation-invariant measure on  $\mathcal{O}_{\vec{a}}$ . Since our ultimate goal is to strip away Lie algebras from our calculations, we will present another proof, which is essentially based on the combinatorics of commutators.

*Combinatorial proof of Theorem 2.3.* Since  $\exp(\mathfrak{r}) \subset \mathfrak{G}$  does not form a subgroup, then the invariance proof is a bit more subtle than the standard proof of the left-invariance of measures like (2.47) on the whole group  $\mathfrak{G}$ . We have to learn how to compose  $F_{\vec{y}_1}$  and  $F_{\vec{y}_2}$ . A relation

$$e^{\vec{y}_2} e^{\vec{y}_1} = e^{F_{\vec{y}_2}(\vec{y}_1)} e^{\tilde{F}_{\vec{y}_2}(\vec{y}_1)} \quad (2.48)$$

implies that

$$F_{\vec{y}_2} \circ F_{\vec{y}_1} = F_{F_{\vec{y}_2}(\vec{y}_1)} \circ \text{Ad}_{\exp(\tilde{F}_{\vec{y}_2}(\vec{y}_1))}. \quad (2.49)$$

Since  $\text{Ad}_{\exp(\tilde{F}_{\vec{y}_2}(\vec{y}_1))} : \mathfrak{r} \rightarrow \mathfrak{r}$  is a linear map and  $\det_{\mathfrak{r}} \text{Ad}_{\exp(\tilde{F}_{\vec{y}_2}(\vec{y}_1))} = 1$ , we conclude that

$$\det F'_{\vec{y}_2} \det F'_{\vec{y}_1} = \det F'_{F_{\vec{y}_2}(\vec{y}_1)} \quad (2.50)$$

and in particular

$$\det F'_{\vec{y}_2}(\vec{y}_1) \det F'_{\vec{y}_1}(\vec{0}) = \det F'_{F_{\vec{y}_2}(\vec{y}_1)}(\vec{0}). \quad (2.51)$$

The latter relation means that the measure (2.47) is indeed adjoint-invariant.  $\square$

Now let us calculate the determinant in the measure (2.47). For a stationary phase calculation it is sufficient to express it as a power series in  $\vec{x}$ . Actually, it is easier to calculate the determinant of the inverse operator  $F'_{-\vec{x}}(\vec{x})$ . In order to find the action of that operator on  $\Delta\vec{x} \in \mathfrak{r}$ , we perform the following approximate calculation up to  $\mathcal{O}(\Delta\vec{x}^2)$

$$\begin{aligned} e^{-\vec{x}} e^{\vec{x} + \Delta\vec{x}} &\approx \exp\left(\frac{I - e^{-\text{ad}_{\vec{x}}}}{\text{ad}_{\vec{x}}} \Delta\vec{x}\right) \\ &\approx \exp\left(P_{\mathfrak{r}} \frac{I - e^{-\text{ad}_{\vec{x}}}}{\text{ad}_{\vec{x}}} \Delta\vec{x}\right) \exp\left(P_{\mathfrak{h}} \frac{I - e^{-\text{ad}_{\vec{x}}}}{\text{ad}_{\vec{x}}} \Delta\vec{x}\right), \end{aligned} \quad (2.52)$$

where  $I$  is the identity operator and by definition

$$\frac{I - e^{-\text{ad}_{\vec{x}}}}{\text{ad}_{\vec{x}}} = \sum_{n=0}^{\infty} \frac{(-\text{ad}_{\vec{x}})^n}{(n+1)!} = I + \sum_{n=1}^{\infty} \frac{(-\text{ad}_{\vec{x}})^n}{(n+1)!}, \quad (2.53)$$

while  $P_{\mathfrak{r}}, P_{\mathfrak{h}}$  are the orthogonal projectors of  $\mathfrak{g}$  onto  $\mathfrak{r}$  and  $\mathfrak{h}$ . Equation (2.52) means that

$$F'_{-\vec{x}}(\vec{x}) = P_{\mathfrak{r}} \frac{I - e^{-\text{ad}_{\vec{x}}}}{\text{ad}_{\vec{x}}}. \quad (2.54)$$

in the sense that the *r.h.s.* of this formula maps  $\mathfrak{r} \subset \mathfrak{g}$  into  $\mathfrak{r}$ . We see from eqs. (2.53) and (2.54) that as an operator on  $\mathfrak{r}$

$$F'_{-\vec{x}}(\vec{x}) = I + P_{\mathfrak{r}} \sum_{n=1}^{\infty} \frac{(-\text{ad}_{\vec{x}})^n}{(n+1)!}, \quad (2.55)$$

so we can calculate its determinant as a power series in  $\vec{x}$  through the formula

$$\det(I + A) = \exp \left( \text{Tr} \log(I + A) \right) = \exp \left( - \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{Tr} A^n \right) \quad (2.56)$$

Thus combining eqs. (2.44), (2.47), (2.55) and (2.56) we find that

$$\mu_{\vec{a}}(\vec{x}) = (-2\pi)^{-|\Delta_+|} d_{\mathfrak{g}}(\vec{a}) \exp \left[ - \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{Tr}_{\mathfrak{g}} \left( \sum_{m=1}^{\infty} \frac{P_{\mathfrak{r}}(-\text{ad}_{\vec{x}})^m}{(m+1)!} \right)^n \right]. \quad (2.57)$$

Note that all calculations with operators in this equation are performed in  $\mathfrak{g}$ , while the projector  $P_{\mathfrak{r}}$  essentially reduces  $\text{Tr}_{\mathfrak{g}}$  to  $\text{Tr}_{\mathfrak{r}}$ .

## 2.4 $\mathfrak{g}$ -based $U(1)$ -RCC invariant as a stationary phase integral

Having presented the integration measure for the substitution (2.43) as an exponential of the power series in  $\vec{x}$ , we proceed to substitute

$$\vec{X}_j = R_{\vec{a}_j}(\vec{x}_j) \quad (1 \leq j \leq L, j \neq k), \quad \vec{X}_k = \vec{a}_k \quad (2.58)$$

in the exponent of (2.36), expand it in powers of  $\vec{x}$  and combine this expansion with the exponent of eq.(2.57). This results in a formula

$$\begin{aligned} I_{\mathfrak{g}}^{\mathfrak{r}}(\mathcal{L}; \vec{\mathfrak{a}}; \hbar, k) &= \hbar^{-L|\Delta_+|} \frac{d_{\mathfrak{g}}(\vec{a}_k)}{d_{\mathfrak{g}}(\vec{\rho})} \\ &\times \int_{[\vec{\mathfrak{x}}_{(k)}=0]} d\vec{\mathfrak{x}}_{(k)} \left[ \exp \left( \sum_{m,n=0}^{\infty} L_{m,n}(\mathcal{L}; R_{\vec{a}_1}(\vec{x}_1), \dots, R_{\vec{a}_L}(\vec{x}_L)) \hbar^{n-1} \right) \prod_{j=1}^L \mu_{\vec{a}_j}(\vec{x}_j) \right] \Big|_{\vec{x}_k=0} \\ &= \hbar^{-|\Delta_+|} (-2\pi\hbar)^{(1-L)|\Delta_+|} \frac{d_{\mathfrak{g}}(\vec{\mathfrak{a}})}{d_{\mathfrak{g}}(\vec{\rho})} \\ &\times \int_{[\vec{\mathfrak{x}}_{(k)}=0]} d\vec{\mathfrak{x}}_{(k)} \exp \left( \frac{1}{\hbar} \sum_{m,n=0}^{\infty} L_{m,n}^{\mathfrak{R}}(\mathcal{L}; \vec{\mathfrak{a}}; \vec{\mathfrak{x}}_{(k)}) \hbar^n \right). \end{aligned} \quad (2.59)$$

Here  $d_{\mathfrak{g}}(\vec{\mathbf{a}}) = \prod_{j=1}^L d_{\mathfrak{g}}(\vec{a}_j)$ ,  $\int_{[\vec{\mathbf{x}}_{(k)}=0]}$  denotes the stationary phase contribution of the point  $\vec{\mathbf{x}}_{(k)} = \vec{0}$  to the integral, while  $L_{m,n}^{\mathbf{R}}(\mathcal{L}; \vec{\mathbf{a}}; \vec{\mathbf{x}}_{(k)})$  are homogeneous polynomials of  $\vec{\mathbf{x}}_{(k)}$  of degree  $m$  whose coefficients are formal power series of  $\vec{\mathbf{a}}$ :  $L_{m,n}^{\mathbf{R}}(\mathcal{L}; \vec{\mathbf{a}}; \vec{\mathbf{x}}_{(k)}) \in \mathbb{Q}[[\vec{\mathbf{a}}]][\vec{\mathbf{x}}_{(k)}]$ .

Let us check the structure of the terms  $L_{m,0}^{\mathbf{R}}(\mathcal{L}; \vec{\mathbf{a}}; \vec{\mathbf{x}}_{(k)})$ ,  $m \leq 2$  which are crucial for the stationary phase approximation. We will need two simple lemmas. Let  $D$  be a connected graph with  $m$  legs from  $\mathcal{B}_{\mathfrak{L}}$ , all of its legs having distinct labels, and let  $\mathcal{T}_{\mathfrak{g}}D$  be its image in  $((\mathbf{U}\mathfrak{g})^{\otimes m})^{\mathfrak{G}}$ . Since  $\mathcal{T}_{\mathfrak{g}}D$  is of degree  $m$ , then we may consider it to be an  $m$ -linear  $\mathfrak{G}$ -invariant function on  $\mathfrak{g}^*$ , which we denote as  $\mathcal{T}_{\mathfrak{g}}D(\vec{X}_1, \dots, \vec{X}_m)$ .

**Lemma 2.4** *If  $D$  is a tree graphs,  $\vec{a}_2, \dots, \vec{a}_m \in \mathfrak{h}$  and  $\vec{X} \in \mathfrak{g}$ , then  $\mathcal{T}_{\mathfrak{g}}D(\vec{X}, \vec{a}_2, \dots, \vec{a}_m) = 0$ , unless  $D$  is a strut.*

*Proof.* If  $D$  is not a strut, then out of any  $m-1$  of its  $m$  1-valent vertices one can always find a pair of 1-valent vertices connected to the same 3-valent vertex. This means that each term in the expression of  $\mathcal{T}_{\mathfrak{g}}D(\vec{X}, \vec{a}_2, \dots, \vec{a}_m)$  contains a commutator of the elements  $\vec{a}_j$  attached to those 1-valent vertices. Then  $\mathcal{T}_{\mathfrak{g}}D(\vec{X}, \vec{a}_2, \dots, \vec{a}_m) = 0$ , because all elements of  $\mathfrak{h}$  commute.  $\square$

**Lemma 2.5** *If  $\vec{a}_1, \dots, \vec{a}_m \in \mathfrak{h}$  and  $\vec{X} \in \mathfrak{g}$ , then*

$$\mathcal{T}_{\mathfrak{g}}D([\vec{X}, \vec{a}_1], \vec{a}_2, \dots, \vec{a}_m) = 0. \quad (2.60)$$

*Proof.* Since  $\mathcal{T}_{\mathfrak{g}}D(\vec{X}_1, \dots, \vec{X}_m)$  is invariant under adjoint action of  $\mathfrak{G}$  on its arguments, then

$$\mathcal{T}_{\mathfrak{g}}D([\vec{X}, \vec{a}_1], \vec{a}_2, \dots, \vec{a}_m) = - \sum_{j=2}^m \mathcal{T}_{\mathfrak{g}}D(\vec{X}, \vec{a}_2, \dots, [\vec{a}_j, \vec{a}_1], \dots, \vec{a}_m). \quad (2.61)$$

The *r.h.s.* of this equation is zero, since all  $[\vec{a}_j, \vec{a}_1] = \vec{0}$ .  $\square$

Obviously,  $L_{0,0}^{\mathbf{R}}(\mathcal{L}; \vec{\mathbf{a}}; \vec{\mathbf{x}}_{(k)}) = \sum_{m=2}^{\infty} L_{m,0}(\mathcal{L}; \vec{\mathbf{a}})$  (we dropped  $m=0, 1$  in the sum, because for them  $L_{m,0}(\mathcal{L}; \vec{\mathbf{X}}) = 0$ ). The polynomials  $L_{m,0}(\mathcal{L}; \vec{\mathbf{X}})$  come from  $\mathcal{T}_{\mathfrak{g}}$  acting on the tree graphs of  $W^{\Omega}(\mathcal{L})$ , which coincide with the tree graphs of  $W(\mathcal{L})$ , since wheeling does not produce new tree graphs. Therefore, according to Lemma 2.4, the only contribution to  $L_{0,0}^{\mathbf{R}}(\mathcal{L}; \vec{\mathbf{a}}; \vec{\mathbf{x}}_{(k)})$  comes from the strut part of  $W(\mathcal{L})$ , and in view of eq.(2.7),

$$L_{0,0}^{\mathbf{R}}(\mathcal{L}; \vec{\mathbf{a}}; \vec{\mathbf{x}}_{(k)}) = \frac{1}{2} \sum_{i,j=1}^L \ell_{ij}(\vec{a}_i, \vec{a}_j) \stackrel{\text{def}}{=} \text{lk}(\mathcal{L}, \vec{\mathbf{a}}). \quad (2.62)$$

The terms  $L_{1,n}^{\mathbf{R}}(\mathcal{L}; \vec{\mathbf{a}}; \vec{\mathbf{x}}_{(k)})$  (and in particular  $L_{1,0}^{\mathbf{R}}(\mathcal{L}; \vec{\mathbf{a}}; \vec{\mathbf{x}}_{(k)})$ ) come from all graphs of  $W^{\Omega}(\mathcal{L})$ . The formula  $R_{\vec{a}}(\vec{x}) = \vec{a} + [\vec{x}, \vec{a}] + \mathcal{O}(\vec{x}^2)$  suggests that the contribution of a graph  $D$

from  $W^\Omega(\mathcal{L})$  is calculated by placing  $[\vec{x}_j, \vec{a}_j]$  at one of its 1-valent vertices and  $\vec{a}$  at all other 1-valent vertices. Then Lemma 2.5 says that

$$L_{1,n}^R(\mathcal{L}; \vec{a}; \vec{x}_{(k)}) = 0 \quad \text{for all } n. \quad (2.63)$$

In particular, this means that there are no terms of order  $\hbar^{-1}$  in the exponent of (2.59) which are linear in  $\vec{x}_{(k)}$ . Thus we proved the following

**Lemma 2.6** *The point (2.40) is a stationary point of the exponent of (2.36).*

Now we turn to  $L_{2,0}^R(\mathcal{L}; \vec{a}; \vec{x}_{(k)})$ . Since  $\vec{X} = \vec{a} + [\vec{x}, \vec{a}] + (1/2)[\vec{x}, [\vec{x}, \vec{a}]] + \mathcal{O}(\vec{x}^3)$ , the terms of  $L_{2,0}^R(\mathcal{L}; \vec{a}; \vec{x}_{(k)})$  come from tree graphs of  $W(\mathcal{L})$  in two ways: either by placing the double commutator  $(1/2)[\vec{x}_j, [\vec{x}_j, \vec{a}_j]]$  at one of their 1-valent vertices and  $\vec{a}$  at the others, or by placing  $[\vec{x}_j, \vec{a}_j]$  at two 1-valent vertices and  $\vec{a}$  at the others. According to Lemma 2.4, only the strut graphs may contribute in the first way, so the contribution of the first way is

$$\frac{1}{4} \sum_{\substack{1 \leq i \neq j \leq L \\ i \neq k}} l_{ij}([\vec{x}_i, [\vec{x}_i, \vec{a}_i], \vec{a}_j]) = \frac{1}{4} \sum_{\substack{1 \leq i \neq j \leq L \\ i \neq k}} l_{ij}(\text{ad}_{\vec{a}_i} \text{ad}_{\vec{a}_j} \vec{x}_i, \vec{x}_i) \quad (2.64)$$

(we neglected the first way contribution of strut graphs  $j \text{ --- } j$  since it is canceled by their second way contribution). It is easy to see that the only graphs contributing to  $L_{2,0}^R(\mathcal{L}; \vec{a}; \vec{x}_{(k)})$  a term proportional to  $\vec{x}_i \vec{x}_j$  are the ones which have the ‘haircomb’ shape of Fig. 1 up to the cyclic order at 3-valent vertices (note that the graph of Fig. 1 contributes also to the terms  $\vec{x}_{j_1} \vec{x}_j$ ,  $\vec{x}_i \vec{x}_{j_m}$  and  $\vec{x}_{j_1} \vec{x}_{j_m}$ ). The contribution of the graph of Fig. 1 is




Figure 1: A haircomb graph.

$$([\vec{a}_{j_m}, [\dots [\vec{a}_{j_2}, [\vec{a}_{j_1}, [\vec{x}_i, \vec{a}_i] \dots], [\vec{x}_j, \vec{a}_j]]) = -(\text{ad}_{\vec{a}_j} \text{ad}_{\vec{a}_{j_m}} \dots \text{ad}_{\vec{a}_{j_1}} \text{ad}_{\vec{a}_i} \vec{x}_i, \vec{x}_j) \quad (2.65)$$

Thus adding up expression (2.64) and the contributions (2.65) we obtain  $L_{2,0}^R(\mathcal{L}; \vec{a}; \vec{x}_{(k)})$  as a quadratic form of  $\vec{x}_{(k)}$ .

Now we are almost ready to write a well-defined expression for the stationary phase integral (2.59). With a slight abuse of notation, let us think of  $\vec{x}_{(k)}$  as a single vector in a direct sum of  $L - 1$  spaces  $\mathfrak{r}$ :

$$\vec{x}_{(k)} = \sum_{\substack{1 \leq j \leq L \\ j \neq k}} \vec{x}_j \in \mathfrak{r}^{\oplus(L-1)}. \quad (2.66)$$

Then we can write

$$L_{2,0}^R(\mathcal{L}; \vec{\mathbf{a}}; \vec{\mathbf{x}}_{(k)}) = \frac{1}{2} \sum_{\substack{1 \leq i, j \leq N \\ i, j \neq k}} \left( \vec{x}_i, (Q_{\mathbf{g},k}(\mathcal{L}; \vec{\mathbf{a}}))_{ij} \vec{x}_j \right), \quad (2.67)$$

where  $Q_{\mathbf{g},k}(\mathcal{L}; \vec{\mathbf{a}})$  is a linear operator acting on  $\mathfrak{r}^{\oplus(L-1)}$ , whose coefficients  $(Q_{\mathbf{g},k}(\mathcal{L}; \vec{\mathbf{a}}))_{ij}$  are formal power series in  $\vec{\mathbf{a}}$ . Let us assume that  $Q_{\mathbf{g},k}(\mathcal{L}; \vec{\mathbf{a}})$  is non-degenerate, or in other words, that

$$\det Q_{\mathbf{g},k}(\mathcal{L}; \vec{\mathbf{a}}) \neq 0. \quad (2.68)$$

We denote by  $Q_{\mathbf{g},k}^{-1}(\mathcal{L}; \vec{\mathbf{a}})$  the inverse operator. Then the formula for the stationary phase integral (2.59) is

$$I_{\mathbf{g}}^r(\mathcal{L}; \vec{\mathbf{a}}; \hbar, k) = \hbar^{-|\Delta_+|} \frac{d_{\mathbf{g}}(\vec{\mathbf{a}})}{d_{\mathbf{g}}(\vec{\rho})} e^{\frac{1}{\hbar} \text{lk}(\mathcal{L}, \vec{\mathbf{a}})} \left( \det Q_{\mathbf{g},k}(\mathcal{L}; \vec{\mathbf{a}}) \right)^{-1/2} e^{L_{0,1}^R(\mathcal{L}; \vec{\mathbf{a}}; \vec{\mathbf{x}}_{(k)})} \text{FD}(\mathcal{L}; \vec{\mathbf{a}}), \quad (2.69)$$

where  $\text{lk}(\mathcal{L}, \vec{\mathbf{a}})$  was defined by eq.(2.62), while the most interesting factor  $\text{FD}(\mathcal{L}; \vec{\mathbf{a}})$  is defined by the formula

$$\begin{aligned} \text{FD}(\mathcal{L}; \vec{\mathbf{a}}) &= \\ &= \exp \frac{1}{\hbar} \left( \sum_{m=3}^{\infty} L_{m,0}^R(\mathcal{L}; \vec{\mathbf{a}}; \partial_{\vec{\mathbf{x}}_{(k)}}) + \sum_{m,n=1}^{\infty} L_{m,n}^R(\mathcal{L}; \vec{\mathbf{a}}; \partial_{\vec{\mathbf{x}}_{(k)}}) \hbar^n \right) \exp \left( -\frac{\hbar}{2} (\vec{\mathbf{x}}_{(k)}, Q_{\mathbf{g},k}^{-1}(\mathcal{L}; \vec{\mathbf{a}}) \vec{\mathbf{x}}_{(k)}) \right) \Big|_{\vec{\mathbf{x}}_{(k)}=0} \end{aligned} \quad (2.70)$$

and is known in the context of Quantum Field Theory as a sum of Feynman diagrams (hence the name FD). It is easy to see that  $\text{FD}(\mathcal{L}; \vec{\mathbf{a}})$  is a well-defined element of  $\mathcal{Q}(\mathbb{Q}[[\vec{\mathbf{a}}]])[[\hbar]]$ , where for a ring  $R$ ,  $\mathcal{Q}(R)$  denotes its field of fractions. The coefficients of  $L_{m,n}^R(\mathcal{L}; \vec{\mathbf{a}}; \vec{\mathbf{x}}_{(k)})$  and  $Q_{\mathbf{g},k}(\mathcal{L}; \vec{\mathbf{a}})$  belong to  $\mathbb{Q}[[\vec{\mathbf{a}}]]$ , so denominators in  $\text{FD}(\mathcal{L}; \vec{\mathbf{a}})$  are due exclusively to the denominators of  $Q_{\mathbf{g},k}^{-1}(\mathcal{L}; \vec{\mathbf{a}})$  which are, of course, equal to the determinant  $\det Q_{\mathbf{g},k}(\mathcal{L}; \vec{\mathbf{a}})$ . Then an easy combinatorics (which we will review in subsection 6.3) demonstrates that  $\text{FD}(\mathcal{L}; \vec{\mathbf{a}})$  can be put in a form

$$\text{FD}(\mathcal{L}; \vec{\mathbf{a}}) = \exp \left( \sum_{n=1}^{\infty} \frac{p_{Q,n}(\mathcal{L}; \vec{\mathbf{a}})}{(\det Q_{\mathbf{g},k}(\mathcal{L}; \vec{\mathbf{a}}))^{3n}} \hbar^n \right), \quad p_{Q,n}(\mathcal{L}; \vec{\mathbf{a}}) \in \mathbb{Q}[[\vec{\mathbf{a}}]]. \quad (2.71)$$

**Remark 2.7** Since the sum in the exponent of eq.(2.57) starts at  $n = 1$ , then it does not contain the terms of 0-th order in  $\vec{x}$ . Therefore, the term  $L_{0,1}^R(\mathcal{L}; \vec{\mathbf{a}}; \vec{\mathbf{x}}_{(k)})$ , which is singled out in eq.(2.69), comes exclusively from the polynomials  $L_{m,1}(\mathcal{L}; \vec{\mathbf{X}})$

$$L_{0,1}^R(\mathcal{L}; \vec{\mathbf{a}}; \vec{\mathbf{x}}_{(k)}) = \sum_{m=0}^{\infty} L_{m,1}(\mathcal{L}; \vec{\mathbf{a}}) \quad (2.72)$$

(in fact, one can show that  $L_{0,1}(\mathcal{L}; \vec{\mathbf{a}}) = L_{1,1}(\mathcal{L}; \vec{\mathbf{a}}) = 0$ , so the sum in eq.(2.72) begins at  $m = 2$ ).

In [13] we studied  $I_{\mathfrak{g}}^r(\mathcal{L}; \vec{\mathbf{a}}; \hbar, k)$  when  $\mathfrak{g} = su(2)$ . Since  $su(2)$  has a 1-dimensional Cartan subalgebra, then in that case it is convenient to introduce the variables  $\mathbf{a} = (\vec{\mathbf{a}}, \vec{\lambda})$ , where  $\vec{\lambda}$  is the single positive root of  $su(2)$ . We proved in [13] that

$$I_{su(2)}^r(\mathcal{L}; \mathbf{a}; \hbar, k) = \exp \left( \frac{1}{4\hbar} \sum_{i,j=1}^L \ell_{ij} a_i a_j \right) \check{J}^{(r)}(\mathcal{L}; e^{\mathbf{a}}; e^{\hbar} - 1), \quad (2.73)$$

where  $\check{J}^{(r)}(\mathcal{L}; t_1, \dots, t_L; \hbar)$  is defined by eq.(1.4). By comparing eqs.(2.69) and (1.4) through the relation (2.73) and taking into account that in the case of  $su(2)$   $d_{\mathfrak{g}}(\vec{\mathbf{a}}) = \mathbf{a}$  and  $d_{\mathfrak{g}}(\vec{\rho}) = 1$ , we find that

$$\left( \prod_{j=1}^L a_j \right)^{-1} \left( Q_{su(2),k}(\mathcal{L}; \mathbf{a}) \right)^{1/2} e^{-L_{0,1}^R(\mathcal{L}; \mathbf{a})} = \nabla_A(\mathcal{L}; e^{\mathbf{a}}), \quad (2.74)$$

or, in view of eq.(2.72),

$$\left( \prod_{j=1}^L a_j \right)^{-1} \left( Q_{su(2),k}(\mathcal{L}; \mathbf{a}) \right)^{1/2} \exp \left( - \sum_{m=0}^{\infty} L_{m,1}(\mathcal{L}; \mathbf{a}) \right) = \nabla_A(\mathcal{L}; e^{\mathbf{a}}). \quad (2.75)$$

Let us introduce a notation

$$F_{\nabla, \mathfrak{g}}(\mathcal{L}, \vec{\mathbf{a}}) = \prod_{\vec{\lambda} \in \Delta_+} \nabla_A(\mathcal{L}; e^{(\vec{\mathbf{a}}, \vec{\lambda})}). \quad (2.76)$$

In subsection 6.3 we will use the properties of the universal  $U(1)$ -RCC invariant in order to prove the following

**Theorem 2.8** *If  $\nabla_A(\mathcal{L}; \mathbf{t}) \neq 0$ , then the  $\mathfrak{g}$ -based  $U(1)$ -RCC invariant defined by eqs. (2.69) and (2.70) is well-defined and does not depend on a presentation of  $\mathcal{L}$  as a dotted Morse link and on a choice of  $k$  ( $1 \leq k \leq L$ ) thus being a topological invariant of an oriented link  $\mathcal{L}$ . It can be presented in a form*

$$I_{\mathfrak{g}}^r(\mathcal{L}; \vec{\mathbf{a}}; \hbar) = \frac{\hbar^{-|\Delta_+|}}{d_{\mathfrak{g}}(\vec{\rho})} \frac{e^{\frac{1}{\hbar} \text{lk}(\mathcal{L}, \vec{\mathbf{a}})}}{F_{\nabla, \mathfrak{g}}(\mathcal{L}, \vec{\mathbf{a}})} \exp \left( \sum_{n=1}^{\infty} \frac{p_n(\mathcal{L}; \vec{\mathbf{a}})}{[d_{\mathfrak{g}}(\vec{\mathbf{a}}) F_{\nabla, \mathfrak{g}}(\mathcal{L}, \vec{\mathbf{a}})]^{3n}} \hbar^n \right), \quad (2.77)$$

where the series

$$p_n(\mathcal{L}; \vec{\mathbf{a}}) \in \mathbb{Q}[[\vec{\mathbf{a}}]] \quad (2.78)$$

are invariants of oriented links.

In subsection 6.4 we will formulate the rationality conjecture for links and derive from it the following corollary which sharpens the formula (2.77).

**Corollary 2.9 (corollary of the rationality conjecture)** *The  $\mathfrak{g}$ -based  $U(1)$ -RCC invariant can be presented in a form*

$$I_{\mathfrak{g}}^r(\mathcal{L}; \vec{\mathbf{a}}; \hbar) = \frac{\hbar^{-|\Delta_+|}}{d_{\mathfrak{g}}(\vec{\rho})} \frac{e^{\frac{1}{\hbar} \text{lk}(\mathcal{L}, \vec{\mathbf{a}})}}{F_{\nabla, \mathfrak{g}}(\mathcal{L}, \vec{\mathbf{a}})} \exp \left( \sum_{n=1}^{\infty} \frac{P_n \left( \mathcal{L}; (e^{(\vec{a}_i, \vec{\lambda}_j)})_{\substack{1 \leq i \leq L \\ 1 \leq j \leq r}} \right)}{F_{\nabla, \mathfrak{g}}^{3n}(\mathcal{L}, \vec{\mathbf{a}})} \hbar^n \right), \quad (2.79)$$

where  $\vec{\lambda}_1, \dots, \vec{\lambda}_r$  are simple roots of  $\mathfrak{g}$  and

$$P_n \left( \mathcal{L}; (t_{ij})_{\substack{1 \leq i \leq L \\ 1 \leq j \leq r}} \right) \in \mathbb{Q} \left[ (t_{ij}^{\pm 1})_{\substack{1 \leq i \leq L \\ 1 \leq j \leq r}} \right]. \quad (2.80)$$

### 3 Graph algebras $\mathcal{B}_{\mathfrak{L}}^{\mathbb{C}}$ , $\mathcal{D}_{\mathfrak{L}}^{\mathbb{C}}$ and $\mathcal{QD}_{\mathfrak{L}}^{\mathbb{C}}$

#### 3.1 Root and Cartan edges of algebra $\mathcal{B}_{\mathfrak{L}}^{\mathbb{C}}$

Now our main goal is to rewrite the calculations of the previous section solely in terms of (1,3)-valent graphs while avoiding any use of Lie algebras. However, it seems that the algebra  $\mathcal{B}_{\mathfrak{L}}$  is not sufficient for this purpose, because the integration in eq.(2.59) does not go over the whole Lie algebra  $\mathfrak{g}$  but only over its space of roots  $\mathfrak{r}$ . Therefore we will construct a bigger algebra  $\mathcal{B}_{\mathfrak{L}}^{\mathbb{C}}$  which reflects a distinction between the root space and Cartan subalgebra.




Figure 2: Types of edges in graphs of  $\mathcal{B}^{\mathbb{C}}$

We begin by defining a bigger space  $\tilde{\mathcal{B}}_{\mathfrak{L}}^{\mathbb{C}}$ . Similarly to  $\mathcal{B}_{\mathfrak{L}}$ , it consists of the formal linear combinations of (1,3)-valent graphs with cyclic order fixed at 3-valent vertices, while the graphs which differ by the cyclic orientation at one vertex, are considered opposites of each other (the AS relation). As usual, the 1-valent vertices of the graphs are labeled by the elements of  $\mathfrak{L}$ . However, in contrast to  $\mathcal{B}_{\mathfrak{L}}$ , we allow two different basic types of edges: Cartan edges (which are depicted by dotted lines) and root edges (which are depicted by dashed lines). For convenience, we define two ‘auxiliary’ types of edges which are expressed in terms of the basic types. First, a ‘total’ edge (which we depict by a standard thin line) is a ‘sum’ of the root edge and the Cartan edge. In other words, a graph with a total edge is



a sum of two graphs in which the total edge is replaced by the root edge and by the Cartan edge. Second, an ‘any’ edge can be of any type, we depict it by a thick solid line. All types of edges are depicted in Fig. 2.



Figure 3: Cartan-commutator relations  $CC_1$  and  $CC_2$

Next, we exclude the ‘Cartan-commutator’ or CC graphs. A graph is called  $CC_1$  if it contains a 3-vertex which is incident to 3 Cartan edges. We call a subgraph  $D'$  of  $D$  *proper*, if all vertices of  $D'$  are either 1-valent or 3-valent. A graph is called  $CC_2$ , if it contains a proper subgraph, all of whose legs are Cartan except for one leg, which is root. A graph is called CC, if it is either  $CC_1$  or  $CC_2$ , and we exclude all these graphs from  $\tilde{\mathcal{B}}_{\mathfrak{L}}^C$ .

Eliminating CC graphs from the set of all graphs can be equivalently described by setting these graphs to zero through factoring over their span. Namely, let  $\tilde{\mathcal{B}}_{\mathfrak{L},CC}^C$  be the span of CC relations, then  $\tilde{\mathcal{B}}_{\mathfrak{L}}^C$  is the span of all possible graphs with Cartan and root edges and  $\tilde{\mathcal{B}}_{\mathfrak{L}}^C$  is defined as its quotient over  $\tilde{\mathcal{B}}_{\mathfrak{L},CC}^C$ . This point of view is useful, for example, when a CC graph appears in an IHX relation.

For future calculations it is convenient to supplement the set of graphs defining  $\tilde{\mathcal{B}}_{\mathfrak{L}}^C$  with two more objects, which are not actually graphs (although we also call them graphs): a root circle  $\bigcirc$  and a Cartan circle  $\bigodot$ . They will be needed in particular in 4.5 in order to define determinants of matrices in graph calculus.

The set of labels  $\mathfrak{L}$  splits into three subsets:  $\mathfrak{L} = \mathfrak{L}_r \cup \mathfrak{L}_c \cup \mathfrak{L}_t$  of root, Cartan and total labels. If a root label is placed on a Cartan leg, or a Cartan label is placed on a root leg, then the graph is considered to be zero. A total label can be placed on both types of legs.

Thus we defined the space  $\tilde{\mathcal{B}}_{\mathfrak{L}}^C$ . We define a familiar subspace  $\tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX}}^C$  as a span of linear combination of the graphs of Fig. 4. As we just mentioned, if a CC graph appears there, then it is considered to be equal to zero. Now we define  $\mathcal{B}_{\mathfrak{L}}^C$  as a quotient space

$$\mathcal{B}_{\mathfrak{L}}^C = \tilde{\mathcal{B}}_{\mathfrak{L}}^C / \tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX}}^C. \quad (3.1)$$




Figure 4: The IHX relation

**Lemma 3.1 (Cartan commutativity)** *Any graph having a 3-vertex incident to two Cartan edges, is CC and therefore is zero in  $\mathcal{B}_{\mathfrak{g}}^{\mathbb{C}}$ .*

*Proof.* If the third edge is Cartan, then the graph is  $\text{CC}_1$ , and if the third edge is root, then the graph is  $\text{CC}_2$ .  $\square$

A  $\mathfrak{g}$ -based weight system for  $\mathcal{B}_{\mathfrak{g}}$  is a linear map

$$\mathcal{T}_{\mathfrak{g}}: \mathcal{B}_{\mathfrak{g}}^{\mathbb{C}} \rightarrow (S\mathfrak{h})^{\otimes |\mathfrak{L}_{\mathbb{C}}|} \otimes ((S\mathfrak{r})^{\otimes |\mathfrak{L}_{\mathfrak{r}}|} \otimes (S\mathfrak{g})^{\otimes |\mathfrak{L}_{\mathfrak{t}}|})^{\mathfrak{T}}, \quad (3.2)$$

which is similar to the second map of (2.10). In order to define it, we modify (2.12). Projectors  $P_{\mathfrak{r}}$  and  $P_{\mathfrak{h}}$  project  $S^2\mathfrak{g}$  naturally onto  $_{\text{str}}$  and  $S^2\mathfrak{h}$ . Thus we construct an element  $h_D^{-1} \in (S^2\mathfrak{g})^{\otimes n_e(D)}$  by taking a tensor product of  $h_{\mathfrak{r}}^{-1} = P_{\mathfrak{r}}(h_{\mathfrak{g}}^{-1})$  for every root edge and  $h_{\mathfrak{h}}^{-1} = P_{\mathfrak{h}}(h_{\mathfrak{g}}^{-1})$  for every Cartan edge and then we change the definition (2.12):

$$f_{\mathfrak{g}}^{\otimes n_3(D)} \otimes h_D^{-1} \xrightarrow{\mathcal{C}_D} \tilde{\mathcal{T}}_{\mathfrak{g}}(D) \in \mathfrak{h}^{\otimes n_{1,\mathfrak{h}}(D)} \otimes \mathfrak{r}^{\otimes n_{1,\mathfrak{r}}(D)} \subset \mathfrak{g}^{\otimes n_1(D)}, \quad (3.3)$$

where  $n_{1,\mathfrak{h}}(D)$  and  $n_{1,\mathfrak{r}}(D)$  are the numbers of Cartan and root legs of  $D$ . The symmetrization over 1-valent vertices which have the same label, projects  $\tilde{\mathcal{T}}_{\mathfrak{g}}(D)$  to  $\mathcal{T}_{\mathfrak{g}}(D) \in (S\mathfrak{h})^{\otimes |\mathfrak{L}_{\mathbb{C}}|} \otimes ((S\mathfrak{r})^{\otimes |\mathfrak{L}_{\mathfrak{r}}|} \otimes (S\mathfrak{g})^{\otimes |\mathfrak{L}_{\mathfrak{t}}|})^{\mathfrak{T}}$ .

For special circle graphs we define  $\mathcal{T}_{\mathfrak{g}}(\text{---}) = \dim \mathfrak{r}$  and  $\mathcal{T}_{\mathfrak{g}}(\text{---}) = \dim \mathfrak{h}$ .

## 3.2 Injections

In this subsection we adopt the definition of  $\tilde{\mathcal{B}}_{\mathfrak{g}}^{\mathbb{C}}$ , in which it includes the CC graphs, so that now

$$\mathcal{B}_{\mathfrak{g}}^{\mathbb{C}} = \tilde{\mathcal{B}}_{\mathfrak{g}}^{\mathbb{C}} / (\tilde{\mathcal{B}}_{\mathfrak{g},\text{IHX}}^{\mathbb{C}} + \tilde{\mathcal{B}}_{\mathfrak{g},\text{CC}}^{\mathbb{C}}), \quad \tilde{\mathcal{B}}_{\mathfrak{g},\text{CC}}^{\mathbb{C}} = \tilde{\mathcal{B}}_{\mathfrak{g},\text{CC}_1}^{\mathbb{C}} + \tilde{\mathcal{B}}_{\mathfrak{g},\text{CC}_2}^{\mathbb{C}}. \quad (3.4)$$

where  $\tilde{\mathcal{B}}_{\mathfrak{L}, \text{CC}_1}^{\text{C}}$  and  $\tilde{\mathcal{B}}_{\mathfrak{L}, \text{CC}_2}^{\text{C}}$  are the spans of  $\text{CC}_1$  and  $\text{CC}_2$  graphs. Also, as usual,  $\tilde{\mathcal{B}}_{\mathfrak{L}}$  is the space of all (1,3)-valent graphs before it was factored over its IHX relation subspace  $\tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}}$ .

Let us consider two injective linear maps

$$\tilde{f}_{\text{trv}}: \tilde{\mathcal{B}}_{\mathfrak{L}} \rightarrow \tilde{\mathcal{B}}_{\mathfrak{L}}^{\text{C}}, \quad \mathfrak{L} = \mathfrak{L}_{\text{t}}, \quad (3.5)$$

$$\tilde{f}_{\text{Crtn}}: \tilde{\mathcal{B}}_{\mathfrak{L}} \rightarrow \tilde{\mathcal{B}}_{\mathfrak{L}}^{\text{C}}, \quad \mathfrak{L} = \mathfrak{L}_{\text{c}}, \quad (3.6)$$

which map each graph  $D$  of  $\tilde{\mathcal{B}}_{\mathfrak{L}}$  into the same graph of  $\tilde{\mathcal{B}}_{\mathfrak{L}}^{\text{C}}$ , such that all edges of  $\tilde{f}_{\text{trv}}(D)$  are total, while all internal edges of  $\tilde{f}_{\text{Crtn}}(D)$  are total and all legs are Cartan. Since  $\tilde{f}_{\text{trv}}(\tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}}), \tilde{f}_{\text{Crtn}}(\tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}}) \subset \tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}}^{\text{C}}$ , then these maps extend to

$$f_{\text{trv}}: \mathcal{B}_{\mathfrak{L}} \rightarrow \mathcal{B}_{\mathfrak{L}}^{\text{C}}, \quad \mathfrak{L} = \mathfrak{L}_{\text{t}}, \quad (3.7)$$

$$f_{\text{Crtn}}: \mathcal{B}_{\mathfrak{L}} \rightarrow \mathcal{B}_{\mathfrak{L}}^{\text{C}}, \quad \mathfrak{L} = \mathfrak{L}_{\text{c}}. \quad (3.8)$$

The weight system (3.2) obviously commutes with  $f_{\text{trv}}$ .

**Theorem 3.2** *The map  $f_{\text{trv}}$  is an injection. If  $\mathfrak{L}$  consists of a single element, then the map  $f_{\text{Crtn}}$  is also an injection.*

In order to prove this Theorem, we need a couple of lemmas about the structure of  $\mathcal{B}_{\mathfrak{L}}$ .

**Lemma 3.3** *If a graph  $D$  of  $\tilde{\mathcal{B}}_{\mathfrak{L}}$  contains a proper non-strut subgraph with a single leg, then this graph belongs to  $\tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}}$  and therefore is zero in  $\mathcal{B}_{\mathfrak{L}} = \tilde{\mathcal{B}}_{\mathfrak{L}} / \tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}}$ .*

*Proof.* We follow the  $\mathcal{B}_{\mathfrak{L}}$  transformations of Fig. 5. Since the subgraph is non-strut, then we




Figure 5: The IHX slide of the asterisk-marked edge onto itself

can ‘pull out’ a 3-valent vertex from it (first equation). The IHX relations in  $\mathcal{B}_{\mathfrak{L}}^{\text{C}}$  allow us to slide the asterisk-marked edge all the way through the grey box onto itself (second equation).

Finally, any graph which has a proper subgraph consisting of a circle with a single leg is equal to zero in view of the AS relation (third equation).  $\square$

Recall that an edge of a graph is called *a bridge*, if it connects two otherwise disconnected components (or, in other words, if its removal increases the number of connected components of the graph). Note that a leg is always a bridge.

**Lemma 3.4** *If  $\mathcal{L}$  consists of a single element, then a graph of  $\tilde{\mathcal{B}}_{\mathcal{L}}$ , in which a 3-valent vertex is incident to 3 bridges, belongs to  $\tilde{\mathcal{B}}_{\mathcal{L},\text{IHX}}$  and therefore is zero in  $\mathcal{B}_{\mathcal{L}} = \tilde{\mathcal{B}}_{\mathcal{L}}/\tilde{\mathcal{B}}_{\mathcal{L},\text{IHX}}$ .*

*Proof.* The proof relies on IHX-slides depicted in Fig. 6. First, we IHX-slide one of the




Figure 6: The IHX-slide of a bridge and the IHX slide of a leg

bridges onto the legs. The summands in the *r.h.s.* of the first equation of Fig. 6 have a 3-vertex, which is incident to a leg and two bridges. Now we IHX-slide that leg onto the legs, as in the second equation of Fig. 6. Each graph in the *r.h.s.* of the second equation has a 3-vertex, which is incident to two legs. It is easy to see that, due to the anti-symmetry of cyclic orientations at 3-vertices, a graph which has a 3-vertex incident to two legs of the same label, is equal to zero.  $\square$

*Proof of Theorem 3.2.* <sup>4</sup> Our strategy for proving the injectivity of the maps  $f_{\text{trv}}, f_{\text{Crtn}}$  is to define the left-inverses of  $\tilde{f}_{\text{trv}}$  and  $\tilde{f}_{\text{Crtn}}$ :

$$\tilde{f}_{\text{trv}}^{-1}, \tilde{f}_{\text{Crtn}}^{-1} : \tilde{\mathcal{B}}_{\Sigma}^{\text{C}} \rightarrow \tilde{\mathcal{B}}_{\Sigma}, \quad \tilde{f}_{\text{trv}}^{-1} \circ \tilde{f}_{\text{trv}} = \tilde{f}_{\text{Crtn}}^{-1} \circ \tilde{f}_{\text{Crtn}} = I. \quad (3.9)$$

Then we will show that

$$\tilde{f}_{\text{trv}}^{-1}(\tilde{\mathcal{B}}_{\Sigma, \text{IHx}}^{\text{C}}), \tilde{f}_{\text{trv}}^{-1}(\tilde{\mathcal{B}}_{\Sigma, \text{CC}}^{\text{C}}) \subset \tilde{\mathcal{B}}_{\Sigma, \text{IHx}}, \quad \tilde{f}_{\text{Crtn}}^{-1}(\tilde{\mathcal{B}}_{\Sigma, \text{IHx}}^{\text{C}}), \tilde{f}_{\text{Crtn}}^{-1}(\tilde{\mathcal{B}}_{\Sigma, \text{CC}}^{\text{C}}) \subset \tilde{\mathcal{B}}_{\Sigma, \text{IHx}}. \quad (3.10)$$

This implies that the maps  $\tilde{f}_{\text{trv}}^{-1}$  and  $\tilde{f}_{\text{Crtn}}^{-1}$  can be extended to the maps  $f_{\text{trv}}^{-1}, f_{\text{Crtn}}^{-1} : \mathcal{B}_{\Sigma}^{\text{C}} \rightarrow \mathcal{B}_{\Sigma}$ , which in view of eq.(3.9) are left-inverses of  $f_{\text{trv}}$  and  $f_{\text{Crtn}}$ :

$$f_{\text{trv}}^{-1} \circ f_{\text{trv}} = f_{\text{Crtn}}^{-1} \circ f_{\text{Crtn}} = I. \quad (3.11)$$

The latter equation means that  $f_{\text{trv}}$  and  $f_{\text{Crtn}}$  are injective and thus proves the theorem.

Thus it remains to define the maps  $\tilde{f}_{\text{trv}}^{-1}$  and  $\tilde{f}_{\text{Crtn}}^{-1}$  which satisfy eqs.(3.9), and prove the inclusions (3.10).

In order to define the maps  $\tilde{f}_{\text{trv}}^{-1}$  and  $\tilde{f}_{\text{Crtn}}^{-1}$  we pick a particular basis in  $\tilde{\mathcal{B}}_{\Sigma}^{\text{C}}$ : it consists of graphs each of whose edges is either root or Cartan. By definition,  $\tilde{f}_{\text{trv}}^{-1}$  maps a basis graph of  $\tilde{\mathcal{B}}_{\Sigma}^{\text{C}}$  into the same graph of  $\tilde{\mathcal{B}}_{\Sigma}$ , if all of its edges are total, and maps it to zero otherwise. Then, obviously,  $\tilde{f}_{\text{trv}}^{-1} \circ \tilde{f}_{\text{trv}} = I$ . The space  $\tilde{\mathcal{B}}_{\Sigma, \text{IHx}}^{\text{C}}$  is a span of relations of Fig. 4, in which all participating graphs are basis. Depending on the nature of their common edges,  $\tilde{f}_{\text{trv}}^{-1}$  either maps the graphs in each triplet to zero, or to the corresponding graphs of  $\tilde{\mathcal{B}}_{\Sigma}$ , which form the  $\tilde{\mathcal{B}}_{\Sigma, \text{IHx}}$  triplets. Therefore,  $\tilde{f}_{\text{trv}}^{-1}(\tilde{\mathcal{B}}_{\Sigma, \text{IHx}}^{\text{C}}) \subset \tilde{\mathcal{B}}_{\Sigma, \text{IHx}}$ . Since  $\tilde{f}_{\text{trv}}^{-1}$  maps all graphs, which have at least one Cartan edge, to zero, then  $\tilde{f}_{\text{trv}}^{-1}(\tilde{\mathcal{B}}_{\Sigma, \text{CC}_1}^{\text{C}}) = 0$ . Almost all  $\text{CC}_2$  graphs also have Cartan edges and hence are also mapped to zero. The only exception are those  $\text{CC}_2$  graphs, whose  $\text{CC}_2$  subgraph has only one leg (which is root). But  $\tilde{f}_{\text{trv}}^{-1}$ -image of such graph is proportional to a graph of  $\tilde{\mathcal{B}}_{\Sigma}$  which, according to Lemma 3.3, belongs to  $\tilde{\mathcal{B}}_{\Sigma, \text{IHx}}$ .

By definition,  $\tilde{f}_{\text{Crtn}}^{-1}$  maps a basis graph of  $\tilde{\mathcal{B}}_{\Sigma}^{\text{C}}$  into the same graph of  $\tilde{\mathcal{B}}_{\Sigma}$  if all of its non-bridge edges are total, and maps it to zero otherwise. Obviously,  $\tilde{f}_{\text{Crtn}}^{-1} \circ \tilde{f}_{\text{Crtn}} = I$ . Also  $\tilde{f}_{\text{Crtn}}^{-1}(\tilde{\mathcal{B}}_{\Sigma, \text{IHx}}^{\text{C}}) \subset \tilde{\mathcal{B}}_{\Sigma, \text{IHx}}$  for the same reason as in the case of  $\tilde{f}_{\text{trv}}^{-1}$ . The  $\tilde{f}_{\text{Crtn}}^{-1}$  image of a  $\text{CC}_1$  graph can be non-zero only if all three Cartan edges of its  $\text{CC}_1$  subgraph are bridges, but according to Lemma 3.4, the image of such graphs belongs to  $\tilde{\mathcal{B}}_{\Sigma, \text{IHx}}$ , so  $\tilde{f}_{\text{Crtn}}^{-1}(\tilde{\mathcal{B}}_{\Sigma, \text{CC}_1}^{\text{C}}) \subset \tilde{\mathcal{B}}_{\Sigma, \text{IHx}}$ . Similarly, the  $\tilde{f}_{\text{Crtn}}^{-1}$  image of a  $\text{CC}_2$  graph can be non-zero only if all Cartan legs of its  $\text{CC}_2$  subgraph are bridges of the graph, but then the root leg of that subgraph must also be a bridge. However, the  $\tilde{f}_{\text{Crtn}}^{-1}$  image of a graph with a root bridge is zero (indeed, we can

---

<sup>4</sup>We are very thankful to D. Thurston, who considerably streamlined the proof of this theorem.

present it as a difference of two graphs, in which this bridge is total and Cartan, but their  $\tilde{f}_{\text{Crtn}}^{-1}$  images are the same, since the action of  $\tilde{f}_{\text{Crtn}}^{-1}$  on basis elements does not depend on the nature of their bridge edges).  $\square$

### 3.3 Symmetric algebras of cohomologies of graphs

For our purposes it will be more convenient to use a different presentation for the space  $\mathcal{B}_{\mathfrak{L}}^{\text{C}}$ . Let  $\mathfrak{L}'_{\text{c}} \subset \mathfrak{L}_{\text{c}}$  be a subset of Cartan labels. We are going to convert Cartan legs with labels of  $\mathfrak{L}'_{\text{c}}$  into the elements of cohomology of (1,3)-valent graphs in exactly the same way as we did it with all legs of the graphs of  $\mathcal{B}$  in [14]. In other words, we will define a new algebra  $\mathcal{D}_{\mathfrak{L}}^{\text{C}}$  and then show that it is canonically isomorphic to  $\mathcal{B}_{\mathfrak{L}}^{\text{C}}$ . The only difference between our case and that of [14] is that the space  $\mathcal{D}$  is based there on 3-valent graphs and ours will be based on (1,3)-valent graphs, while the leg commutativity lemma 3.3 of [14] is replaced by Cartan leg commutativity lemma 3.1. Thus all proofs remain exactly the same, but we will repeat them here for convenience.

Let  $D$  be a general graph. Thinking of it as a  $CW$ -complex, its boundary  $\partial D$  being the set of 1-valent vertices, we may consider a rational relative cohomology space  $H^1(D, \partial D)$ . An oriented edge  $e$  of  $D$  represents an integral element  $\hat{e} \in H^1(D, \partial D)$ , a pairing between  $\hat{e}$  and a cycle of  $D$  being the coefficient at  $e$  in a presentation of the cycle as a linear combination of edges. Moreover,  $H^1(D, \partial D)$  can be presented as a quotient of a linear space, whose basis elements are oriented edges of  $D$ . Let us fix an orientation of the edges of  $D$  and let  $\mathbf{E}(D)$  be the set of these oriented edges.  $E(D)$  denotes the corresponding linear space, whose basis is formed by the elements of  $\mathbf{E}(D)$  (we also assume that edges with opposite orientation represent the opposite elements of  $E(D)$ ). For a vertex  $v$  and an oriented edge  $e$  we define an incidence number  $a_{v,e}$  to be 0 if  $e$  is not incident to  $v$  or if it is a loop attached to  $v$ ,  $a_{v,e} = 1$  if  $e$  goes into  $v$  and  $a_{v,e} = -1$  if  $e$  goes out of  $v$ . Then

$$H^1(D, \partial D) = E(D)/E_{\text{v}}(D), \quad (3.12)$$

where

$$E_{\text{v}}(D) = \bigoplus_{m \geq 2} \text{span} \left( \sum_{e \in \mathbf{E}(D)} a_{v,e} e \mid v \in \mathbf{V}_m(D) \right), \quad (3.13)$$

and  $\mathbf{V}_m(D)$  denotes the set of  $m$ -valent vertices of  $D$ . We excluded the 1-valent vertices from the *r.h.s.* of eq.(3.12), because they represent the boundary of  $D$  and we are interested in the relative cohomology.

Now let  $D$  be a (1,3)-valent graph of  $\mathcal{B}_{\mathcal{L}}^{\mathcal{C}}$  which has no  $\mathcal{L}'$  Cartan legs and let  $H_{\mathcal{C}}^1(D) \subset H^1(D, \partial D)$  denote a linear span of Cartan edges of  $D$ . We will use a quotient space

$$\tilde{H}^1(D) = H^1(D, \partial D) / H_{\mathcal{C}}^1(D). \quad (3.14)$$

It has an alternative description. Let  $\tilde{D}$  be the graph (or, rather, the cell complex) constructed from  $D$  by removing Cartan edges. The  $\text{CC}_2$  condition guarantees that no 3-valent vertices of  $D$  are incident to two Cartan edges. Therefore all 1-valent vertices of  $\tilde{D}$  come from those of  $D$  and as a result in view of eq.(3.12)

$$\tilde{H}^1(D) = H^1(\tilde{D}, \partial \tilde{D}). \quad (3.15)$$

**Remark 3.5** By using the cohomology spaces that we have just introduced, we can rewrite condition  $\text{CC}_2$  of Fig. 4. Namely, a graph  $D$  contains a proper  $\text{CC}_2$  subgraph iff it has a root edge  $e$  such that  $\hat{e} \in H_{\mathcal{C}}^1(D)$ .

Let  $G_D$  be the symmetry group of  $D$ :  $G_D$  maps edges to edges of the same type, 3-vertices to 3-vertices and 1-vertices to 1-vertices of the same label. The elements of  $G_D$  may change orientation at 3-valent vertices.  $G_D$  acts on  $H^1(D, \partial D)$ , and  $H_{\mathcal{C}}^1(D)$  is invariant under that action, so the action of  $G_D$  on  $\tilde{H}^1(D)$  is well-defined. This action can be extended to the algebra  $(S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'|}$ , which is the  $|\mathcal{L}'|$ -th tensor power of the symmetric algebra of  $\tilde{H}^1(D)$ . We modify this action of  $G_D$  by multiplying the action of  $g \in G_D$  by a sign factor  $(-1)^{|g|}$ , where  $|g|$  is the number of 3-vertices of  $D$  whose cyclic order was changed by  $g$ . Now we define the space  $\mathcal{H}_{\mathcal{L}'}(D)$  associated to a graph  $D$  as the  $G_D$ -invariant part of  $(S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'|}$

$$\mathcal{H}_{\mathcal{L}'}(D) = \left( (S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'|} \right)^{G_D}, \quad (3.16)$$

if  $D$  does not contain proper  $\text{CC}$  subgraphs, and  $\mathcal{H}_{\mathcal{L}'}(D)$  is by definition 0-dimensional otherwise. In the former case there is also a natural symmetrization projector

$$(S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'|} \xrightarrow{P_{G_D}} \mathcal{H}_{\mathcal{L}'}(D). \quad (3.17)$$

We define the space  $\mathcal{H}_{\mathcal{L}'}$  also for four special ‘graphs’. The space  $\mathcal{H}_{\mathcal{L}'}(D)$  for the first two of them is defined in accordance with eq.(3.16). The first special graph is a root circle  $\langle \bigcirc \rangle$ , so  $\tilde{H}^1(\langle \bigcirc \rangle) = H^1(\langle \bigcirc \rangle)$  and the symmetry group is  $G_{\langle \bigcirc \rangle} = \{1, -1\}$ , its element  $-1$  flipping the circle. Therefore  $\mathcal{H}_{\mathcal{L}'}(\langle \bigcirc \rangle) = S^{\text{even}} \mathbb{Q}^{|\mathcal{L}'|}$ . The second special graph is a Cartan circle  $\langle \bigodot \rangle$ , so  $\dim \tilde{H}^1(\langle \bigodot \rangle) = 0$  and by definition  $\mathcal{H}_{\mathcal{L}'}(\langle \bigodot \rangle) = \mathbb{Q}$ . The third special graph is an empty unlabeled dot  $\circ$ , it consists of a single vertex and has no edges. We define

$\mathcal{H}_{\mathcal{L}'_c}(\circ) = S^2(\mathbb{Q}^{|\mathcal{L}'_c|})$ . The fourth special graph is a filled labeled dot  $\bullet b$ , it consists of a single vertex labeled by elements of  $\mathcal{L}_c \setminus \mathcal{L}'_c$  and we define  $\mathcal{H}_{\mathcal{L}'_c}(\bullet b) = \mathbb{Q}^{|\mathcal{L}'_c|}$ . If  $D$  is a disjoint union of  $m_1$  graphs  $\langle \_ \rangle$ ,  $m_2$  graphs  $\langle \_ \rangle$ ,  $m_3$  graphs  $\circ$ ,  $m_4(b)$  graphs  $\bullet b$ ,  $b \in \mathcal{L}_c \setminus \mathcal{L}'_c$  and a graph  $D'$  of  $\mathcal{B}_{\mathcal{L}}^C$ , which has no  $\mathcal{L}'_c$  Cartan legs, then we define

$$\begin{aligned} \mathcal{H}_{\mathcal{L}'_c}(D) = & S^{m_1} \mathcal{H}_{\mathcal{L}'_c}(\langle \_ \rangle) \otimes S^{m_2} \mathcal{H}_{\mathcal{L}'_c}(\langle \_ \rangle) \otimes S^{m_3} \mathcal{H}_{\mathcal{L}'_c}(\circ) \\ & \otimes \left( \bigotimes_{b \in \mathcal{L}_c \setminus \mathcal{L}'_c} S^{m_4(b)} \mathcal{H}_{\mathcal{L}'_c}(\bullet b) \right) \otimes \tilde{H}^1(D'). \end{aligned} \quad (3.18)$$

Let  $\mathbf{D}_{\mathcal{L}, \mathcal{L}'_c}$  be the set of all such graphs  $D$ . Then we define the space

$$\tilde{\mathcal{D}}_{\mathcal{L}, \mathcal{L}'_c}^C = \bigoplus_{D \in \mathbf{D}_{\mathcal{L}, \mathcal{L}'_c}} \mathcal{H}_{\mathcal{L}'_c}(D). \quad (3.19)$$

**Remark 3.6** We will abbreviate the notations  $\mathcal{H}_{\mathcal{L}'_c}(D)$ ,  $\mathbf{D}_{\mathcal{L}, \mathcal{L}'_c}$  and  $\tilde{\mathcal{D}}_{\mathcal{L}, \mathcal{L}'_c}^C$  down to  $\mathcal{H}(D)$ ,  $\mathbf{D}_{\mathcal{L}}$  and  $\tilde{\mathcal{D}}_{\mathcal{L}}^C$  when it is clear that a particular set  $\mathcal{L}'_c$  has been chosen.

$\tilde{\mathcal{D}}_{\mathcal{L}}^C$  has a commutative algebra structure. In order to define it, we observe that if  $D_1 \amalg D_2$  is a disjoint union of the graphs  $D_1$  and  $D_2$ , then  $\tilde{H}^1(D_1 \amalg D_2) = \tilde{H}^1(D_1) \oplus \tilde{H}^1(D_2)$  and as a result  $S^* \tilde{H}^1(D_1 \amalg D_2) = S^* \tilde{H}^1(D_1) \otimes S^* \tilde{H}^1(D_2)$  and  $(S^* \tilde{H}^1(D_1 \amalg D_2))^{\otimes |\mathcal{L}'_c|} = (S^* \tilde{H}^1(D_1))^{\otimes |\mathcal{L}'_c|} \otimes (S^* \tilde{H}^1(D_2))^{\otimes |\mathcal{L}'_c|}$ . Therefore  $G_{D_1 \amalg D_2}$  acts on  $(S^* \tilde{H}^1(D_1))^{\otimes |\mathcal{L}'_c|} \otimes (S^* \tilde{H}^1(D_2))^{\otimes |\mathcal{L}'_c|}$  and since  $G_{D_1} \times G_{D_2} \subset G_{D_1 \amalg D_2}$ , then in view of the definition (3.16),  $\mathcal{H}(D_1 \amalg D_2) \subset \mathcal{H}(D_1) \otimes \mathcal{H}(D_2)$  and the symmetrization projector of the type (3.17) establishes a projection

$$\mathcal{H}(D_1) \otimes \mathcal{H}(D_2) \xrightarrow{P_{G_{D_1 \amalg D_2}}} \mathcal{H}(D_1 \amalg D_2). \quad (3.20)$$

The latter map defines the multiplication structure of  $\tilde{\mathcal{D}}_{\mathcal{L}}^C$ .

The spaces  $\mathcal{H}_{\mathcal{L}'_c}(D)$  inherit  $|\mathcal{L}'_c|$  independent gradings  $\deg_{(j)}$  ( $1 \leq j \leq |\mathcal{L}'_c|$ ) from the individual symmetric algebras  $S^* \tilde{H}^1(D)$ . We define  $|\mathcal{L}'_c|$  parity operators  $P_j$  acting on an element  $x \in \mathcal{H}_{\mathcal{L}'_c}(D)$  with definite degree  $\deg_{(j)}$  as

$$P_j(x) = (-1)^{\deg_{(j)}(x)} x. \quad (3.21)$$

Also, we define a grading  $\deg_2$  of  $\tilde{\mathcal{D}}_{\mathcal{L}, \mathcal{L}'_c}^C$  by assigning

$$\deg_2 \mathcal{H}_{\mathcal{L}'_c}(D) = \chi(D) \quad (3.22)$$

(cf. eq.(2.3)). All gradings  $\deg_{(j)}$  as well as  $\deg_2$  are respected by the multiplication (3.20).



Suppose that  $D \in \mathbf{D}_{\mathfrak{L}, \mathfrak{L}'_c}$  and  $D'$  is obtained from  $D$  by replacing one of its root edges  $e$  with a Cartan edge (if  $e$  is a leg, then we assign to it a color from  $\mathfrak{L} \setminus \mathfrak{L}'_c$ ). Obviously,  $\tilde{D}' \subset \tilde{D}$ , so there is a natural map

$$\tilde{H}^1(D) \xrightarrow{f_e} \tilde{H}^1(D'), \quad (3.23)$$

which can be extended to an algebra homomorphism  $S^*\tilde{H}^1(D) \longrightarrow S^*\tilde{H}^1(D')$  and further to

$$\mathcal{H}(D) \xrightarrow{f_e} \mathcal{H}(D'), \quad (3.24)$$

which is a combination of the map  $: \mathcal{H}(D) \rightarrow (S^*\tilde{H}^1(D))^{\otimes |\mathfrak{L}'_c|}$  and a symmetrization projection  $P_{G_{D'}} : (S^*\tilde{H}^1(D))^{\otimes |\mathfrak{L}'_c|} \rightarrow \mathcal{H}(D')$ . If  $D'$  is CC, then by definition  $f_e = 0$ .

The map (3.24) allows us to define the space  $\mathcal{H}(D)$  for a graph  $D$  which, in addition to root and Cartan edges, may also have total edges. In order to simplify notations, suppose that  $D$  has exactly one total edge  $e$ . Let  $D_r$  and  $D_c$  be the graphs constructed from  $D$  by replacing the total edge by a root edge and by a Cartan edge. We define  $\mathcal{H}(D)$  as a subspace of  $\mathcal{H}(D_r) \oplus \mathcal{H}(D_c) \subset \tilde{\mathcal{D}}_{\mathfrak{L}}^C$ :

$$\mathcal{H}(D) = \begin{cases} \mathcal{H}(D_c) \xrightarrow{0 \oplus I} \mathcal{H}(D_r) \oplus \mathcal{H}(D_c), & \text{if } D_r \text{ contains a CC subgraph,} \\ \mathcal{H}(D_r) \xrightarrow{I \oplus f_e} \mathcal{H}(D_r) \oplus \mathcal{H}(D_c) & \text{otherwise,} \end{cases} \quad (3.25)$$

where  $I$  is the identity map and  $f_e$  is the map (3.24). Thus  $\mathcal{H}(D)$  is naturally injected into  $\tilde{\mathcal{D}}_{\mathfrak{L}}^C$ .

Now we define an ideal  $\tilde{\mathcal{D}}_{\mathfrak{L}, \text{IHx}}^C \subset \tilde{\mathcal{D}}_{\mathfrak{L}}^C$ . Consider a (1,3,4)-valent graph  $D$  which has a single 4-valent vertex. We assume that a cyclic order is fixed at its 3- and 4-valent vertices, that 1-valent vertices are colored by the elements of  $\mathfrak{L}$ , that its edges are either root or Cartan and that it has no  $\mathfrak{L}'_c$  Cartan legs. We can ‘resolve’ a 4-valent vertex into two 3-valent vertices connected by a total edge  $e$  in three different ways depicted in Fig. 4. We denote the corresponding graphs of  $\mathbf{D}_{\mathfrak{L}}$  as  $D_i$  ( $i = 1, 2, 3$ ). The graphs  $D_{i,r}$  and  $D_{i,c}$  are constructed from  $D_i$  by declaring  $e$  to be either root or Cartan. For any  $i$ ,  $\tilde{D}$  can be constructed from  $\tilde{D}_{i,r}$  by contracting the new root edge and also from  $\tilde{D}_{i,c}$  by gluing together two 2-vertices, which were 3-vertices of  $D_{i,c}$  incident to the Cartan edge. Therefore, there are natural maps

$$\begin{aligned} H^1(\tilde{D}, \partial \tilde{D}) &\xrightarrow{f_{i,r}} H^1(\tilde{D}_{i,r}, \partial \tilde{D}_{i,r}), \\ H^1(\tilde{D}, \partial \tilde{D}) &\xrightarrow{f_{i,c}} H^1(\tilde{D}_{i,c}, \partial \tilde{D}_{i,c}), \end{aligned} \quad (3.26)$$

which can be extended to symmetric algebra homomorphisms composed with symmetry projectors

$$\begin{aligned} S^*\tilde{H}^1(D_{i,r}) &\xrightarrow{P_{G_{D_{i,r}}}} \mathcal{H}(D_{i,r}) \\ S^*\tilde{H}^1(D_{i,c}) &\xrightarrow{P_{G_{D_{i,c}}}} \mathcal{H}(D_{i,c}). \end{aligned} \quad (3.27)$$

Since, according to the definition (3.25), the spaces  $\mathcal{H}(D_i)$  are either  $\mathcal{H}(D_{i,r})$  or  $\mathcal{H}(D_{i,c})$  (depending on whether  $D_{i,r}$  contains a CC subgraph), the maps (3.27) define maps

$$(S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|} \xrightarrow{f_i} \mathcal{H}(D_i), \quad i = 1, 2, 3. \quad (3.28)$$

Note that if  $D_{i,r}$  does not contain a CC subgraph, then  $f_{i,c} = f_e \circ f_{i,r}$  and therefore a composition of maps

$$(S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|} \xrightarrow{f_i} \mathcal{H}(D_i) \hookrightarrow \mathcal{D}_{\mathcal{L}}^C$$

is equal to the sum  $f_{i,r} + f_{i,c}$  of maps (3.27).

The ideal  $\tilde{\mathcal{D}}_{\mathcal{L},\text{IHX}}^C$  is defined as a span of all elements  $f_1(x) - f_2(x) - f_3(x)$  for all graphs  $D$  and all elements  $x \in (S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'|}$ . Finally, we define the algebra  $\mathcal{D}_{\mathcal{L}}^C$  as a quotient

$$\mathcal{D}_{\mathcal{L}}^C = \tilde{\mathcal{D}}_{\mathcal{L}}^C / \tilde{\mathcal{D}}_{\mathcal{L},\text{IHX}}^C. \quad (3.29)$$

### 3.4 Converting Cartan legs into cohomology

Now we will show that  $\mathcal{D}_{\mathcal{L}}^C$  is canonically isomorphic to the algebra  $\mathcal{B}_{\mathcal{L}}^C$ . First of all, we define a map from the graphs of  $\tilde{\mathcal{B}}_{\mathcal{L}}^C$  to the graphs of  $\mathbf{D}_{\mathcal{L},\mathcal{L}'_c}$ , which removes the  $\mathcal{L}'_c$  legs. A Cartan strut with two  $\mathcal{L}'_c$  labels maps to  $\circ$ . A Cartan strut with one  $\mathcal{L}'_c$  label and one  $\mathcal{L}_c \setminus \mathcal{L}'_c$  label  $b$  maps to  $\bullet b$ . In any other graph,  $\mathcal{L}'_c$  legs are attached to root edges. We remove these legs and ‘dissolve’ their incident vertices, thus obtaining the graphs of  $\mathbf{D}_{\mathcal{L},\mathcal{L}'_c}$ . We call them  $\mathcal{L}'_c$ -frames of the graphs of  $\mathcal{B}_{\mathcal{L}}^C$ . A removal of  $\mathcal{L}'_c$  legs preserves the homotopy class of the CW-complex and hence its Euler characteristic, so if  $D'$  is a  $\mathcal{L}'_c$ -frame of  $D$ , then, according to definitions (2.3) and (3.22),  $\deg_2 D = \deg_2 D'$ .

Let  $\tilde{\mathcal{B}}_{\mathcal{L}}^C(D) \subset \tilde{\mathcal{B}}_{\mathcal{L}}^C$  denote the span of all graphs whose  $\mathcal{L}'_c$ -frame is  $D \in \mathbf{D}_{\mathcal{L},\mathcal{L}'_c}$ . Then  $\tilde{\mathcal{B}}_{\mathcal{L}}^C = \bigoplus_{D \in \mathbf{D}_{\mathcal{L},\mathcal{L}'_c}} \tilde{\mathcal{B}}_{\mathcal{L}}^C(D)$ .

Let  $\tilde{\mathcal{B}}_{\mathcal{L},\text{IHX},i}^C \subset \tilde{\mathcal{B}}_{\mathcal{L},\text{IHX}}^C$  ( $0 \leq i \leq 4$ ) denote a span of IHX relations of Fig. 4, in which  $i$  connecting edges of the IHX graphs are  $\mathcal{L}'_c$  legs. If at least 3 connecting edges are Cartan legs, then the corresponding IHX graphs are CC, so  $\tilde{\mathcal{B}}_{\mathcal{L},\text{IHX},3}^C = \tilde{\mathcal{B}}_{\mathcal{L},\text{IHX},4}^C = 0$  and  $\tilde{\mathcal{B}}_{\mathcal{L},\text{IHX}}^C = \bigoplus_{i=0}^{i=2} \tilde{\mathcal{B}}_{\mathcal{L},\text{IHX},i}^C$ . All graphs in 2 and 3  $\mathcal{L}'_c$ -legged  $\tilde{\mathcal{B}}_{\mathcal{L}}^C$  IHX triplets reduce to the same  $\mathbf{D}_{\mathcal{L},\mathcal{L}'_c}$  graph after the removal of  $\mathcal{L}'_c$  legs. Therefore, if we define

$$\tilde{\mathcal{B}}_{\mathcal{L}}^C(D) = \tilde{\mathcal{B}}_{\mathcal{L}}^C(D) / \tilde{\mathcal{B}}_{\mathcal{L},\text{IHX},2}^C(D), \quad (3.30)$$

where  $\tilde{\mathcal{B}}_{\mathcal{L},\text{IHX},2}^C(D) = \tilde{\mathcal{B}}_{\mathcal{L}}^C(D) \cap \tilde{\mathcal{B}}_{\mathcal{L},\text{IHX},2}^C$ , and if we also define

$$\begin{aligned} \mathcal{B}_{\mathcal{L}}^C(D) &= \tilde{\mathcal{B}}_{\mathcal{L}}^C(D) / (\tilde{\mathcal{B}}_{\mathcal{L}}^C(D) \cap (\tilde{\mathcal{B}}_{\mathcal{L},\text{IHX},1}^C + \tilde{\mathcal{B}}_{\mathcal{L},\text{IHX},2}^C)) \\ &= \tilde{\mathcal{B}}_{\mathcal{L}}^C(D) / \tilde{\mathcal{B}}_{\mathcal{L},\text{IHX},1}^C(D), \end{aligned} \quad (3.31)$$

where  $\tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX},1}^{\mathcal{C}}(D) = (\tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX},1}^{\mathcal{C}} / (\tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX},1}^{\mathcal{C}} \cap \tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX},2}^{\mathcal{C}})) \cap \mathcal{B}_{\mathfrak{L}}^{\mathcal{C}}(D)$ , then

$$\tilde{\mathcal{B}}_{\mathfrak{L}}^{\mathcal{C}} / \tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX},2}^{\mathcal{C}} = \bigoplus_{D \in \mathbf{D}_{\mathfrak{L},\mathfrak{L}'_c}} \tilde{\mathcal{B}}_{\mathfrak{L}}^{\mathcal{C}}(D), \quad (3.32)$$

$$\tilde{\mathcal{B}}_{\mathfrak{L}}^{\mathcal{C}} / (\tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX},1}^{\mathcal{C}} + \tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX},2}^{\mathcal{C}}) = \bigoplus_{D \in \mathbf{D}_{\mathfrak{L},\mathfrak{L}'_c}} \mathcal{B}_{\mathfrak{L}}^{\mathcal{C}}(D) \quad (3.33)$$

and therefore

$$\mathcal{B}_{\mathfrak{L}}^{\mathcal{C}} = \left( \bigoplus_{D \in \mathbf{D}_{\mathfrak{L},\mathfrak{L}'_c}} \mathcal{B}_{\mathfrak{L}}^{\mathcal{C}}(D) \right) / \left( \tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX},0}^{\mathcal{C}} / ((\tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX},1}^{\mathcal{C}} + \tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX},2}^{\mathcal{C}}) \cap \tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX},0}^{\mathcal{C}}) \right). \quad (3.34)$$

In each triplet of IHX graphs with two connecting edges being  $\mathfrak{L}'_c$  legs, one graph has these legs attached to the same 3-vertex, and so it is CC. The remaining two graphs differ only in order in which the two  $\mathfrak{L}'_c$  legs are attached to the (root) edge. Therefore the graphs which differ only in the order in which their  $\mathfrak{L}'_c$  legs are attached to root edges, are equal in  $\tilde{\mathcal{B}}_{\mathfrak{L}}^{\mathcal{C}} / \tilde{\mathcal{B}}_{\mathfrak{L},\text{IHX},2}^{\mathcal{C}}$ .

Next, we establish the isomorphisms  $f_{\mathcal{H}}: \mathcal{H}(D) \rightarrow \mathcal{B}_{\mathfrak{L}}^{\mathcal{C}}(D)$  for all  $D \in \mathbf{D}_{\mathfrak{L},\mathfrak{L}'_c}$ . First, we define it for  $\circ$  and  $\bullet b$ . According to our definition,  $\mathcal{H}_{\mathfrak{L}'_c}(\bullet b) = \mathbb{Q}^{|\mathfrak{L}'_c|}$ . We choose the basis  $e_a$ ,  $a \in \mathfrak{L}'_c$  in this space and then define

$$e_a \xrightarrow{f_{\mathcal{H}}} a \cdots \cdots b.$$

Similarly, we choose the basis  $e_{a_1} e_{a_2}$ ,  $a_1, a_2 \in \mathfrak{L}'_c$  of  $\mathcal{H}_{\mathfrak{L}'_c}(\circ) = \mathbb{S}^2(\mathbb{Q}^{|\mathfrak{L}'_c|})$  and define

$$e_{a_1} e_{a_2} \xrightarrow{f_{\mathcal{H}}} a_1 \cdots \cdots a_2.$$

For the rest of the graphs  $D \in \mathbf{D}_{\mathfrak{L},\mathfrak{L}'_c}$  we define  $f_{\mathcal{H}}$  with the help of the following commutative diagram:

$$\begin{array}{ccccc} & & (S^* E(\tilde{D}))^{\otimes |\mathfrak{L}'_c|} & \xleftarrow{\tilde{f}_{\mathcal{H},s}^{-1}} & \\ & \swarrow P_{G_D} & \downarrow & \searrow \tilde{f}_{\mathcal{H}} & \\ ((S^* E(\tilde{D}))^{\otimes |\mathfrak{L}'_c|})^{G_D} & \xrightarrow{\quad \quad \quad} & \tilde{\mathcal{B}}_{\mathfrak{L}}^{\mathcal{C}}(D) & & \\ & \downarrow f_1 & & & \\ & (S^* \tilde{H}^1(D))^{\otimes |\mathfrak{L}'_c|} & & & \\ & \swarrow P_{G_D} & \downarrow f_{\mathcal{H}} & \searrow f_2 & \\ \mathcal{H}(D) & \xrightarrow{\quad \quad \quad} & \mathcal{B}_{\mathfrak{L}}^{\mathcal{C}}(D) & & \end{array}$$

(3.35)

Recall that  $E(\tilde{D})$  is a linear space, whose basis vectors are oriented root edges of  $D$ . We define the map

$$(S^*E(\tilde{D}))^{\otimes |\mathcal{L}'_c|} \xrightarrow{\tilde{f}_{\mathcal{H}}} \tilde{\mathcal{B}}_{\mathcal{L}}^C(D) \quad (3.36)$$

by its action on the monomials of  $(S^*E(\tilde{D}))^{\otimes |\mathcal{L}'_c|}$ : a monomial  $\prod_{e \in \mathbf{E}(\tilde{D})} \prod_{a \in \mathcal{L}'_c} (e_a)^{m(e_a)}$  is mapped to a graph constructed from  $D$  by attaching  $m(e_a)$  legs of color  $a \in \mathcal{L}'_c$  to the right side of each oriented root edge  $e$ . This map is well-defined, because the order of attaching  $\mathcal{L}'_c$  legs to a root edge does not matter after we take a quotient over  $\tilde{\mathcal{B}}_{\mathcal{L}, \text{IH}, 2}^C$ . The map

$$((S^*E(\tilde{D}))^{\otimes |\mathcal{L}'_c|})^{G_D} \xrightarrow{\tilde{f}_{\mathcal{H}}} \tilde{\mathcal{B}}_{\mathcal{L}}^C(D) \quad (3.37)$$

is the restriction of the former map to the subspace  $((S^*E(\tilde{D}))^{\otimes |\mathcal{L}'_c|})^{G_D} \subset (S^*E(\tilde{D}))^{\otimes |\mathcal{L}'_c|}$ .

**Lemma 3.7** *The map (3.37) is an isomorphism.*

*Proof.* Suppose that  $D'$  is a graph of  $\tilde{\mathcal{B}}_{\mathcal{L}}^C(D)$ . Generally, there may be many ways of identifying its  $\mathcal{L}'_c$ -frame  $D$  with a ‘standard copy’ of  $D$ , but they all differ by a composition with the elements of  $G_D$ . Let  $s$  be a way of identifying the  $\mathcal{L}'_c$ -frames of every graph of  $\tilde{\mathcal{B}}_{\mathcal{L}}^C(D)$  with the standard copy of  $D$ . Then we can construct the left-inverse  $\tilde{\mathcal{B}}_{\mathcal{L}}^C(D) \xrightarrow{\tilde{f}_{\mathcal{H}, s}^{-1}} (S^*E(\tilde{D}))^{\otimes |\mathcal{L}'_c|}$  of  $\tilde{f}_{\mathcal{H}}$ . The composition  $\tilde{f}_{\mathcal{H}}^{-1} = P_{G_D} \circ \tilde{f}_{\mathcal{H}, s}^{-1}$  does not depend on the choice of  $s$  and serves as the left-inverse of (3.37). Since the images of  $\tilde{f}_{\mathcal{H}, s}^{-1}$  for all possible identifications  $s$  span the whole  $(S^*E(\tilde{D}))^{\otimes |\mathcal{L}'_c|}$ , then  $\tilde{f}_{\mathcal{H}}^{-1}$  is surjective, hence (3.37) is an isomorphism.  $\square$

Let us check that the spaces of the lower triangle of the diagram (3.35) are the quotients of the spaces of the upper triangle. According to the definition (3.31),  $\mathcal{B}_{\mathcal{L}}^C(D) = \tilde{\mathcal{B}}_{\mathcal{L}}^C(D)/\tilde{\mathcal{B}}_{\mathcal{L}, \text{IH}, 1}^C(D)$ . According to (3.12),  $\tilde{H}^1(D) = E(\tilde{D})/E_v(\tilde{D})$ , so  $(S^*\tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|} = (S^*E(\tilde{D}))^{\otimes |\mathcal{L}'_c|}/I_v(\tilde{D})$ , where  $I_v(\tilde{D}) \subset (S^*E(\tilde{D}))^{\otimes |\mathcal{L}'_c|}$  is the ideal generated by  $E_v(\tilde{D})$ . Since  $E_v(\tilde{D})$  is invariant under the action of  $G_D$ , then

$$\begin{aligned} \mathcal{H}(D) &= \left( (S^*\tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|} \right)^{G_D} = ((S^*E(\tilde{D}))^{\otimes |\mathcal{L}'_c|})^{G_D} / \left( ((S^*E(\tilde{D}))^{\otimes |\mathcal{L}'_c|})^{G_D} \cap I_v(\tilde{D}) \right). \\ &= ((S^*E(\tilde{D}))^{\otimes |\mathcal{L}'_c|})^{G_D} / (I_v(\tilde{D}))^{G_D}. \end{aligned} \quad (3.38)$$

This follows from a simple

**Lemma 3.8** *Suppose that a finite group  $G$  acts on a linear space  $V$  and its subspace  $W \subset V$  is invariant under this action. Then  $(V/W)^G = V^G/(V^G \cap W) \subset V/W$ .*

*Proof.* Consider a commutative diagram

$$\begin{array}{ccc} V & \xrightarrow{P_G} & V^G \subset V \\ \downarrow f & & \downarrow f \\ V/W & \xrightarrow{P_G} & V^G/(V^G \cap W) \subset V/W \end{array} \quad (3.39)$$

where  $f$  maps  $V$  naturally onto its quotient  $V/W$  and  $P_G$ , as usual, is a  $G$ -symmetrization projector. Since all elements of  $V^G/(V^G \cap W)$  are  $G$ -invariant, then the lower  $P_G$  is surjective, hence  $V^G/(V^G \cap W) = (V/W)^G$ .  $\square$

Thus, the vertical maps  $f_1, f_2$  of the diagram (3.35) are surjections of linear spaces onto their quotients. It is easy to see, that  $\tilde{f}_{\mathcal{H}}$  maps the  $E_v(\tilde{D})$ -generated ideal  $I_v(\tilde{D})$  into the subspace  $\tilde{\mathcal{B}}_{\mathfrak{L}}^{\mathcal{C}}(D) \cap \tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}, 1}^{\mathcal{C}} \subset \tilde{\mathcal{B}}_{\mathfrak{L}}^{\mathcal{C}}(D)$ . Therefore  $\tilde{f}_{\mathcal{H}}$  of (3.37) descends to the map  $(S^* \tilde{H}^1(D))^{\otimes |\mathfrak{L}'_c|} \xrightarrow{f_{\mathcal{H}}} \tilde{\mathcal{B}}_{\mathfrak{L}}^{\mathcal{C}}(D)$ .

**Lemma 3.9** *The restriction of  $f_{\mathcal{H}}$*

$$\mathcal{H}(D) \xrightarrow{f_{\mathcal{H}}} \mathcal{B}_{\mathfrak{L}}^{\mathcal{C}}(D) \quad (3.40)$$

*is an isomorphism.*

*Proof.* In view of Lemma (3.7), it is sufficient to show that the inverse map  $\tilde{f}_{\mathcal{H}}^{-1}$  descends to a map  $\mathcal{B}_{\mathfrak{L}}^{\mathcal{C}}(D) \xrightarrow{f_{\mathcal{H}}^{-1}} \mathcal{H}(D)$ . The latter statement would follow from the fact that  $\tilde{f}_{\mathcal{H}}^{-1}(\ker f_2) \subset \ker f_1$ . In order to see this, observe, that  $\ker f_2 = \tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}, 1}^{\mathcal{C}}(D)$ . Consider a triplet of IHX graphs from  $\tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}, 1}^{\mathcal{C}}(D)$ . Suppose that an identification assignment  $s$  identifies their  $\mathfrak{L}'_c$ -frames consistently with the standard copy of  $D$ . Then  $\tilde{f}_{\mathcal{H}, s}^{-1}$  maps their IHX linear combination into  $I_v(\tilde{D})$ . Since  $\tilde{f}_{\mathcal{H}}^{-1} = P_{G_D} \circ \tilde{f}_{\mathcal{H}, s}^{-1}$  does not depend on the choice of  $s$ , then all  $\tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}, 1}^{\mathcal{C}}(D)$  triplets are mapped into  $(I_v(\tilde{D}))^{G_D} = \ker f_1$ .  $\square$

Thus, if we apply a sum of individual maps  $f_{\mathcal{H}}$  to the direct sum  $\bigoplus_{D \in \mathbf{D}_{\mathfrak{L}, \mathfrak{L}'_c}} \mathcal{H}(D)$ , then in view of relations (3.19) and (3.33) we establish an isomorphism

$$\tilde{\mathcal{D}}_{\mathfrak{L}}^{\mathcal{C}} \xrightarrow{f_{\mathcal{H}}} \tilde{\mathcal{B}}_{\mathfrak{L}}^{\mathcal{C}} / (\tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}, 1}^{\mathcal{C}} + \tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}, 2}^{\mathcal{C}}). \quad (3.41)$$

We leave it for the reader to verify the following

**Lemma 3.10** *The map (3.41) establishes the isomorphism between the IHX subspaces  $\tilde{\mathcal{D}}_{\mathfrak{L}, \text{IHX}}^{\mathcal{C}} \subset \tilde{\mathcal{D}}_{\mathfrak{L}}^{\mathcal{C}}$  and  $\tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}, 0}^{\mathcal{C}} / ((\tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}, 1}^{\mathcal{C}} + \tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}, 2}^{\mathcal{C}}) \cap \tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}, 0}^{\mathcal{C}}) \subset \tilde{\mathcal{B}}_{\mathfrak{L}}^{\mathcal{C}} / (\tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}, 1}^{\mathcal{C}} + \tilde{\mathcal{B}}_{\mathfrak{L}, \text{IHX}, 2}^{\mathcal{C}})$ .*

*Proof.* Recall the definition of  $\tilde{\mathcal{D}}_{\Sigma, \text{IHX}}^{\text{C}}$ . Let  $D$  be a (1,3)-valent graph without  $\Sigma'_c$  legs and with a single 4-vertex. Consider a monomial

$$\prod_{e \in \mathbf{E}(\tilde{D})} \prod_{a \in \Sigma'_c} (\hat{e}_a)^{m(e_a)} \in (S^* \tilde{H}^1(D))^{\otimes |\Sigma'_c|} \quad (3.42)$$

and let  $D'$  be a graph representing its  $f_{\mathcal{H}}$  image (in other words,  $D'$  is a result of attaching  $\Sigma'_c$  legs to the root edges of  $D$  according to the powers  $m(e_a)$ ). Let us compare the IHX elements constructed from the monomial and from  $D'$ .

The root edges of  $D$  are naturally identified with the root edges of the graphs  $D_{i,r}$  and  $D_{i,c}$ . The naturality of the maps (3.26) guarantees that if a root edge  $e$  of  $D$  corresponds to a root edge  $e'$  of  $D_{i,r}$  (or  $D_{i,c}$ ), then  $f_{i,r}(\hat{e}) = \hat{e}'$  (or  $f_{i,c}(\hat{e}) = \hat{e}'$ ). This means that if we apply the maps (3.28) to the monomial (3.42) and then apply  $f_{\mathcal{H}}$  to the resulting monomials, the result will be the graphs constructed by IHX resolving the 4-vertex of  $D'$ . This proves that

$$f_{\mathcal{H}}(\tilde{\mathcal{D}}_{\Sigma, \text{IHX}}^{\text{C}}) = \tilde{\mathcal{B}}_{\Sigma, \text{IHX}, 0}^{\text{C}} / ((\tilde{\mathcal{B}}_{\Sigma, \text{IHX}, 1}^{\text{C}} + \tilde{\mathcal{B}}_{\Sigma, \text{IHX}, 2}^{\text{C}}) \cap \tilde{\mathcal{B}}_{\Sigma, \text{IHX}, 0}^{\text{C}}). \quad (3.43)$$

□

Now, in view of the definition (3.29) and relation (3.34) we come to the following

**Theorem 3.11** *The map  $\tilde{f}_{\mathcal{H}}$  of (3.36) descends to the isomorphism*

$$\mathcal{D}_{\Sigma}^{\text{C}} \xrightarrow{f_{\mathcal{H}}} \mathcal{B}_{\Sigma}^{\text{C}}, \quad (3.44)$$

*which preserves the grading  $\deg_2$ .*

### 3.5 Edge-related denominators

For the purpose of describing the universal  $U(1)$ -RCC invariant we have to consider a modified algebra  $\mathcal{QD}_{\Sigma, \Sigma'_c}^{\text{C}}$  (which we abbreviate down to  $\mathcal{QD}_{\Sigma}^{\text{C}}$ ). This algebra is similar to  $\mathcal{D}_{\Sigma}^{\text{C}}$ , except that in constructing the analogs of the spaces  $\mathcal{H}(D)$  we allow the ‘edge related’ denominators. As a result, whenever we define the analogs of operations in  $\mathcal{D}_{\Sigma}^{\text{C}}$ , we have to make sure that we do not produce zeroes in these denominators.

We begin by defining the analogs of the spaces  $\mathcal{H}(D)$ , which we call  $\mathcal{QH}(D)$ . Let  $D \in \mathbf{D}_{\Sigma, \Sigma'_c}$  be a regular graph. If it contains a CC subgraph, then  $\mathcal{QH}(D)$  is by definition zero-dimensional. Suppose now that  $D$  does not contain CC subgraphs. According to Remark 3.5, this means that for any root edge  $e$  of  $D$ ,  $\hat{e} \notin H_C^1(D)$ . Therefore, as an element of the quotient

space  $H_C^1(D)$ ,  $\hat{e} \neq 0$ , hence any polynomial of  $\hat{e}_{(x)}$  is non-zero as an element of  $(S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|}$ . Thus we can define  $\mathcal{Q}_e((S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|})$  as an extension of the algebra  $(S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|}$ , which is a linear span of fractions  $\frac{f}{g}$ , where  $f \in (S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|}$  and

$$g = \prod_{e \in \mathbf{E}(\tilde{D})} p_e((\hat{e}_a)_{a \in \mathcal{L}'_c}), \quad (3.45)$$

where  $p_e((\hat{e}_a)_{a \in \mathcal{L}'_c}) \in (S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|}$  are formal power series of  $(\hat{e}_a)_{a \in \mathcal{L}'_c}$  (that is, each formal power series depends on a particular root edge of all possible  $\mathcal{L}'_c$  labels). We denote the  $G_D$ -invariant part of this algebra as

$$\mathcal{QH}_{\mathcal{L}'_c}(D) = \mathcal{Q}_e((S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|})^{G_D} \quad (3.46)$$

Let us consider a couple of useful examples of these spaces. First, let  $D$  be a root strut with different colors at 1-vertices:  $D = x_1 - - - x_2$ . The single edge of  $D$  is a natural basis element of  $H^1(D, \partial D)$ ,  $H_C^1(D)$  is trivial and  $G_D$  is also trivial. Hence there is a canonical isomorphism

$$\mathcal{QH}(x_1 - - - x_2) = \mathcal{Q}(\mathbb{Q}[[\mathbf{a}]]), \quad (3.47)$$

where  $\mathbf{a} = (a_1, \dots, a_{|\mathcal{L}'_c|})$  and  $\mathcal{Q}(R)$  denotes a field of quotients of a ring  $R$ . Consider a map  $D \rightarrow D$  which flips the root strut. This map is not a symmetry of the labeled graph, but it acts on  $H^1(D, \partial D)$  by changing the sign of the basis element. We extend its action to the whole space  $\mathcal{QH}(D)$ . We denote the image of  $l \in \mathcal{QH}(D)$  as  $l^*$ . If we represent  $l$  as an element of  $\mathcal{Q}(\mathbb{Q}[[\mathbf{a}]])$ , then obviously

$$l^*(\mathbf{a}) = l(-\mathbf{a}). \quad (3.48)$$

Our second example is a root strut with the same colors at 1-vertices:  $D = x - - - x$ . The difference with the first example is that the graph symmetry group is non-trivial:  $G_D = \{1, -1\}$  where the  $-1$  element reverses the edge. Therefore,  $\mathcal{QH}(D)$  is just the (simultaneously) even part of the algebra (3.47)

$$\mathcal{QH}(x - - - x) = \left[ \mathcal{Q}(\mathbb{Q}[[\mathbf{a}]]) \right]^{\text{even}}. \quad (3.49)$$

It remains to define  $\mathcal{QH}_{\mathcal{L}'_c}(D)$  for special ‘graphs’. We define

$$\mathcal{QH}^1(\text{ } \circ \text{ } ) = \tilde{H}^1(\text{ } \circ \text{ } ), \quad \mathcal{QH}_{\mathcal{L}'_c}(\circ) = \mathcal{H}_{\mathcal{L}'_c}(\circ), \quad \mathcal{QH}_{\mathcal{L}'_c}(\bullet b) = \mathcal{H}_{\mathcal{L}'_c}(\bullet b). \quad (3.50)$$

Since the cohomology and symmetry of the root circle  $\text{ } \circ \text{ }$  are exactly the same as those of  $x - - - x$ , then we define

$$\mathcal{QH}(\text{ } \circ \text{ } ) = \left[ \mathcal{Q}(\mathbb{Q}[[\mathbf{a}]]) \right]^{\text{even}}. \quad (3.51)$$

**Remark 3.12** There is an obvious inclusion and projection which relate the algebras (3.47), (3.49) and (3.51)

$$\left[ \mathcal{Q}(\mathbb{Q}[[\mathbf{a}]]) \right]^{\text{even}} \hookrightarrow \mathcal{Q}(\mathbb{Q}[[\mathbf{a}]]) , \quad \mathcal{Q}(\mathbb{Q}[[\mathbf{a}]]) \longrightarrow \left[ \mathcal{Q}(\mathbb{Q}[[\mathbf{a}]]) \right]^{\text{even}} . \quad (3.52)$$

**Remark 3.13** Actually, we assume that the algebras  $\mathcal{QH}(\begin{smallmatrix} \square \\ \circ \end{smallmatrix})$  and  $\mathcal{QH}(\begin{smallmatrix} \square \\ \circ \end{smallmatrix})$  are extended by the logarithms of  $\left[ \mathcal{Q}(\mathbb{Q}[[\mathbf{a}]]) \right]^{\text{even}}$  and  $\mathbb{Q}$ .

Now we define the analog of  $\tilde{\mathcal{D}}_{\mathfrak{L}, \mathfrak{L}'_c}^C$

$$\mathcal{Q}\tilde{\mathcal{D}}_{\mathfrak{L}, \mathfrak{L}'_c}^C = \bigoplus_{D \in \mathbf{D}_{\mathfrak{L}, \mathfrak{L}'_c}} \mathcal{QH}_{\mathfrak{L}'_c}(D). \quad (3.53)$$

As usual, we will tend to drop  $\mathfrak{L}'_c$  from these notations.

Similarly to  $\tilde{\mathcal{D}}_{\mathfrak{L}}^C$ ,  $\mathcal{Q}\tilde{\mathcal{D}}_{\mathfrak{L}}^C$  has a structure of a graded commutative algebra, its multiplication and grading  $\deg_2$  being defined by the obvious analogs of (3.20) and (3.22). Although the spaces  $\mathcal{QH}(D)$  do not have gradings  $\deg_{(j)}$  (because of denominators), we can still define the action of parity operators  $P_j$  of eq.(3.21) on  $\mathcal{Q}\tilde{\mathcal{D}}_{\mathfrak{L}}^C$  by the formula

$$P_j(x/y) = P_j(x)/P_j(y), \quad x, y \in \mathcal{H}(D). \quad (3.54)$$

The definition of  $\mathcal{Q}\tilde{\mathcal{D}}_{\mathfrak{L}, \text{IHx}}^C$  involves a total edge and therefore requires some care. Let  $e$  be a root edge of a graph  $D$ . Let  $D'$  denote a graph constructed from  $D$  by replacing  $e$  with a Cartan edge. We call an element  $x \in \mathcal{QH}(D)$  *non-singular at a root edge  $e$  of  $D$*  if either  $D'$  contains a  $\text{CC}_2$  subgraph or  $x$  can be presented as a sum of fractions, whose denominators do not belong to the kernel of the unsymmetrized version of the map  $f_e$  of (3.24):

$$(\mathbb{S}^* \tilde{H}^1(D))^{\otimes |\mathfrak{L}'_c|} \xrightarrow{f_e} (\mathbb{S}^* \tilde{H}^1(D'))^{\otimes |\mathfrak{L}'_c|}. \quad (3.55)$$

**Lemma 3.14** *If for a root edge  $e_0$  of  $D$ , an element  $x \in \mathcal{QH}(D)$  has a presentation as a sum of fractions such that*

$$p_{e_0}((\hat{e}_{0,a})_{a \in \mathfrak{L}'_c}) \Big|_{\hat{e}_{0,a}=0 \text{ for all } a \in \mathfrak{L}'_c} \neq 0 \quad (3.56)$$

*in all their denominators (3.45), then  $x$  is non-singular at  $e_0$ .*

*Proof.* A product (3.45) belongs to the kernel of (3.55), if at least one of the polynomials  $p_e((\hat{e}_a)_{a \in \mathfrak{L}'_c})$  belongs to it, and that happens only if the corresponding edge  $\hat{e}$  belongs to the kernel of the map (3.23). That would mean that  $\hat{e} \in H_C^1(D')$ , where  $D'$  is the graph



constructed from  $D$  by declaring  $e_0$  to be Cartan. But then, according to Remark 3.5,  $D'$  contains a  $\text{CC}_2$  subgraph, so  $x$  is still non-singular at  $e_0$ .  $\square$

All elements of  $\mathcal{QH}(D)$ , which are non-singular at  $e$ , form a subalgebra  $\mathcal{QH}_e(D) \subset \mathcal{QH}(D)$ . The map (3.24) extends to  $\mathcal{QH}_e(D)$

$$\mathcal{QH}_e(D) \xrightarrow{f_e} \mathcal{QH}(D'), \quad (3.57)$$

if we define  $f_e$  to be indentially zero if  $D'$  contains a  $\text{CC}_2$  subgraph (this is the only possible definition because in this case the space  $\mathcal{QH}(D')$  is trivial).

The map (3.57) allows us to define a space  $\mathcal{QH}(D)$  for a graph  $D$  with a total edge  $e$  as a subspace of  $\mathcal{QH}(D_r) \oplus \mathcal{QH}(D_c)$ :

$$\mathcal{QH}(D) = \begin{cases} \mathcal{QH}(D_c) \xrightarrow{0 \oplus I} \mathcal{QH}(D_r) \oplus \mathcal{QH}(D_c), & \text{if } D_r \text{ contains a CC subgraph} \\ \mathcal{QH}_e(D_r) \xrightarrow{I \oplus f_e} \mathcal{QH}(D_r) \oplus \mathcal{QH}(D_c) & \text{otherwise,} \end{cases} \quad (3.58)$$

where  $D_r$  and  $D_c$  are again the graphs constructed from  $D$  by declaring  $e$  to be root or Cartan.

Now we define  $\mathcal{QD}_{\Sigma, \text{IHx}}^C$  by copying the definition of  $\tilde{\mathcal{D}}_{\Sigma, \text{IHx}}^C$ . Again, let  $D$  be a graph with a 4-valent vertex and  $D_1, D_2$  and  $D_3$  be (1,3)-valent graphs of Fig. 4 with a single total edge  $e$  constructed by resolving the 4-valent vertex of  $D$ . This time we extend the linear maps (3.26) to the maps

$$\begin{aligned} \mathcal{Q}_e((S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|}) &\xrightarrow{f_{i,c}} \mathcal{QH}(D_{i,c}) && \text{if } D_{i,r} \text{ contains a CC subgraph} \\ \mathcal{Q}_e((S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|}) &\xrightarrow{f_{i,r}} \mathcal{QH}_e(D_{i,r}) && \text{otherwise.} \end{aligned} \quad (3.59)$$

**Lemma 3.15** *The homomorphisms (3.59) are well-defined.*

*Proof.* If  $D_{i,r}$  contains a  $\text{CC}$  subgraph, then  $\hat{e} \in H_C^1(D_{i,r})$  and this means that the map  $f_{i,c}$  of (3.26) has no kernel. Therefore, denominators of the elements of  $\mathcal{Q}_e((S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|})$  do not map into zero, and the map  $f_{i,c}$  of (3.59) is well-defined.

Lemma 3.14 guarantees that the image of the map  $f_{i,r}$  is contained in  $\mathcal{QH}_e(D_{i,r}) \subset \mathcal{QH}(D_{i,r})$ , because the edge  $e$  is not present in  $D$  and therefore does not contribute polynomials to the denominators of the elements of  $\mathcal{Q}_e((S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|})$ .  $\square$

The definition (3.58) allows us to combine the homomorphisms (3.59) into the homomorphisms

$$\mathcal{Q}_e((S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|}) \xrightarrow{f_i} \mathcal{QH}(D_i), \quad i = 1, 2, 3. \quad (3.60)$$

Then the ideal  $\mathcal{Q}\tilde{\mathcal{D}}_{\Sigma, \text{IHX}}^{\mathcal{C}} \subset \mathcal{Q}\tilde{\mathcal{D}}_{\Sigma}^{\mathcal{C}}$  is a span of all elements  $f_1(x) - f_2(x) - f_3(x)$  for all graphs  $D$  with a single 4-valent vertex, and we define the new algebra

$$\mathcal{QD}_{\Sigma}^{\mathcal{C}} = \mathcal{Q}\tilde{\mathcal{D}}_{\Sigma}^{\mathcal{C}} / \mathcal{Q}\tilde{\mathcal{D}}_{\Sigma, \text{IHX}}^{\mathcal{C}}. \quad (3.61)$$

Naturally,  $\mathcal{H}(D) \subset \mathcal{QH}(D)$  for any  $D \in \mathbf{D}_{\Sigma, \mathcal{G}_{\mathcal{C}}}$ , so  $\tilde{\mathcal{D}}_{\Sigma}^{\mathcal{C}}$  is a subalgebra of  $\mathcal{Q}\tilde{\mathcal{D}}_{\Sigma}^{\mathcal{C}}$ . It is easy to see that  $\tilde{\mathcal{D}}_{\Sigma, \text{IHX}}^{\mathcal{C}} \subset \mathcal{Q}\tilde{\mathcal{D}}_{\Sigma, \text{IHX}}^{\mathcal{C}}$ , therefore there is a natural map

$$\mathcal{D}_{\Sigma}^{\mathcal{C}} \xrightarrow{f_{\mathcal{Q}}} \mathcal{QD}_{\Sigma}^{\mathcal{C}}, \quad (3.62)$$

and, in fact, we have a sequence of natural maps

$$\mathcal{B}_{\Sigma} \xrightarrow{f_{\text{trv}}} \mathcal{B}_{\Sigma}^{\mathcal{C}} \xrightarrow{f_{\mathcal{QH}}} \mathcal{QD}_{\Sigma}^{\mathcal{C}}, \quad \text{where } f_{\mathcal{QH}} = f_{\mathcal{Q}} \circ f_{\mathcal{H}}^{-1}. \quad (3.63)$$

**Conjecture 3.16**  $\tilde{\mathcal{D}}_{\Sigma, \text{IHX}}^{\mathcal{C}} = \mathcal{Q}\tilde{\mathcal{D}}_{\Sigma, \text{IHX}}^{\mathcal{C}} \cap \tilde{\mathcal{D}}_{\Sigma}^{\mathcal{C}}$ , and hence the map  $f_{\mathcal{Q}}$  of (3.62) is an injection.

### 3.6 Gluing of legs

The spaces  $\mathcal{D}_{\Sigma}^{\mathcal{C}}$  and  $\mathcal{QD}_{\Sigma}^{\mathcal{C}}$  have algebra structure, but for our future purposes we will need another useful operation on graph spaces: the gluing of legs. The definition of gluing  $\flat$  is very straightforward on  $\mathcal{B}_{\Sigma}$  (see *e.g.* [3]) and on  $\mathcal{B}_{\Sigma}^{\mathcal{C}}$ . First, we define it as a unary operation (that is, linear map)  $\mathcal{B}_{\Sigma}^{\mathcal{C}} \xrightarrow{\flat_{x,y;m}} \mathcal{B}_{\Sigma}^{\mathcal{C}}$  by its action on graphs. Suppose that we have a graph  $D$  with one 1-valent vertex labeled  $x$  and another labeled  $y$ . Then  $\flat_{x,y;1}(D)$  is a graph constructed by joining and ‘dissolving’ these vertices. In other words, we join two legs into a single edge. More precisely, if both legs are of the same type, then we glue them, if one leg is root and the other Cartan, then  $\flat_{x,y;1}(D) = 0$  by definition. If one leg is either root or Cartan and the other is total, then since a total edge is a sum of a root edge and a Cartan edge, then we must define  $\flat_{x,y;1}D$  by gluing them and declaring the resulting edge correspondingly either root or Cartan.

For a positive integer  $m$ , if  $D$  has at least  $m$  legs of each label  $x$  and  $y$ , then  $\flat_{x,y;m}(D)$  is a sum of graphs constructed by gluing  $m$  legs  $x$  with  $m$  legs  $y$  in all the possible ways, and if the number of legs is insufficient, then  $\flat_{x,y;m}(D) = 0$ . If a graph  $D$  is CC, then so are all the graphs of  $\flat_{x,y;m}(D)$ . Therefore  $\flat_{x,y;m}$  is defined as an operation on  $\tilde{\mathcal{B}}_{\Sigma}^{\mathcal{C}}$ . If  $D$  is a graph with a single 4-valent vertex, which generates an IHX triplet  $D_1 - D_2 - D_3$  of Fig. 4, then the graphs of  $\flat_{x,y;m}(D)$  generate  $\flat_{x,y;m}(D_1 - D_2 - D_3)$ . Hence,  $\flat_{x,y;m}(\tilde{\mathcal{B}}_{\Sigma, \text{IHX}}^{\mathcal{C}}) \subset \tilde{\mathcal{B}}_{\Sigma, \text{IHX}}^{\mathcal{C}}$  and  $\flat_{x,y;m}$  is well-defined as an operation on  $\mathcal{B}_{\Sigma}^{\mathcal{C}} = \tilde{\mathcal{B}}_{\Sigma}^{\mathcal{C}} / \tilde{\mathcal{B}}_{\Sigma, \text{IHX}}^{\mathcal{C}}$ .

The binary operation  $D_1 \flat_{x,y;m} D_2$  is a composition of two operations: first, taking a product (that is, disconnected union) of  $D_1$  and  $D_2$  and then applying the unary  $\flat_{x,y;m}$  to it.

**Remark 3.17** Gluing changes the grading  $\deg_2$  in a simple way:

$$\deg_2 \flat_{x,y;m}(D) = \deg_2 D + m. \quad (3.64)$$

Now we have to define  $\flat$  on  $\mathcal{QD}_{\mathfrak{L}, \mathfrak{L}'_c}^C$  in such a way that its restriction to  $\mathcal{D}_{\mathfrak{L}, \mathfrak{L}'_c}^C$  coincides with  $\flat$  on  $\mathcal{B}_{\mathfrak{L}}^C$  transferred by the isomorphism (3.44). We will define the gluing of only the legs which are either root or Cartan of labels  $\mathfrak{L} \setminus \mathfrak{L}'_c$ . So let  $D$  be a graph of  $\mathbf{D}_{\mathfrak{L}, \mathfrak{L}'_c}$  which has at least  $m$  legs labeled  $x$  and at least  $m$  legs of the same type which are labeled by  $y$ . Let  $D'_j$   $j = 1, \dots$  be the graphs constructed by gluing  $m$  legs of labels  $x$  and  $y$  pairwise in all possible ways,  $j$  indexing particular ways of gluing. The map

$$\mathcal{QH}(D) \xrightarrow{\flat_{x_1, x_2; m}} \bigoplus_j \mathcal{QH}(D'_j). \quad (3.65)$$

is a sum of individual maps

$$\mathcal{QH}(D) \xrightarrow{\flat_{x_1, x_2; m|j}} \mathcal{QH}(D'_j). \quad (3.66)$$

These maps are defined in the following way. If a graph  $D'_j$  has a proper CC subgraph, then the space  $\mathcal{QH}(D'_j)$  is zero-dimensional and we define the image of (3.66) as zero. If  $D'_j$  does not contain proper CC subgraphs, then we consider a natural map

$$H_1(D'_j, \partial D'_j) \xrightarrow{f_{\flat, j}} H_1(D, \partial D), \quad (3.67)$$

which simply cuts the cycles at gluing points. Then we extend the dual map

$$H^1(D, \partial D) \xrightarrow{f_{\flat, j}} H^1(D'_j, \partial D'_j) \quad (3.68)$$

to the algebra homomorphism  $(S^* \tilde{H}^1(D))^{\otimes |\mathfrak{L}'_c|} \xrightarrow{\flat_{x_1, x_2; m|j}} (S^* \tilde{H}^1(D'_j))^{\otimes |\mathfrak{L}'_c|}$  and then to (3.66). We have to check that (3.66) is defined correctly. More precisely, we have to verify that this map does not produce zeroes in the denominators of  $x \in \mathcal{QH}(D)$  in case when  $D_j$  does not contain proper CC subgraphs. In other words, we have to verify that for any denominator (3.45),

$$\flat_{x_1, x_2; m|j} \left( \prod_{e \in \mathbf{E}(\tilde{D})} p_e((\hat{e}_a)_{a \in \mathfrak{L}'_c}) \right) \neq 0. \quad (3.69)$$

Zero may appear only if  $\flat_{x_1, x_2; m|j}(\hat{e}) = 0$  for some root edge  $e$  of  $D$ , but that would mean that the corresponding root edge of  $D_j$  is in  $H_C^1(D_j)$ . Then, according to Remark 3.5, the graph  $D_j$  has a  $\text{CC}_2$  subgraph and therefore the map (3.66) is defined as identical zero irrespective of denominators. Thus we defined a unary operation  $\tilde{\mathcal{D}}_{\mathfrak{L}}^C \xrightarrow{\flat_{x, y; m}} \tilde{\mathcal{D}}_{\mathfrak{L}}^C$ .

Gluing can also be defined naturally for graphs containing total internal edges, because it does not introduce zeroes into denominators. This is guaranteed by the following

**Lemma 3.18** *For a root edge  $e$  of  $D$ , let  $e'$  denote the corresponding root edge of  $D_j$ . Then  $\flat_{x_1, x_2; m|j}(\mathcal{QH}_e(D)) \subset \mathcal{QH}_e(D_j)$ .*

*Proof.* Let us denote by  $D'$  and  $D'_j$  the graphs obtained from  $D$  and  $D_j$  by declaring  $e$  and  $e'$  to be Cartan. If  $D'_j$  contains a proper  $\text{CC}_2$  subgraph, then all of  $\mathcal{QH}(D)$  is non-singular by definition. Therefore we assume that neither  $D_j$  nor  $D'_j$  contain a proper  $\text{CC}_2$  subgraph. If  $D'_j$  does not contain a proper  $\text{CC}_2$  subgraph, then neither does  $D'$ , because proper  $\text{CC}_2$  subgraphs survive gluing. The maps (3.23) and (3.68) commute, hence their symmetric algebra extensions (3.55) and (3.66) also commute:

$$\begin{array}{ccc} (S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|} & \xrightarrow{\flat_{x_1, x_2; m|j}} & (S^* \tilde{H}^1(D_j))^{\otimes |\mathcal{L}'_c|} \\ \downarrow f_e & & \downarrow f_e \\ (S^* \tilde{H}^1(D'))^{\otimes |\mathcal{L}'_c|} & \xrightarrow{\flat_{x_1, x_2; m|j}} & (S^* \tilde{H}^1(D'_j))^{\otimes |\mathcal{L}'_c|} \end{array} \quad (3.70)$$

Since  $D'$  and  $D'_j$  do not contain  $\text{CC}_2$  subgraphs, then no root edges belong to the kernels of the horizontal maps. Therefore since the denominators of  $x$  do not belong to the kernel of the left map, then their image by the upper map can not belong to the kernel of the right map. Therefore  $\flat_{x_1, x_2; m|j}(x)$  is non-singular at  $e'$ .  $\square$

**Lemma 3.19** *A gluing  $\flat_{x, y; m}$  maps the IHX ideal  $\tilde{\mathcal{D}}_{\mathcal{L}, \text{IHX}}^{\text{C}}$  into itself:*

$$\flat_{x, y; m}(\tilde{\mathcal{D}}_{\mathcal{L}, \text{IHX}}^{\text{C}}) \subset \tilde{\mathcal{D}}_{\mathcal{L}, \text{IHX}}^{\text{C}}. \quad (3.71)$$

*Proof.* Let  $D$  be a graph with a single 4-vertex, and it produces 3 IHX related graphs  $D_i$ ,  $i = 1, 2, 3$ . There is a natural identification between the legs of all 4 graphs  $D, D_1, D_2, D_3$ , so we can index consistently the ways in which the  $x$  and  $y$  legs are glued for each of these graphs. Let us fix a particular gluing  $j$ . Obviously, the graph  $D' = \flat_{x_1, x_2; m|j}(D)$  produces the IHX triplet of graphs  $D'_i = \flat_{x_1, x_2; m|j}(D_i)$ ,  $i = 1, 2, 3$ .

Let  $w \in \mathcal{Q}_e((S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|})$ . Since maps (3.26) commute with the maps (3.68), then the maps (3.60) and (3.66) also commute. Therefore, if  $x \in \mathcal{Q}_e((S^* \tilde{H}^1(D))^{\otimes |\mathcal{L}'_c|})$ , then  $f_i(\flat_{x_1, x_2; m|j}(w)) = \flat_{x_1, x_2; m|j}(f_i(w))$ , so

$$\flat_{x_1, x_2; m|j}(f_1(w) - f_2(w) - f_3(w)) = f_1(z) - f_2(z) - f_3(z), \quad (3.72)$$

where  $z = \flat_{x, y; m}(w)$ . Equation (3.72) means that a glued IHX triplet is itself an IHX triplet, which implies (3.71).  $\square$

The latter lemma indicates that gluing  $\flat_{x, y; m}$  is well-defined on the quotient space  $\mathcal{D}_{\mathcal{L}}^{\text{C}} = \tilde{\mathcal{D}}_{\mathcal{L}}^{\text{C}} / \tilde{\mathcal{D}}_{\mathcal{L}, \text{IHX}}^{\text{C}}$ .

**Remark 3.20** The maps (3.63) commute with leg gluing.

### 3.7 Weight system

Finally, let us describe an algebra homomorphism

$$\mathcal{QD}_{\mathfrak{g}}^{\mathbb{C}} \xrightarrow{\mathcal{T}_{\mathfrak{g}}} \mathcal{Q}((S\mathfrak{h})^{\otimes |\mathfrak{L}'_c|}) \otimes (S\mathfrak{h})^{\otimes |\mathfrak{L}_c \setminus \mathfrak{L}'_c|} \otimes ((S\mathfrak{r})^{\otimes |\mathfrak{L}_r|} \otimes (S\mathfrak{g})^{\otimes |\mathfrak{L}_t|})^{\mathfrak{T}}, \quad (3.73)$$

which coincides with the homomorphism  $\mathcal{T}_{\mathfrak{g}}$  of (3.2) on  $\mathcal{D}_{\mathfrak{g}}^{\mathbb{C}} \subset \mathcal{QD}_{\mathfrak{g}}^{\mathbb{C}}$ . Let us orient the edges of a graph  $D \in \mathbf{D}_{\mathfrak{g}, \mathfrak{L}'_c}$ . We denote by  $\mathbf{E}_{r,D}$  the set of root edges of  $D$  and we denote by  $\Delta_{\mathfrak{g}} \subset \mathfrak{h}$  the set of roots of  $\mathfrak{g}$ . A map  $c: \mathbf{E}_{r,D} \rightarrow \Delta_{\mathfrak{g}}$  is called a root assignment. An assignment is called consistent if the sum of incoming roots is equal to the sum of outgoing roots at every 3-valent vertex of  $D$ . Let  $\mathbf{S}_D$  be the set of all consistent root assignments for  $D$ . For a root  $\vec{\lambda} \in \Delta_{\mathfrak{g}}$  let  $\mathfrak{r}_{\vec{\lambda}} \subset \mathfrak{r}_{\mathbb{C}}$  be the corresponding root space. Obviously,  $\mathfrak{r}_{\mathbb{C}} = \bigoplus_{\vec{\lambda} \in \Delta_{\mathfrak{g}}} \mathfrak{r}_{\vec{\lambda}}$ . We denote the projection of  $h_{\mathfrak{r}}^{-1} \in \mathfrak{r}_{\mathbb{C}}^{\otimes 2}$  onto  $\mathfrak{r}_{\vec{\lambda}} \otimes \mathfrak{r}_{-\vec{\lambda}}$  as  $h_{\mathfrak{r}_{\vec{\lambda}}}^{-1}$  and for a consistent assignement  $c \in \mathbf{S}_D$  we construct a tensor  $h_{\mathfrak{r},c}^{-1} = (\bigotimes_{e \in \mathbf{E}_{r,D}} h_{\mathfrak{r}_{c(e)}}^{-1}) \otimes h_{\mathfrak{h}}^{-1}$ .

Think of  $D$  as a cell complex. Let  $D'$  denote a complex that is obtained from  $D$  by removing all Cartan edges. A consistent assignment  $c$  defines an element  $\hat{c} \in H_1(D', \partial D') \otimes \mathfrak{h}$ . According to the definition (3.14),  $\tilde{H}^1(D) = H^1(D', \partial D')$ , hence an element  $x \in \mathcal{QH}(D) \subset \mathcal{Q}_e((S^* \tilde{H}^1(D))^{\otimes |\mathfrak{L}'_c|})$  can be evaluated on  $\hat{c}^{\otimes |\mathfrak{L}'_c|}$  to produce an element  $x(\hat{c}) \in \mathcal{Q}((S\mathfrak{h})^{\otimes |\mathfrak{L}'_c|})$ . Then we can define a map

$$\mathcal{QH}(D) \xrightarrow{\tilde{\mathcal{T}}_{\mathfrak{g}}} \mathcal{Q}((S\mathfrak{h})^{\otimes |\mathfrak{L}'_c|}) \otimes \mathfrak{h}^{\otimes n_{1,\mathfrak{h}}(D)} \otimes \mathfrak{r}^{\otimes n_{1,\mathfrak{r}}(D)} \quad (3.74)$$

by the formula

$$\tilde{\mathcal{T}}_{\mathfrak{g}}(x) = \sum_{c \in \mathbf{S}_D} x(\hat{c}) \otimes \mathcal{C}_D(f_{\mathfrak{g}}^{\otimes n_3(D)} \otimes h_{\mathfrak{r},c}^{-1}), \quad (3.75)$$

where  $\mathcal{C}_D$  is the contraction map (2.11). The symmetrization over 1-valent vertices of  $D$  which have the same label, projects  $\tilde{\mathcal{T}}_{\mathfrak{g}}(x)$  to  $\mathcal{T}_{\mathfrak{g}}(x) \in \mathcal{Q}((S\mathfrak{h})^{\otimes |\mathfrak{L}'_c|}) \otimes (S\mathfrak{h})^{\otimes |\mathfrak{L}_c \setminus \mathfrak{L}'_c|} \otimes ((S\mathfrak{r})^{\otimes |\mathfrak{L}_r|} \otimes (S\mathfrak{g})^{\otimes |\mathfrak{L}_t|})^{\mathfrak{T}}$ . If we restrict  $\mathcal{T}_{\mathfrak{g}}$  to  $\mathcal{H}(D) \subset \mathcal{QH}(D)$ , then it takes the values in

$$(S\mathfrak{h})^{\otimes |\mathfrak{L}'_c|} \otimes (S\mathfrak{h})^{\otimes |\mathfrak{L}_c \setminus \mathfrak{L}'_c|} \otimes ((S\mathfrak{r})^{\otimes |\mathfrak{L}_r|} \otimes (S\mathfrak{g})^{\otimes |\mathfrak{L}_t|})^{\mathfrak{T}} \cong (S\mathfrak{h})^{\otimes |\mathfrak{L}_c|} \otimes ((S\mathfrak{r})^{\otimes |\mathfrak{L}_r|} \otimes (S\mathfrak{g})^{\otimes |\mathfrak{L}_t|})^{\mathfrak{T}}$$

and coincides with its previous definition which followed (3.3).

Let us describe explicitly the action of  $\mathcal{T}_{\mathfrak{g}}$  on spaces associated to special ‘graphs’. An element  $x \in \mathcal{H}_{\mathfrak{L}'_c}(\circ) = S^2(\mathbb{Q}^{|\mathfrak{L}'_c|})$  can be presented as a symmetric matrix  $\|l_{ij}\|$ ,  $1 \leq i, j \leq |\mathfrak{L}'_c|$ . Then  $\mathcal{T}_{\mathfrak{g}}(x) = \sum_{i,j=1}^{|\mathfrak{L}'_c|} l_{ij} h_{\mathfrak{h};i,j}^{-1} \in S^2\left(\bigoplus_{j=1}^{|\mathfrak{L}'_c|} \mathfrak{h}_j\right)$ , where  $h_{\mathfrak{h};i,j}^{-1} \in \mathfrak{h}_i \otimes \mathfrak{h}_j$  is a copy of  $h_{\mathfrak{h}}^{-1}$ . For  $x \in \mathcal{H}_{\mathfrak{L}'_c}(\begin{smallmatrix} \circ & \circ \\ \vdots & \vdots \\ \circ & \circ \end{smallmatrix}) = \mathbb{Q}$  we define  $\mathcal{T}_{\mathfrak{g}}(x) = x \in \mathbb{Q}$ . Finally, if we think of  $x \in \mathcal{QH}(\begin{smallmatrix} \circ & \circ \\ \vdots & \vdots \\ \circ & \circ \end{smallmatrix}) = \left[\mathcal{Q}(\mathbb{Q}[[\mathbf{a}]])\right]_{\text{even}}$  as a ‘function’ of  $\mathbf{a}$ , then

$$\mathcal{T}_{\mathfrak{g}}(x) = \sum_{\vec{\lambda} \in \Delta_{\mathfrak{g}} \subset \mathfrak{h}} x(\vec{\lambda}_1, \dots, \vec{\lambda}_{|\mathfrak{L}'_c|}) \in \mathcal{Q}((S\mathfrak{h})^{\otimes |\mathfrak{L}'_c|}), \quad (3.76)$$

where  $\vec{\lambda}_j$  is a copy of  $\vec{\lambda}$  in the  $j$ -th factor of  $(S\mathfrak{h})^{\otimes |\mathcal{L}_c|}$ . Since the function  $x(\mathbf{a})$  is even, then the last expression can be written as a sum over only positive roots

$$\mathcal{T}_{\mathfrak{g}}(x) = 2 \sum_{\vec{\lambda} \in \Delta_+} x(\vec{\lambda}_1, \dots, \vec{\lambda}_{|\mathcal{L}_c|}). \quad (3.77)$$

## 4 Calculus and differential geometry of graph algebras

### 4.1 Functions and general tensor fields

We have to introduce a few basic calculus definitions and theorems for algebras  $\mathcal{B}_{\mathcal{L}}$ ,  $\mathcal{B}_{\mathcal{L}}^C$ ,  $\mathcal{D}_{\mathcal{L}, \mathcal{L}'_c}^C$  and  $\mathcal{QD}_{\mathcal{L}, \mathcal{L}'_c}^C$  in order to define a universal graph version of the stationary phase integral of subsection 2.4 and prove its invariance under the shifts of marks on the components of a marked Morse link. Our general strategy is to translate as much differential geometry of  $\mathfrak{g}^*$  and of coadjoint orbits  $\mathcal{O}_{\vec{\alpha}}$  as possible into the language of graph algebras. We are going to define a graph algebra version of functions, vector fields and matrix fields as well as related operations such as a Lie derivative, a determinant and a (stationary phase) integral. An application of a weight system should translate graph objects into their Lie coalgebra relatives in such a way that the operations on graphs become the usual operations of differential geometry of  $\mathfrak{g}^*$  and  $\mathcal{O}_{\vec{\alpha}}$ . Actually, the definition of the stationary phase (*i.e.* gaussian) integral was already given in [3]. Fubini's theorem and the integration by parts formula were also proved there. In [15] we hope to introduce a graph version of differential forms with the help of grassman variables and to use them to prove the Duistermaat-Heckmann theorem for coadjoint orbits.

All our objects are defined as graphs with a particular labeling of legs. Most of our operations are defined either as relabeling or as gluing. Such definitions work equally well for all graph algebras  $\mathcal{B}_{\mathcal{L}}$ ,  $\mathcal{B}_{\mathcal{L}}^C$ ,  $\mathcal{D}_{\mathcal{L}, \mathcal{L}'_c}^C$  and  $\mathcal{QD}_{\mathcal{L}, \mathcal{L}'_c}^C$ . We just assume that all edges in the definitions are of 'any' type and whenever there is gluing, we invoke an appropriate definition. However, the definition of an inverse map, a determinant and a gaussian integral require inverting the coefficients at strut graphs. As a result, these objects are not well-defined on  $\mathcal{D}_{\mathcal{L}, \mathcal{L}'_c}^C$ , but are well-defined on other three graph algebras. We will work with the space  $\mathcal{QD}_{\mathcal{L}, \mathcal{L}'_c}^C$ , because it includes  $\mathcal{D}_{\mathcal{L}, \mathcal{L}'_c}^C$ , while  $\mathcal{D}_{\mathcal{L}, \mathcal{L}'_c}^C$  coincides literally with  $\mathcal{B}_{\mathcal{L}}^C$  if we take  $\mathcal{L}'_c = \emptyset$ . Whenever necessary, we will explain how to modify a definition for the case of  $\mathcal{B}_{\mathcal{L}}$ .

It will be obvious that our definitions and operations commute with the maps (3.63). Only determinants and gaussian integrals present a slight problem. Their definition involves an explicit separation of a 'strut' part of a linear combination of graphs. However, the

definition of struts is not respected by the isomorphism  $f_{\mathcal{QH}}$ : the latter ‘shaves off’ some legs and thus creates new struts out of some haircomb graphs (Fig. 1) of  $\mathcal{B}_{\mathcal{L}}^C$ . We will address this issue separately.

All our objects are graphs and their linear combinations. When we say ‘graph’ in a definition, we actually mean a (possibly infinite) linear combination of graphs. The distinction between the objects is made by the way in which we label their legs. We also allow a labeling of a leg by a linear combination of labels. In this case, following [3], we assume that the result is a linear combination of graphs in which this leg is labeled by individual labels. In particular, this means that if a leg of  $D$  is labeled with 0, then the whole graph is equal to 0.

The legs of our graphs will be labeled with variables  $x, y, \dots$  as well as with ‘differentials’  $dx, dy, \dots$  and with derivatives  $\partial_x, \partial_y, \dots$ . The variables, as labels, may be either root, or Cartan, or total. A total variable  $x$  is essentially a sum of a root variable  $x_r$  and a Cartan variable  $x_c$ . We just have to remember that the spaces  $\mathfrak{r}$  and  $\mathfrak{h}$  related to  $x_r$  and  $x_c$  through the application of the weight system  $\mathcal{T}_{\mathfrak{g}}$ , are parts of the same space  $\mathfrak{g}$ .

Let  $\mathbf{x} = (x_1, \dots, x_L)$  be our ‘coordinate’ labels. They may be of any type. Other variable colors (such as  $y, z, \dots$ ) may be interpreted as parameters. A *function*  $f(\mathbf{x})$  is a graph whose legs are labeled by the variables. Let  $\mathfrak{v}_j$  ( $1 \leq j \leq L$ ) be the spaces  $\mathfrak{g}$ ,  $\mathfrak{r}$  or  $\mathfrak{h}$  depending on whether  $x_j$  is a total, root or Cartan variable. A weight system (3.73) maps  $f$  into an element

$$\mathcal{T}_{\mathfrak{g}}f \in \mathcal{Q}((S\mathfrak{h})^{\otimes |\mathcal{L}'_c|}) \otimes \bigotimes_{j=1}^L S^* \mathfrak{v}_j, \quad (4.1)$$

which can be also interpreted as a formal power series  $\mathcal{T}_{\mathfrak{g}}f(\vec{\mathbf{x}}; \vec{\mathbf{a}}) \in \mathcal{Q}(\mathbb{Q}[[\vec{\mathbf{a}}]])[[\vec{\mathbf{x}}]]$ , where  $\vec{x}_j \in \mathfrak{v}^*$  ( $1 \leq j \leq L$ ) and  $\vec{a}_j \in \mathfrak{h}^*$  ( $1 \leq j \leq |\mathcal{L}'_c|$ ). Thus if we forget about a distinction between a formal power series and a function, then we may think of  $\mathcal{T}_{\mathfrak{g}}F$  as a function of variables  $\vec{\mathbf{x}}$  which also depends on Cartan parameters  $\vec{\mathbf{a}}$ . Obviously,  $\mathcal{T}_{\mathfrak{g}}$  converts a product of graphs in  $\mathcal{QD}_{\mathcal{L}, \mathcal{L}'_c}^C$  into a product of functions.

For positive integer numbers  $m_j, n_j$  ( $1 \leq j \leq L$ ) we define a *tensor field*  $T(\mathbf{x})$  of ranks  $\mathbf{m}, \mathbf{n}$  in  $\mathbf{x}$  as a graph which has exactly one leg labeled by each of the colors  $dx_j^{(i)}$  ( $1 \leq i \leq m_j$ ,  $1 \leq j \leq L$ ),  $\partial_{x_j^{(i)}}$  ( $1 \leq i \leq n_j$ ,  $1 \leq j \leq L$ ), all other legs being labeled by  $\mathbf{x}$ . When applying the weight system  $\mathcal{T}_{\mathfrak{g}}$  to  $T$ , we use extra metric tensors  $h$  in order to convert the spaces  $\mathfrak{v}$  of  $\partial_x$  legs into  $\mathfrak{v}^*$ . Thus,  $\mathcal{T}_{\mathfrak{g}}T \in \mathcal{Q}(\mathbb{Q}[[\vec{\mathbf{a}}]])[[\vec{\mathbf{x}}]] \otimes \bigotimes_{j=1}^L (\mathfrak{v}^{\otimes m_j} \otimes (\mathfrak{v}^*)^{\otimes n_j})$  is indeed a tensor field (formal power series) of variables  $\vec{\mathbf{x}}$ .  $\mathcal{T}_{\mathfrak{g}}$  converts a product of graphs in  $\mathcal{QD}_{\mathcal{L}, \mathcal{L}'_c}^C$  into a pointwise tensor product of tensor fields.

Next, we define a *contraction of indices*  $dx_j^{(i_1)}$  and  $\partial_{x_j^{(i_2)}}$  of  $A$  simply as gluing of the corresponding legs

$$T \longmapsto \mathfrak{b}_{dx_j^{(i_1)}, \partial_{x_j^{(i_2)}}; 1}(T). \quad (4.2)$$

**Lemma 4.1** *The weight system  $\mathcal{T}_{\mathfrak{g}}$  turns this operation into a standard definition of contracting upper and lower indices of a tensor  $\mathfrak{v}_j \otimes \mathfrak{v}_j^* \longrightarrow \mathbb{Q}$ .*

*Proof.* This is obvious, since, according to our convention, a weight system puts a tensor  $h^{-1}$  only on the  $dx$  leg, effectively omitting it on the  $\partial_x$  leg, while a new internal edge produced by gluing carries the same tensor  $h^{-1}$ .  $\square$

The action of the maps (3.63) on a tensor field  $T$  is prescribed by their definitions. In particular, to define the action of  $f_{\mathcal{H}}^{-1}$ , we suppose that the labels in the set  $\mathcal{L}'_c$  are some Cartan variables  $\mathbf{a} = (a_1, \dots, a_L)$ . Assume that the graphs of a tensor  $T(\mathbf{x}, \mathbf{a})$  of  $\mathcal{B}_{\mathcal{L}}^C$  do not have labels  $\partial_{\mathbf{a}}$  or  $d\mathbf{a}$ . Then the action of  $f_{\mathcal{H}}^{-1}$  on  $T$  ‘shaves off’ the legs  $\mathbf{a}$ , converting them into the elements of the spaces (3.46). Thus,  $\mathbf{a}$  are no longer treated as coordinates, and we can write

$$T(\mathbf{x}, \mathbf{a}) \xrightarrow{f_{\mathcal{H}}^{-1}} T(\mathbf{x}) \in \mathcal{D}_{\mathcal{L}}^C, \quad (4.3)$$

keeping the same name  $T$  for the image of the tensor field in  $\mathcal{D}_{\mathcal{L}, \mathcal{L}'_c}^C$ .

A *1-form*  $\omega(\mathbf{x})$  is a tensor field all legs of which are colored by coordinates except for one leg which is colored by one of the differentials  $dx_j$  ( $1 \leq j \leq L$ ).

A *partial differential*  $\partial_{x_j} T$  of a tensor field  $A$  is a sum of graphs constructed from  $T$  by changing a color at one of the legs of  $T$  from  $x_j$  to  $dx_j$  in all the possible ways. Also we define a ‘total differential’

$$\nabla T = \sum_{j=1}^L \partial_{x_j} T. \quad (4.4)$$

For a function  $f$  we define a *differential*  $df = \nabla f$  which is a 1-form.

## 4.2 Vector fields

A *vector field*  $\xi(\mathbf{x})$  is a tensor field all of whose legs are colored by the variables and parameters except for one leg which is colored by a derivative  $\partial_{x_j}$  ( $1 \leq j \leq L$ ). We define a



pairing  $(\omega, \xi)$  between a 1-form  $\omega$  and a vector field  $\xi$  as a contraction between the  $dx$  and  $\partial_x$  indices in the tensor product  $\omega \xi$

$$(\omega, \xi) = \sum_{j=1}^L \omega \flat_{dx_j, \partial_{x_j}; 1} \xi. \quad (4.5)$$

In other words, if  $\omega$  and  $\xi$  are both presented by a single graph, then  $(\omega, \xi)$  is constructed by gluing the differential leg of  $\omega$  with the derivative leg of  $\xi$  if their variables match, and  $(\omega, \xi) = 0$  otherwise. If  $\omega$  and  $\xi$  are sums of graphs, then  $(\omega, \xi)$  is defined by bilinearity.

A vector field  $\xi$  defines a derivation  $\nabla_\xi$  acting on any object  $T$  by the formula

$$\nabla_\xi T = \sum_{j=1}^L \xi \flat_{\partial_{x_j}, x_j; 1} T. \quad (4.6)$$

Obviously, for a function  $f$

$$\nabla_\xi f = (df, \xi). \quad (4.7)$$

A *divergence* of a vector field  $\xi(\mathbf{x})$  (relative to the coordinates  $\mathbf{x}$ ) is a result of gluing the  $\partial_{\mathbf{x}}$  leg of  $\xi$  to one of its  $\mathbf{x}$  legs in all the possible ways:

$$\text{div}_{\mathbf{x}} \xi = \flat_{\partial_{\mathbf{x}}, \mathbf{x}; 1}(\xi). \quad (4.8)$$

$\mathcal{T}_{\mathfrak{g}}$  transforms this divergence into a standard divergence of a vector field.

### 4.3 Diffeomorphisms

Next, we define differentiable maps and diffeomorphisms which we will also call substitutions. For two sets of coordinate labels  $\mathbf{x}, \mathbf{y}$ , a *map*  $\mathbf{y} = F(\mathbf{x})$  is defined as a sum of graphs all legs of which are labeled by  $\mathbf{x}$  and parameters except for one leg which is labeled by (one of the coordinates)  $\mathbf{y}$ . In applying  $\mathcal{T}_{\mathfrak{g}}$  to  $F$  we treat the  $\mathbf{x}$  legs similar to  $\partial_x$  legs in tensor fields: we apply an extra metric tensor  $h$  to the  $\mathbf{y}$  legs so that  $\mathcal{T}_{\mathfrak{g}} F_j \in \mathcal{Q}(\mathbb{Q}[[\vec{\mathbf{a}}]])[[\vec{\mathbf{x}}]] \otimes \mathfrak{v}_j^*$ , where  $F_j$  is the sum of graphs of  $\mathbf{y} = F(\mathbf{x})$  which have the  $y_j$  leg. Thus a set of functions  $\mathcal{T}_{\mathfrak{g}} F_j$  defines a local smooth map (or, more precisely, a formal power series) from  $\vec{\mathbf{x}}$  to  $\vec{\mathbf{y}}$ .

A definition of a composition of maps is obvious, but we will write it in a formal way which is a bit messy. First, for an element  $T \in \mathcal{QD}_{\mathfrak{L}, \mathfrak{L}'_c}^C$  we define a gluing operation

$$\flat_{x, y; \text{LR}}(T) = \sum_{m=0}^{\infty} \flat_{x, y; m}(T)|_{x=y=0}. \quad (4.9)$$

In other words,  $\flat_{x,y;\text{LR}}$  glues all  $x$  legs to all  $y$  legs and it yields 0 if the numbers of  $x$  and  $y$  legs of  $T$  do not match. We also define a multiple gluing  $\flat_{\mathbf{x},\mathbf{y};\text{LR}}$  as a composition of individual gluings  $\flat_{x_j,y_j;\text{LR}}$ ,  $1 \leq j \leq L$ . Then for a map  $\mathbf{y} = F(\mathbf{x})$  we define an element

$$F^{(\text{exp})} = \sum_{j=1}^L \sum_{n=0}^{\infty} \frac{(F_j)^n}{n!}. \quad (4.10)$$

Now a composition of maps  $\mathbf{y} = F(\mathbf{x})$  and  $\mathbf{z} = G(\mathbf{y})$  is a map  $\mathbf{z} = G \circ F(\mathbf{x})$  defined by the formula

$$G \circ F = G \flat_{\mathbf{y},\mathbf{y};\text{LR}} F^{(\text{exp})}, \quad (4.11)$$

which says that all  $\mathbf{y}$  legs of the graphs of  $G$  have to be glued to the matching  $\mathbf{y}$  legs of  $F$  in all possible ways. There exists an identity map

$$\text{Id} = \mathbf{y} \text{ --- } \mathbf{x}, \quad (4.12)$$

where we introduced an abbreviated multi-index notation

$$\mathbf{y} \text{ --- } \mathbf{x} = \sum_{j=1}^L y_j \text{ --- } x_j. \quad (4.13)$$

Obviously,  $F \circ \text{Id} = \text{Id} \circ F = F$ . It is easy to verify that  $\mathcal{T}_{\mathfrak{g}}$  converts the composition (4.11) into a composition of local maps.

We will use an obvious shortcut notation  $\mathbf{z} = G(F(\mathbf{x}))$  for a composition  $\mathbf{z} = G \circ F(\mathbf{x})$ . Moreover, we will use the same ‘substitution’ notation for any tensor field  $T(\mathbf{y})$  in which  $\mathbf{y}$  is replaced by  $\mathbf{x}$  through  $\mathbf{y} = F(\mathbf{x})$ :

$$T(F(\mathbf{x})) = T \flat_{\mathbf{y},\mathbf{y};\text{LR}} F^{(\text{exp})} \quad (4.14)$$

A map  $\mathbf{y} = F(\mathbf{x})$  is called *linear* if it consists only of struts, one vertex of which is colored by  $\mathbf{x}$  and the other by  $\mathbf{y}$ . Let us assume for a moment that all our coordinates are either root or Cartan (that is, if we have a total coordinate, then we split it into a sum of a root and cartan coordinates) and that the first  $L'$  coordinates ( $0 \leq L' \leq L$ ) are root, while the rest are Cartan. Since  $\mathcal{QH}(y \cdots x) = \mathbb{Q} \subset \mathcal{Q}(\mathbb{Q}[[\mathbf{a}]])$  and  $\mathcal{QH}(y \cdots x) = \mathcal{Q}(\mathbb{Q}[[\mathbf{a}]])$  (cf. eqs.(3.53)), then a linear map has a form

$$F = \sum_{i,j=1}^{L'} l_{ij} y_i \cdots x_j + \sum_{i,j=L'+1}^L l_{ij} y_i \cdots x_j, \quad l_{ij} \in \mathcal{Q}(\mathbb{Q}[[\mathbf{a}]]) \quad (4.15)$$

and we define

$$\det F = \det ||l_{ji}(\mathbf{a})||, \quad (4.16)$$

where  $||l_{ji}(\mathbf{a})||$  denotes an  $L \times L$  matrix with entries  $l_{ji}(\mathbf{a})$ . If  $\det F \neq 0$ , then a linear map  $F$  can be inverted

$$F^{-1} = \sum_{i,j=1}^{L'} l_{ij}^{-1} y_i \text{ --- } x_j + \sum_{i,j=L'+1}^L l_{ij}^{-1} y_i \text{ ..... } x_j, \quad F^{-1} \circ F = \text{Id}, \quad (4.17)$$

where  $||l_{ij}^{-1}(\mathbf{a})|| = ||l_{ij}(\mathbf{a})||^{-1}$ . Note that the block-diagonal structure of  $||l_{ij}(\mathbf{a})||$  implies that  $l_{ij} \in \mathbb{Q}$  for  $i, j > L'$ .

**Remark 4.2** In case of the space  $\mathcal{B}_{\mathcal{L}}$  a linear map has a form

$$F = \sum_{i,j=1}^L l_{ij} x_i \text{ --- } x_j, \quad l_{ij} \in \mathbb{Q} \quad (4.18)$$

and we define  $\det F = \det ||l_{ij}||$ .

Now we again assume that coordinates may be of any type. We call a map  $F$  *strut* if it contains only strut graphs. Those strut graphs are of two types: the linear ones  $y_i \text{ --- } x_j$  and the ‘constant’ ones  $y_i \text{ --- } w_j$ , where  $w_j$  are parameters on which  $F$  depends. For any map  $F$ , let us denote by  $F_{\text{str}}$  the strut part of  $F$  and by  $F_{\text{lin}}$  the linear part of  $F$ . We call a map  $F$  *non-degenerate* iff  $\det F_{\text{lin}} \neq 0$ . The following is obvious

**Lemma 4.3** *A strut map  $F$  is invertible iff it is non-degenerate.*

Now we can prove a general statement

**Theorem 4.4** *A map  $F$  is invertible iff it is non-degenerate.*

*Proof.* Since the operation of taking a linear part of a map commutes with composition, then  $F^{-1} \circ F = \text{Id}$  implies  $F_{\text{lin}}^{-1} \circ F_{\text{lin}} = \text{Id}$ . Thus if  $F$  is invertible, then  $F_{\text{lin}}$  is invertible and hence non-degenerate.

Now suppose that  $F_{\text{lin}}$  is non-degenerate. Then, according to Lemma 4.3,  $F_{\text{str}}$  is invertible. It is easy to see that the composite map  $G = (F_{\text{str}})^{-1} \circ F$  does not have constant struts and  $G_{\text{lin}} = \text{Id}$ . In other words,  $G = \text{Id} + G'$ , and the map  $G'$  contains graphs with at least two edges. Therefore  $G$  can be inverted perturbatively: if  $H^{(n)}$  is such that

$(\text{Id} + H^{(n)}) \circ (\text{Id} + G') = 1 + K^{(n+1)}$ , where  $K^{(n+1)}$  contains only graphs with at least  $n + 1$  edges, then we can choose  $H^{(n+1)} = H^{(n)} + K^{(n+1)}$ . Then

$$G^{-1} = \text{Id} + \sum_{n=2}^{\infty} H^{(n)}, \quad (4.19)$$

and  $F^{-1} = G^{-1} \circ F_{\text{str}}^{-1}$ .  $\square$

**Remark 4.5** Although the map  $f_{\mathcal{QH}}$  of (3.63) related the identity maps of  $\mathcal{B}_{\mathfrak{L}}^{\mathbb{C}}$  and  $\mathcal{QD}_{\mathfrak{L}}^{\mathbb{C}}$  and commutes with the composition of maps, yet the definitions of inverse maps are a bit different: some maps, which are degenerate in  $\mathcal{B}_{\mathfrak{L}}^{\mathbb{C}}$ , may have non-degenerate  $f_{\mathcal{QH}}$  images in  $\mathcal{QD}_{\mathfrak{L}}^{\mathbb{C}}$ . Indeed, if  $F$  is a map in  $\mathcal{B}_{\mathfrak{L}}^{\mathbb{C}}$  and  $F' = f_{\mathcal{QH}}(F) \in \mathcal{QD}_{\mathfrak{L}}^{\mathbb{C}}$ , then  $F'_{\text{lin}}$  has a form (4.15)

$$F'_{\text{lin}} = \sum_{i,j=1}^{L'} l_{ij}(\mathbf{a}) y_i - - - x_j + \sum_{i,j=L'+1}^L l_{ij} y_i \dots\dots\dots x_j, \quad (4.20)$$

while

$$F_{\text{lin}} = \sum_{i,j=1}^{L'} l_{ij}(\mathbf{a})|_{\mathbf{a}=0} y_i - - - x_j + \sum_{i,j=L'+1}^L l_{ij} y_i \dots\dots\dots x_j. \quad (4.21)$$

Thus, generally speaking,  $\det F' \neq \det F$ , rather  $\det F'|_{\mathbf{a}=0} = \det F$ , and the reason for this is that the strut part of  $F'$  includes the haircomb graphs of  $F$  with legs from  $\mathfrak{L}'_{\mathfrak{c}}$  (these legs are shaved off by  $f_{\mathcal{QH}}$  and converted into powers of  $\mathbf{a}$ ). Therefore it may happen that  $\det F = 0$ , whereas  $\det F' \neq 0$  and so  $F'$  is non-degenerate. However, in this case its inverse diffeomorphism  $(F')^{-1}$  is not a  $f_{\mathcal{QH}}$  image of any diffeomorphism of  $\mathcal{B}_{\mathfrak{L}}^{\mathbb{C}}$ .

We call invertible maps *diffeomorphisms* or *substitutions*. Theorem 2.3 shows that diffeomorphisms form a group which we call  $\text{Diff}_L$ .  $\mathcal{T}_{\mathfrak{g}}$  generates a homomorphism of this group into a group of local diffeomorphisms of  $\bigoplus_{j=1}^L \mathfrak{v}_j^*$  depending on parameters  $\mathbf{a}$ . We will define the *contragradient* action of  $\text{Diff}_L$  on some tensor fields. This action is converted by  $\mathcal{T}_{\mathfrak{g}}$  into a natural contragradient action of the group of local diffeomorphisms  $\bigoplus_{j=1}^L \mathfrak{v}_j^*$ .

First of all,  $\text{Diff}_L$  acts naturally on functions: a substitution  $\mathbf{y} = F(\mathbf{x})$  converts a function  $f(\mathbf{y})$  it into a function  $F^*f(\mathbf{x}) = f(F(\mathbf{x}))$  of  $\mathbf{x}$  by gluing the  $\mathbf{y}$  legs of  $F$  to all  $\mathbf{y}$  legs of the graphs of  $f$ :

$$F^*f(\mathbf{x}) = f(F(\mathbf{x})) = f \downarrow_{\mathbf{y}, \mathbf{y}; \text{LR}} F^{(\text{exp})}. \quad (4.22)$$

Formally, we can turn a map  $\mathbf{y} = F(\mathbf{x})$  into a vector field by replacing the labels  $\mathbf{y}$  with  $\partial_{\mathbf{x}}$ . We denote this vector field as  $\vec{F}(\mathbf{x})$ . Conversely, we can turn a vector field  $\xi(\mathbf{x})$  into a map  $\mathbf{y} = \tilde{\xi}(\mathbf{x})$  by replacing the labels  $\partial_{\mathbf{x}}$  with  $\mathbf{y}$ .

The Lie algebra of  $\text{Diff}_L$  can be identified with the space of vector fields  $\text{Vect}_L$ . If a diffeomorphism  $\mathbf{y} = F(\mathbf{x}; t)$  depends on a  $\mathbb{Q}$ -parameter  $t$  and  $F(\mathbf{x}; 0) = \text{Id}$ , then we identify the tangent vector to the curve  $\mathbf{y} = F(\mathbf{x}; t)$  in  $\text{Diff}_L$  at  $t = 0$  with the vector field  $\overrightarrow{\partial_t F}|_{t=0}$ . We leave it for the reader to check that the Lie algebra commutator induced on  $\text{Vect}_L$  coincides with the usual vector field Lie bracket defined as

$$[\xi_1, \xi_2] = \nabla_{\xi_1} \xi_2 - \nabla_{\xi_2} \xi_1. \quad (4.23)$$

Since  $\text{Vect}_L$  is a Lie algebra of  $\text{Diff}_L$ , then the action of vector fields on various objects (such as functions, 1-forms and vector fields) is determined by the action of diffeomorphisms. The corresponding action of a vector field  $\xi$  is called a *Lie derivative* and we denote it as  $L_\xi$ . More precisely, suppose that  $\text{Diff}_L$  acts on a certain type of objects (say, tensor fields). Denote an action of  $F \in \text{Diff}_L$  on  $T$  as  $F^*T$ . For a vector field  $\xi(x)$  we construct a 1-parametric family of diffeomorphisms  $F(x; t) = \text{Id} + t\tilde{\xi}$ , so that if  $\text{Diff}_L$  acts on a tensor field  $T$ , then we define  $L_\xi T = \partial_t F^*(t)T|_{t=0}$ . Obviously,  $L_\xi f = \nabla_\xi f$  for a function  $f$  and  $L_\xi \eta = [\xi, \eta]$  for a vector field  $\eta$ . Also for any 1-parametric family of diffeomorphisms  $y = F(x; t)$ ,

$$\partial_t F^*(t)T = L_{(\partial_t F(t)) \circ F^{-1}(t)}(F^*(t)T). \quad (4.24)$$

**Example 4.6** Consider a map  $y = \text{ad}_z(x)$  defined on Fig. 7,  $x, y$  being total variables. Its




Figure 7: A map  $\text{ad}_z$ , a map  $\text{Ad}_{\exp(z)}$  and a vector field  $\vec{\text{ad}}_z$

composition exponent defines another map  $y = \text{Ad}_{\exp(z)}(x)$  as

$$\text{Ad}_{\exp(z)} = \exp_{\circ}(\text{ad}_z) = \sum_{n=0}^{\infty} \frac{(\text{ad}_z)^{\circ n}}{n!}. \quad (4.25)$$

The maps  $\text{Ad}_{\exp(tz)}$ , where  $t$  is a  $\mathbb{Q}$ -parameter, form a 1-parametric subgroup of  $\text{Diff}_x$ , which is generated by the vector field  $\vec{\text{ad}}_z(x)$  of Fig. 7. This means that  $\text{Ad}_{\exp(tz)} = \exp_{\text{Lie}}(t\vec{\text{ad}}_z)$ , where  $\exp_{\text{Lie}}$  means a standard exponential map from a Lie algebra to a Lie group. More

specifically, the action of  $\text{Ad}_{\exp(tz)}$  on any object is equal to the operator product exponential  $\exp(tL_{\vec{\text{ad}}_z})$ .

The IHX relation says that

$$\text{ad}_{\text{ad}_y(x)} = \text{ad}_y \circ \text{ad}_x - \text{ad}_x \circ \text{ad}_y. \quad (4.26)$$

Then it is easy to see that

$$\text{ad}_{\text{Ad}_{\exp(y)}x} = \text{Ad}_{\exp(y)} \circ \text{ad}_x \circ \text{Ad}_{\exp(-y)}, \quad (4.27)$$

and with the help of eq.(4.25) we derive a useful formula

$$\text{Ad}_{\exp(\text{Ad}_{\exp(y)}x)} = \text{Ad}_{\exp(y)} \circ \text{Ad}_{\exp(x)} \circ \text{Ad}_{\exp(-y)}. \quad (4.28)$$

The following lemma follows easily from IHX and CC relations.

**Lemma 4.7** *If a function  $f(x)$  depends only on a single coordinate  $x$  (that is, all its legs are labeled by  $x$ ), then*

$$\nabla_{\vec{\text{ad}}_y} f(x) = 0, \quad (4.29)$$

*if one of the two conditions holds: either if all edges of the graphs of  $f$  are total (that is  $f$  belongs to the image of the map (3.7) and  $x$  is total), or if  $y$  is Cartan.*

**Remark 4.8** If  $f_1(x)$  and  $f_2(x)$  are two functions which may depend on other parameters, then

$$\nabla_{\vec{\text{ad}}_y} \left( f_1(x) \text{ } \text{ } f_2(x) \right) = \left( \nabla_{\vec{\text{ad}}_y} f_1(x) \right) \text{ } \text{ } f_2(x) + f_1(x) \text{ } \text{ } \left( \nabla_{\vec{\text{ad}}_y} f_2(x) \right). \quad (4.30)$$

## 4.4 Derivatives and matrix fields

Let us define a derivative of a diffeomorphism. A derivative is a ‘matrix field’. A *matrix field*  ${}^y_x M(\mathbf{x})$  is a sum of graphs all of whose legs are labeled by coordinates  $\mathbf{x}$  and parameters except for two legs, one carrying a color  $dx_j$  ( $1 \leq j \leq L$ ) and the other carrying a color  $\partial_{y_i}$  ( $1 \leq i \leq L$ ). We will use an abbreviated notation  $M(\mathbf{x})$  for  ${}^x_x M(\mathbf{x})$ . A derivative of a map  $y = F(x)$  is a matrix field  ${}^y_x F'(x)$  whose graphs are constructed from the graphs of  $F$  by replacing the colors  $y_i$  with  $\partial_{y_i}$  and a color  $x_j$  at one leg by  $dx_j$  in all possible ways:

$${}^y_x F' = ( \partial_y \text{ } \text{ } y ) \text{ } \text{ } F \text{ } \text{ } ( x \text{ } \text{ } dx ). \quad (4.31)$$

Obviously,  $\mathcal{T}_{\mathbf{g}}$  converts eq.(4.31) into a standard derivative of a diffeomorphism. A vector field  $F^*\xi(\mathbf{x})$  produced by the contragradient (adjoint) action of a diffeomorphism  $\mathbf{y} = F(\mathbf{x})$  on a vector field  $\xi(\mathbf{y})$  can be expressed as

$$F^* \xi(x) = \left( \begin{matrix} x \\ y \end{matrix} (F^{-1})' \right) \begin{matrix} b_{d\mathbf{y}, \partial \mathbf{y}; 1} \\ \xi(y) \end{matrix} \begin{matrix} b_{\mathbf{y}, \mathbf{y}; \text{LR}} \\ F^{(\text{exp})} \end{matrix}, \quad (4.32)$$

where  $\mathbf{x} = F^{-1}(\mathbf{y})$  is the diffeomorphism inverse to  $\mathbf{y} = F(\mathbf{x})$ .

The matrix fields can be multiplied in a matrix way. We denote this product by the symbol  $\times$  in order to distinguish it from the multiplication in the graph algebra. The formula for the product is transparent: for example,

$${}_y^z M_1 \times {}_x^y M_2 = {}_y^z M_1 \circ_{d_{\mathbf{y}, \partial_{\mathbf{y}}; 1}} {}_x^y M_2. \quad (4.33)$$

The identity matrix field is obviously

$${}^x_x I = \partial_x \text{---} d\mathbf{x} . \quad (4.34)$$

**Theorem 4.9 (chain rule)** *For two maps  $\mathbf{y} = F(\mathbf{x})$  and  $\mathbf{z} = H(\mathbf{y})$ , the derivative of their composition is given by the formula*

$$(H \circ F)'(\mathbf{x}) = H'(F(\mathbf{x})) \times F(\mathbf{x}). \quad (4.35)$$

*Proof.* Let us change the order of gluings in the formula for  $(H \circ F)'$

$$\begin{aligned} (H \circ F)' &= (\partial_z \text{---} z) \mid_{z,z;1} (G \mid_{y,y;\text{LR}} F^{(\text{exp})}) \mid_{x,x;1} (x \text{---} dx) \\ &= (\partial_z \text{---} z) \mid_{z,z;1} \left[ G \mid_{y,y;\text{LR}} \left( F^{(\text{exp})} \mid_{x,x;1} (x \text{---} dx) \right) \right]. \end{aligned} \quad (4.36)$$

The gluing  $F^{(\text{exp})} \quad \text{b}_{\mathbf{x}, \mathbf{x}; 1} \quad (\mathbf{x} \text{ --- } d\mathbf{x})$  replaces a label  $\mathbf{x}$  at one of the legs of one of the graphs of  $F^{(\text{exp})}$  with the corresponding label  $d\mathbf{x}$ . Let us relabel the single  $\mathbf{y}$  leg of that same graph with the label  $\partial_{\mathbf{y}}$  and then relabel the  $\mathbf{y}$  leg of  $G$  which was glued to it by  $\text{b}_{\mathbf{y}, \mathbf{x}; \text{LR}}$  with the label  $d\mathbf{y}$ . Then we come directly to the *r.h.s.* of eq.(4.35).  $\square$

The chain rule has an obvious corollary:

**Corollary 4.10 (derivative of the inverse function)** *If a map  $\mathbf{y} = F(\mathbf{x})$  is nondegenerate, then the derivative of the inverse map  $\mathbf{x} = F^{-1}(\mathbf{y})$  is equal to the matrix inverse of the derivative of  $F$*

$$(F^{-1})'(\mathbf{y}) = \left( F'(F^{-1}(\mathbf{y})) \right)^{\times(-1)}. \quad (4.37)$$

A *trace* of a matrix field  ${}^xM$  is a function defined in an obvious way

$$\mathrm{Tr} {}^xM = \flat_{\partial_x, d\mathbf{x}; 1}({}^xM). \quad (4.38)$$

It is easy to see that

$$\mathrm{Tr}({}^xM_1 \times {}^xM_2) = \mathrm{Tr}({}^xM_2 \times {}^xM_1), \quad \mathrm{Tr} [{}^xM_1, {}^xM_2] = 0, \quad (4.39)$$

where the commutator is defined relative to the matrix product  $\times$ .

We define a *transposed matrix field*  ${}^xM^T$  by switching the  $d\mathbf{x}$  and  $\partial_x$  labels at the legs of the graphs of  ${}^xM$ . Obviously,

$$({}^xM_1 \times {}^xM_2)^T = {}^xM_2^T \times {}^xM_1^T, \quad \mathrm{Tr} {}^xM^T = \mathrm{Tr} {}^xM. \quad (4.40)$$

Let  $\xi(\mathbf{x})$  be a vector field. Consider a map  $\mathbf{y} = \tilde{\xi}(\mathbf{x})$  constructed from  $\xi$  by replacing its  $\partial_x$  colors with  $\mathbf{y}$ . Then

$$\mathrm{Tr} \tilde{\xi}' = \mathrm{div}_{\mathbf{x}} \xi. \quad (4.41)$$

## 4.5 Determinant and its properties

A *determinant* of a matrix field is a function which we will define in stages, while making sure that it is multiplicative

$$\det({}^xM_1 \times {}^xM_2) = (\det {}^xM_1) (\det {}^xM_2). \quad (4.42)$$

First, we define a determinant of a *strut matrix field*

$${}^xM = \sum_{i,j=1}^L l_{ji} \partial_{x_j} \text{---} dx_i, \quad (4.43)$$

Similarly to eq.(4.15), we assume that coordinates  $x_j$ ,  $1 \leq j \leq L'$  are root, while coordinates  $x_j$ ,  $L' + 1 \leq j \leq L$  are Cartan, so that

$${}^xM = \sum_{i,j=1}^{L'} l_{ij} \partial_{x_i} \text{---} \text{---} dx_j + \sum_{i,j=L'+1}^L l_{ij} \partial_{x_i} \text{---} \text{---} \text{---} dx_j, \quad l_{ij} \in \begin{cases} \mathcal{Q}(\mathbb{Q}[[\mathbf{a}]]) , & \text{if } 1 \leq i, j \leq L' \\ \mathbb{Q}, & \text{if } L' \leq i, j \leq L \end{cases} \quad (4.44)$$

Then we define  $\det {}^xM$  as

$$\det {}^xM = \exp \left[ \frac{1}{2} \left( \log(\det ||l_{ij}||_r) + \log(\det ||l_{ij}^*||_r) \right) \begin{pmatrix} \text{---} \end{pmatrix} + \log(\det ||l_{ij}||_c) \begin{pmatrix} \text{---} \end{pmatrix} \right], \quad (4.45)$$



if  $\det ||l_{ij}|| \neq 0$ , and  $\det {}^x M = 0$  if  $\det ||l_{ij}|| \equiv 0$ . Here  $||l_{ij}||_r$  and  $||l_{ij}||_c$  are root and Cartan blocks of the block-diagonal matrix  $||l_{ij}||$ . This definition makes sense, because  $\mathcal{QH}(\begin{smallmatrix} \square \\ \square \end{smallmatrix})$  and  $\mathcal{QH}(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix})$  are related as algebras (see Remark 3.12), so we can calculate the products of  $l_{ji}$  in  $\mathcal{Q}(\mathbb{Q}[[\mathbf{a}]])$  and then project them into  $\left[\mathcal{Q}(\mathbb{Q}[[\mathbf{a}]])\right]^{\text{even}}$ , while Remark 3.13 warned the reader that we would have to include logarithms in the spaces  $\mathcal{QH}(\begin{smallmatrix} \square \\ \square \end{smallmatrix})$  and  $\mathcal{QH}(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix})$ . The multiplicativity of the determinants (4.45) follows easily from the multiplicativity of the matrices  $||l_{ij}||_r$  and  $||l_{ij}||_c$ .

**Remark 4.11** If we work in  $\mathcal{B}_{\mathfrak{L}}$ , then for a strut matrix field (4.43) with  $l_{ij} \in \mathbb{Q}$  we define

$$\det {}^x M = \exp \left( \log(\det ||l_{ij}||) \bigcirc \right). \quad (4.46)$$

We call a strut matrix field non-degenerate if its determinant is non-zero. A non-degenerate strut matrix field can be inverted with the help of a formula

$$({}^x M)^{-1} = \sum_{i,j=1}^L l_{ij}^{-1} \partial_{x_i} \text{---} dx_j. \quad (4.47)$$

Next, we define a determinant of a matrix field which is close to identity, by using a well-known formula which holds for ordinary matrices:

$$\det M = \exp(\text{Tr} \log M). \quad (4.48)$$

In order to abbreviate our notations we will temporarily drop the indices  ${}_x$  from the matrix field notation.

A matrix field  $M$  is called a *perturbative identity* (or simply PI) if it is of the form

$$M = I + M', \quad (4.49)$$

where the matrix field  $M'$  contains only the graphs with at least two edges. For such a matrix we can define the logarithm

$$\log_{\times}(M) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{(M')^{\times n}}{n} \quad (4.50)$$

and then define the determinant

$$\det M = \exp \left( \text{Tr} \log_{\times} M \right). \quad (4.51)$$

**Lemma 4.12** *The determinant of PI matrix fields defined by eq. (4.51) is multiplicative.*

*Proof.* Let  $M_i = I + M'_i$  ( $i = 1, 2$ ) be two PI matrices. According to the BCH formula,

$$\begin{aligned} \exp_{\times} \left( \log_{\times}(M_1) \right) \exp_{\times} \left( \log_{\times}(M_2) \right) \\ = \exp_{\times} \left( \log(M_1) + \log(M_2) + (\text{sum of commutators}) \right) \end{aligned} \quad (4.52)$$

Taking the  $\times$ -logarithms of both sides, we find that

$$\log_{\times}(M_1 \times M_2) = \log_{\times}(M_1) + \log_{\times}(M_2) + (\text{sum of commutators}). \quad (4.53)$$

Then in view of eq.(4.39),

$$\text{Tr} \log_{\times}(M_1 \times M_2) = \text{Tr} \log_{\times} M_1 + \text{Tr} \log_{\times} M_2. \quad (4.54)$$

The multiplicativity (4.42) follows from this formula and the definition (4.51).  $\square$

**Lemma 4.13** *Determinant (4.51) is conjugation-invariant: if  $M_1$  is PI and  $M_2$  is a strut matrix field, then*

$$\det(M_2 \times M_1 \times M_2^{-1}) = \det M_1 \quad (4.55)$$

*Proof.* First, note that the matrix field  $M_2 \times M_1 \times M_2^{-1}$  is PI, so eq.(4.55) makes sense. Then it is easy to see from the definition (4.50) of  $\log_{\times}$  that

$$\log_{\times}(M_2 \times M_1 \times M_2^{-1}) = M_2 (\log_{\times} M_1) M_2^{-1}. \quad (4.56)$$

Now it follows from eq.(4.39) that

$$\text{Tr} \log_{\times}(M_2 \times M_1 \times M_2^{-1}) = \text{Tr} \log_{\times} M_1. \quad (4.57)$$

This equation together with eq.(4.51) implies eq.(4.55).  $\square$

For a matrix field  $M$ , let  $M_{\text{str}}$  denote its strut part. We call  $M$  non-degenerate if  $M_{\text{str}}$  is non-degenerate, that is, if  $\det M_{\text{str}} \neq 0$ . If  $M$  is non-degenerate, then it is easy to see that  $M_{\text{str}}^{-1} \times M$  is PI. Then we define

$$\det M = \begin{cases} \det M_{\text{str}} \det(M_{\text{str}}^{-1} \times M), & \text{if } \det M_{\text{str}} \neq 0, \\ 0, & \text{if } \det M_{\text{str}} \equiv 0, \end{cases} \quad (4.58)$$

Let us establish the properties of the graph determinant.

**Theorem 4.14** *For a matrix field  $M$ ,  $\det(\mathcal{T}_{\mathbf{g}} M) = \mathcal{T}_{\mathbf{g}}(\det M)$ .*

*Proof.* Let us first prove this for a strut matrix field (4.43). Let us assume for simplicity that all coordinates are root (a general case is easy to consider, since struts do not mix root and cartan coordinates, so that a general strut matrix field has a block-diagonal form (4.44)). A weight system  $\mathcal{T}_{\mathfrak{g}}$  applied to a root strut matrix field

$$M = \sum_{i,j=1}^L l_{ji} \partial_{x_j} - \dots - dx_i \quad (4.59)$$

takes value in  $\mathcal{Q}((S\mathfrak{h})^{\otimes |\mathcal{L}'_{\mathfrak{c}}|}) \otimes \text{End} \left( \bigoplus_{j=1}^L \mathfrak{r}_j \right)$ , where  $\mathfrak{r}_j$  are  $L$  copies of the space  $\mathfrak{r}$ . In fact, the weight system of a matrix field (4.59) is diagonal with respect to the root spaces  $\mathfrak{r}_{\vec{\lambda}}$ , so

$$\mathcal{T}_{\mathfrak{g}} M = \sum_{\vec{\lambda} \in \Delta_{\mathfrak{g}}} (\mathcal{T}_{\mathfrak{g}} M)_{\vec{\lambda}}, \quad (\mathcal{T}_{\mathfrak{g}} M)_{\vec{\lambda}} = ||l_{ij}(\vec{\lambda})|| \in \mathcal{Q}((S\mathfrak{h})^{\otimes |\mathcal{L}'_{\mathfrak{c}}|}) \otimes \text{End} \left( \bigoplus_{j=1}^L \mathfrak{r}_{\vec{\lambda},j} \right). \quad (4.60)$$

As a result,

$$\det(\mathcal{T}_{\mathfrak{g}} M) = \prod_{\vec{\lambda} \in \Delta_{\mathfrak{g}}} \det(\mathcal{T}_{\mathfrak{g}} M)_{\vec{\lambda}} = \prod_{\vec{\lambda} \in \Delta_{\mathfrak{g}}} \det ||l_{ij}(\vec{\lambda})||. \quad (4.61)$$

A combination of the definition of determinant (4.45) and the formula (3.76) for the application of  $\mathcal{T}_{\mathfrak{g}}$  to  $\mathcal{QH}(\vec{\lambda})$  yields the same expression in view of eq.(4.47) applied to  $||l_{ij}(\vec{\lambda})||$ . Thus we proved the theorem for strut matrix fields.

A general definition of a determinant is based on formulas (4.51) and (4.58). The application of weight system  $\mathcal{T}_{\mathfrak{g}}$  converts eq.(4.51) into eq.(4.47) and converts eq.(4.58) into a multiplicativity of the ordinary determinant. Therefore  $\mathcal{T}_{\mathfrak{g}}$  ‘commutes’ with these defining equations, and hence it commutes with graph determinant.  $\square$

**Theorem 4.15** *The determinant (4.58) is multiplicative.*

*Proof.* Let  $M_1$  and  $M_2$  be non-degenerate matrix fields. Then

$$\det(M_1 \times M_2) = \det(M_{1,\text{str}} \times M_{2,\text{str}}) \det((M_{1,\text{str}} \times M_{2,\text{str}})^{-1} \times M_1 \times M_2). \quad (4.62)$$

Determinant of strut matrix fields is multiplicative, so

$$\det(M_{1,\text{str}} \times M_{2,\text{str}}) = (\det M_{1,\text{str}}) (\det M_{2,\text{str}}). \quad (4.63)$$

As for the second factor in the *r.h.s.* of eq.(4.62),

$$\begin{aligned}
& \det \left( (M_{1,\text{str}} \times M_{2,\text{str}})^{-1} \times M_1 \times M_2 \right) \\
&= \det \left( M_{2,\text{str}}^{-1} \times M_{1,\text{str}}^{-1} \times M_1 \times M_2 \right) \\
&= \det \left[ \left( M_{2,\text{str}}^{-1} \times (M_{1,\text{str}}^{-1} \times M_1) \times M_{2,\text{str}} \right) \times (M_{2,\text{str}}^{-1} \times M_2) \right] \\
&= \det \left( M_{2,\text{str}}^{-1} \times (M_{1,\text{str}}^{-1} \times M_1) \times M_{2,\text{str}} \right) \det(M_{2,\text{str}}^{-1} \times M_2) \\
&= \det(M_{1,\text{str}}^{-1} \times M_1) \det(M_{2,\text{str}}^{-1} \times M_2). \tag{4.64}
\end{aligned}$$

Here the third equality is due to Lemma 4.12 and the fourth equality is due to Lemma 4.13. The multiplicativity (4.42) follows from this equation and the definition (4.58).  $\square$

Since an exponential is non-zero, then it follows from eq.(4.58) that a matrix field  $M_{\text{str}}$  is non-degenerate iff  $\det M_{\text{str}} \neq 0$ .

**Theorem 4.16** *A matrix field  $M$  can be inverted iff it is non-degenerate.*

*Proof.* The easiest way to prove this theorem is to deduce it from Theorem 4.4. Indeed, one just has to declare the coordinate colors  $\mathbf{x}$  to be parameters and call  $\partial_{\mathbf{x}}$  and  $d\mathbf{x}$  new outgoing and incoming coordinates.  $\square$

An explicit formula for the inverse matrix field is

$$M^{-1} = M_{\text{str}}^{-1} \times \sum_{n=0}^{\infty} (-1)^n (M_{\text{str}}^{-1} \times M)^{\times n}. \tag{4.65}$$

**Remark 4.17** A determinant of a non-degenerate matrix field  $M$  is invertible as an element of the graph algebra. Indeed, eqs. (4.58), (4.51) and (4.45) define it as a product of two functions, both of which are exponentials, and an inverse of an exponential function is constructed by adding a minus sign to the exponent.

**Theorem 4.18 (derivative of a determinant)** *If a matrix field  $M(t)$  depends on a  $\mathbb{Q}$ -parameter  $t$  and  $M(t)$  is non-degenerate, then*

$$\partial_t \det M = \det M \operatorname{Tr}(M^{-1} \partial_t M). \tag{4.66}$$

*Proof.* Let us evaluate  $\det M(t + dt)$  to the linear order in  $dt$ :

$$\begin{aligned}
\det M(t + dt) &= \det \left[ M(t) + \left( \partial_t M(t) \right) dt \right] = \det M(t) \det \left[ I + \left( M^{-1}(t) \partial_t M(t) \right) dt \right] \\
&= \left( \det M(t) \right) \left[ 1 + \operatorname{Tr} \left( M^{-1}(t) \partial_t M(t) \right) dt \right], \tag{4.67}
\end{aligned}$$

where the last equation follows from eq.(4.51). Equation (4.66) follows from eq.(4.67).  $\square$

A derivative of a diffeomorphism  $\mathbf{y} = F(\mathbf{x})$  is a matrix field  ${}^y F'(\mathbf{x})$ . We define its determinant by first replacing the colors  $\partial_{\mathbf{y}}$  by the corresponding colors  $\partial_{\mathbf{x}}$  and then applying the definition (4.58).

It is obvious that the definition (4.58) of a determinant commutes with the maps  $f_{\text{trv}}$  and  $f_{\mathcal{Q}}$  of (3.63). However the commutativity with  $f_{\mathcal{QH}}$  is not immediately obvious and requires a special consideration.

**Theorem 4.19** *Let  ${}^x M(\mathbf{x}; \mathbf{a})$  be a matrix field in  $\mathcal{B}_{\mathfrak{L}}^{\mathbb{C}}$ , which depends on Cartan parameters  $\mathbf{a}$ . Suppose that  $\mathbf{a} \in \mathfrak{L}'_{\mathbb{C}}$  and denote  ${}^x M'(\mathbf{x}) = f_{\mathcal{QH}}({}^x M(\mathbf{x}; \mathbf{a}))$  (cf. (4.3), we used notation  ${}^x M'$  for the image of  ${}^x M$  in  $\mathcal{D}_{\mathfrak{L}, \mathfrak{L}'_{\mathbb{C}}}^{\mathbb{C}}$  instead of  ${}^x M$  in order to avoid any mix up). Suppose that  ${}^x M$  is non-degenerate. Then  ${}^x M'$  is also non-degenerate and*

$$\det {}^x M'(\mathbf{x}) = f_{\mathcal{QH}}(\det {}^x M(\mathbf{x}; \mathbf{a})). \quad (4.68)$$

*Proof.* Let us drop  ${}^x$  from our notations of matrix fields. It is easy to see that the strut graphs of  $M'$  come from the haircomb graphs (Fig. 1.1) of  $M$  whose vertical legs are colored by  $\mathbf{a}$  and horizontal legs are colored by  $\partial_{\mathbf{x}}$  and  $d\mathbf{x}$ . Therefore, if following eqs.(4.43) and (4.44) we denote

$$(M'(\mathbf{x}))_{\text{str}} = \sum_{i,j=1}^L l_{ij}(\mathbf{a}) \partial_{x_i} \text{ --- } dx_j, \quad l_{ij}(\mathbf{a}) \in \mathbb{Q}[[\mathbf{a}]], \quad (4.69)$$

then

$$(M(\mathbf{x}; \mathbf{a}))_{\text{str}} = \sum_{i,j=1}^L l_{ij}(\mathbf{a})|_{\mathbf{a}=0} \partial_{x_i} \text{ --- } dx_j. \quad (4.70)$$

As a result,

$$\det M_{\text{str}} = \det M'_{\text{str}}|_{\mathbf{a}=0}, \quad (4.71)$$

and non-degeneracy of  $M(\mathbf{x}; \mathbf{a})$  indeed implies the non-degeneracy of  $M'(\mathbf{x})$ .

Since the determinant is multiplicative in both  $\mathcal{B}_{\mathfrak{L}}^{\mathbb{C}}$  and  $\mathcal{D}_{\mathfrak{L}, \mathfrak{L}'_{\mathbb{C}}}^{\mathbb{C}}$ , we can prove eq. (4.68) separately for  $M'_{\text{str}}$  and for  $(M'_{\text{str}})^{-1} M'$ . Since

$$\det ||l_{ij}(\mathbf{a})|| = \det ||l_{ij}(\mathbf{a})|| \Big|_{\mathbf{a}=0} \exp \left( \text{Tr} \log(||l_{ij}(\mathbf{a})||^{-1}) \Big|_{\mathbf{a}=0} ||l_{ij}(\mathbf{a})|| \right), \quad (4.72)$$

then

$$f_{\mathcal{QH}}^{-1}(\det M'_{\text{str}}(\mathbf{x})) = \det f_{\mathcal{QH}}^{-1}(M'_{\text{str}}(\mathbf{x}; \mathbf{a})). \quad (4.73)$$

Also since the strut parts of both  $(M'_{\text{str}}(\mathbf{x}))^{-1}M'(\mathbf{x})$  and  $f_{\mathcal{QH}}^{-1}((M'_{\text{str}}(\mathbf{x}))^{-1}M'(\mathbf{x}))$  are identity matrices, then their determinants are calculated just by the formula (4.48) and hence these determinants are also related by  $f_{\mathcal{QH}}$ .  $\square$

**Remark 4.20** Equation (4.71) indicates that it may happen that  $\det M(\mathbf{x}; \mathbf{a}) \equiv 0$ , while  $\det M'(\mathbf{x}) \not\equiv 0$ . This means that the definition of a determinant in  $\mathcal{QD}_{\mathfrak{L}, \mathfrak{L}'_c}^C$  is finer than in  $\mathcal{B}_{\mathfrak{L}}^C$ , and the reason for this is that a notion of a strut in  $\mathcal{QD}_{\mathfrak{L}, \mathfrak{L}'_c}^C$  is wider than that in  $\mathcal{B}_{\mathfrak{L}}^C$ , since the former includes the haircomb graphs of  $\mathcal{B}_{\mathfrak{L}}^C$  with legs  $\mathfrak{L}'_c$  legs.

## 4.6 Integration measure

Now we can define an integration measure. As a graph with labeled legs, an *integration measure* is the same as a function. The only difference is in the action of substitutions. A diffeomorphism  $\mathbf{y} = F(\mathbf{x})$  acts on an integration measure  $G(\mathbf{x})$  as

$$G(\mathbf{y}) \xrightarrow{F} F^*G(\mathbf{x}) = G(F(\mathbf{x})) \det F'(\mathbf{x}). \quad (4.74)$$

Chain rule and the multiplicativity property of the determinant indicate that this is a genuine (contragradient) group action. It is easy to verify that a graph algebra product  $f(\mathbf{x})G(\mathbf{x})$  of a function  $f(\mathbf{x})$  and an integration measure  $G(\mathbf{x})$  is an integration measure.

**Theorem 4.21 (Lie derivative of integration measure)** *The action of a vector field  $\xi(\mathbf{x})$  on an integration measure  $G(\mathbf{x})$  is described by a formula*

$$L_{\xi}G = \nabla_{\xi}G + G \operatorname{div}_{\mathbf{x}} \xi. \quad (4.75)$$

*Proof.* We define a family of diffeomorphisms  $\mathbf{y} = F(\mathbf{x}, t)$  by a formula  $F(\mathbf{x}, t) = \operatorname{Id} + t \tilde{\xi}(\mathbf{x})$ , where  $\mathbf{y} = \tilde{\xi}(\mathbf{x})$  is the vector field  $\xi$  converted into a map by replacing the  $\partial_{\mathbf{x}}$  colors with  $\mathbf{y}$ . Then by definition

$$L_{\xi}G = \partial_t \left( G(F(\mathbf{x}, t)) \det F'(\mathbf{x}, t) \right) \Big|_{t=0}. \quad (4.76)$$

As we already know,

$$\partial_t \left( G(F(\mathbf{x}, t)) \right) \Big|_{t=0} = \nabla_{\xi}G. \quad (4.77)$$

At the same time,  $F' = I + t \nabla \xi$ , so  $\det F'(x, 0) = 1$  and according to eqs.(4.66) and (4.41),

$$\partial_t \det F' \Big|_{t=0} = \operatorname{Tr}(\nabla \xi) = \operatorname{div}_{\mathbf{x}} \xi. \quad (4.78)$$

The formula (4.75) follows easily from eqs.(4.76), (4.77) and (4.78).  $\square$

## 4.7 A formal gaussian integral

Now we define the integral. The only integral that we can define is the gaussian or the stationary phase one, because, as it turns out, its definition can be given purely in terms of combinatorics of the integrand without any references to Riemann sums. We will use the names ‘gaussian’ and ‘stationary phase’ as synonyms, although there is a slight difference between them: the stationary phase integral allows infinite formal power series as a preexponential factor.

A gaussian integral for a graph algebra  $\mathcal{B}$  has been already defined in [3]. Its definition can be transferred *verbatim* to any other graph algebra, and we are going to do it. The only difference is that the paper [3] neglected the 1-loop determinant, because it played a rather trivial role in those calculations. However, we must reinstate it, since it participates in the proof of topological invariance of  $U(1)$ -RCC invariant and also contributes to the Alexander polynomial.

We say that a function, a tensor field or an integration measure  $P$  is  $\mathbf{x}$ -substantial if there exists a positive number  $N$  such that  $P$  can be presented as a (possibly infinite) linear combination of elements of spaces  $\mathcal{QH}(D)$  such that each graph  $D$  has at most  $N$  struts  $x_i \text{ --- } x_j$ ,  $1 \leq i, j \leq L$ , among its connected components. In short,  $P$  has a polynomial dependence on these struts. A function, a tensor field or an integration measure  $G$  is called *stationary phase* (SP), if it has a form

$$G(\mathbf{x}) = e^{\frac{1}{2}Q(\mathbf{x})}P(\mathbf{x}), \quad (4.79)$$

where

$$Q(\mathbf{x}) = \sum_{i,j=1}^L l_{ij} \ x_i \text{ --- } x_j, \quad (4.80)$$

while  $P(\mathbf{x})$  is  $\mathbf{x}$ -substantial (we put an extra  $1/2$  in the exponent for future convenience). We choose the coefficients  $l_{ij}$  so that they satisfy a condition

$$l_{ij}^* = l_{ji}. \quad (4.81)$$

We say that  $G$  is *stationary phase non-degenerate* (SPND) if the matrix  $||l_{ij}||$  is non-degenerate. We associate to  $Q(\mathbf{x})$  an ‘inverse quadratic form’

$$Q^{-1}(\mathbf{x}) = \sum_{i,j=1}^L l_{ij}^{-1} \ x_i \text{ --- } x_j \quad (4.82)$$

( $||l_{ij}^{-1}|| = ||l_{ij}||^{-1}$ ) and a matrix field

$$\hat{Q} = \sum_{i,j=1}^L l_{ji} \partial_{x_i} \text{---} dx_j . \quad (4.83)$$

If  $G(\mathbf{x})$  of eq. (4.79) is SPND, then following [3] we define its *formal gaussian* (i.e. stationary phase) integral by the formula

$$\int_{\text{FG}} d\mathbf{x} e^{\frac{1}{2}Q(\mathbf{x})} P(\mathbf{x}) = \left( \det \hat{Q} \right)^{-1/2} \left( P(\mathbf{x}) \text{---} b_{\mathbf{x},\mathbf{x};\text{LR}} \text{---} e^{-\frac{1}{2}Q^{-1}(\mathbf{x})} \right), \quad (4.84)$$

The SPND condition on  $G(\mathbf{x})$  guarantees that the gluing in eq.(4.84) is well-defined, that is, each particular graph in the second brackets can be constructed by gluing the legs of  $P(\mathbf{x})$  and  $e^{-\frac{1}{2}Q^{-1}(\mathbf{x})}$  in only finitely many ways. The difference between our definition and that of [3] is in the ‘1-loop’ determinant factor  $\left( \det \hat{Q} \right)^{-1/2}$  which was dropped in [3].

**Remark 4.22** Condition (4.81) implies that  $\det ||l_{ij}^*|| = \det ||l_{ij}||$ , hence  $\det \hat{Q}$  can be written simply as

$$\det \hat{Q} = \exp \left( \log(\det ||l_{ij}||_{\text{r}}) \text{---} \text{---} + \log(\det ||l_{ij}||_{\text{c}}) \text{---} \text{---} \right) \quad (4.85)$$

(cf. eq.(4.45)).

## 4.8 Properties of the integral

Now let us establish the basic properties of integrals for the definition (4.84). Most of the work in this direction has been already done in [3].

First of all, let us use the weight system  $\mathcal{T}_{\mathfrak{g}}$  in order to relate the graph formula (4.84) to the usual calculus definition of the stationary phase integral. Assume for simplicity that  $G(\mathbf{x})$  and  $P(\mathbf{x})$  are both either functions or integration measures. Then, according to (4.1) we may think of  $\mathcal{T}_{\mathfrak{g}}G$  and  $\mathcal{T}_{\mathfrak{g}}P$  as formal power series of  $\vec{\mathbf{x}}$  (let us ignore their dependence on parameters  $\vec{\mathbf{a}}$ ). Usually in calculus the preexponential factor  $\mathcal{T}_{\mathfrak{g}}P$  would be presented as a power series in  $\hbar$ , so that the whole integral would be well-defined as such a power series. First, let us not commit ourselves to any particular way of inserting  $\hbar$  into  $P(\mathbf{x})$  or  $Q(\mathbf{x})$  and consider the case of a ‘polynomial’  $\mathcal{T}_{\mathfrak{g}}P$ .

**Lemma 4.23** *Suppose that there exists a finite set of graphs  $\mathbf{D}_P$ , such that*

$$P(\mathbf{x}) \in \bigoplus_{D \in \mathbf{D}_P} \mathcal{QH}(D). \quad (4.86)$$



Then  $\mathcal{T}_{\mathfrak{g}}P$  is a polynomial of  $\vec{x}$  and

$$\mathcal{T}_{\mathfrak{g}} \left( \int_{\text{FG}} d\mathbf{x} G(\mathbf{x}) \right) = (-2\pi)^{-\frac{1}{2} \sum_{j=1}^L \dim \mathfrak{v}_j^*} \int d\vec{x} \mathcal{T}_{\mathfrak{g}} G(\vec{x}). \quad (4.87)$$

*Proof.* Obviously,  $\mathcal{T}_{\mathfrak{g}}P$  is a polynomial of  $\mathbf{x}$  of degree which is equal to the maximum number of  $\mathbf{x}$  legs in the graphs of  $\mathbf{D}_P$ .  $\mathcal{T}_{\mathfrak{g}}Q$  is a quadratic form

$$\mathcal{T}_{\mathfrak{g}}Q(\vec{x}) = \sum_{i,j=1}^L l_{ij} (\vec{x}_i, \vec{x}_j), \quad (4.88)$$

so if we assume that this quadratic form is negative-definite, then the *r.h.s.* of eq.(4.84) is a well-defined gaussian integral with a polynomial prefactor. This integral is calculated with the help of an explicit formula

$$(-2\pi)^{-\frac{1}{2} \sum_{j=1}^L \dim \mathfrak{v}^*} \int d\vec{x} e^{\frac{1}{2} \mathcal{T}_{\mathfrak{g}}Q(\vec{x})} \mathcal{T}_{\mathfrak{g}}P(\vec{x}) = \left( \det ||l_{ij}|| \right)^{-1/2} \mathcal{T}_{\mathfrak{g}}P(\partial_{\vec{x}}) e^{-\frac{1}{2} \mathcal{T}_{\mathfrak{g}}Q^{-1}(\vec{x})} \Big|_{\vec{x}=0} \quad (4.89)$$

It is easy to see that the *r.h.s.* of this formula is equal to  $\mathcal{T}_{\mathfrak{g}}$  applied to the *r.h.s.* of eq.(4.84).  $\square$

**Remark 4.24** The prefactor  $(-2\pi)^{-\frac{1}{2} \sum_{j=1}^L \dim \mathfrak{v}^*}$  in equation (4.89) indicates that eq.(4.84) defines  $\int_{\text{FG}} \frac{d\mathbf{x}}{(-2\pi)^{1/2}}$  rather than  $\int_{\text{FG}} d\mathbf{x}$ . Alternatively, we could remove that prefactor by replacing  $\left( \det \hat{Q} \right)^{-1/2}$  in the *r.h.s.* of eq.(4.84) with  $\left( \det(-2\pi \hat{Q}) \right)^{-1/2}$ .

In order to bring in  $\hbar$ , we introduce a weight system  $\mathcal{T}_{\mathfrak{g}}^{2,\hbar}$ , which acts in the same way as  $\mathcal{T}_{\mathfrak{g}}^{1,\hbar}$ , except that it multiplies the graphs by  $\hbar^{\deg_2 D}$  rather than by  $\hbar^{\deg_1 D}$ .

**Theorem 4.25** For a SPND function (4.79)

$$\mathcal{T}_{\mathfrak{g}}^{2,\hbar} \left( \int_{\text{FG}} d\mathbf{x} G(\mathbf{x}) \right) = (-2\pi\hbar)^{-\frac{1}{2} \sum_{j=1}^L \dim \mathfrak{v}_j^*} \int d\vec{x} \mathcal{T}_{\mathfrak{g}}^{2,\hbar} G(\vec{x}). \quad (4.90)$$

*Proof.* Let  $P_N$  denote the part of  $P$  which contains only the contributions of the graphs  $D$  with  $\deg_2 D \leq N$ . Since there are only finitely many graphs with such property for a given  $N$ , then  $P_N$  satisfies the property (4.86) and eq.(4.87) holds for  $G_N = e^{\frac{1}{2} Q(\mathbf{x})} P_N$ . Now eq.(4.90) for  $G_N$  follows from eq.(4.87) and from a relation

$$\mathcal{T}_{\mathfrak{g}}^{2,\hbar} \left( P_N(\mathbf{x}) \mid_{\mathbf{x}, \mathbf{x}; \text{LR}} e^{-\frac{1}{2} Q^{-1}(\mathbf{x})} \right) = \mathcal{T}_{\mathfrak{g}}^{2,\hbar} P_N(\partial_{\vec{x}}) e^{-\frac{\hbar}{2} \mathcal{T}_{\mathfrak{g}} Q^{-1}(\vec{x})} \Big|_{\vec{x}=0}, \quad (4.91)$$

which is established by an easy counting of the powers of  $\hbar$  (note that gluing a strut to two legs of a graph  $D$  reduces  $\deg_2 D$  by 1). Equation (4.90) for  $G$  follows by taking a limit  $N \rightarrow \infty$ , which is well-defined on both sides of this equation.  $\square$

**Theorem 4.26 (parity invariance)** *If  $G(\mathbf{x})$  is SPND, then*

$$\int_{\text{FG}} d\mathbf{x} G(-\mathbf{x}) = \int_{\text{FG}} d\mathbf{x} G(\mathbf{x}) \quad (4.92)$$

*Proof.* This theorem was proved in [3]. We can use this proof verbatim, since changing the sign of  $\mathbf{x}$  does not change the strut part and therefore does not change the 1-loop determinant.  $\square$

**Theorem 4.27 (Fubini's theorem)** *Consider an SP integrand  $G(\mathbf{x}, \mathbf{y})$  which depends on two sets of coordinates  $\mathbf{x}$  and  $\mathbf{y}$ . Suppose that both the full quadratic form  $Q(\mathbf{x}, \mathbf{y})$  and its  $\mathbf{x}$  —  $\mathbf{x}$  part  $Q_x(\mathbf{x})$  are non-degenerate. Then  $\int_{\text{FG}} d\mathbf{x} G(\mathbf{x}, \mathbf{y})$  is a SPND integrand for  $\mathbf{y}$  and*

$$\int_{\text{FG}} d\mathbf{x} d\mathbf{y} G(\mathbf{x}, \mathbf{y}) = \int_{\text{FG}} d\mathbf{y} \left( \int_{\text{FG}} d\mathbf{x} G(\mathbf{x}, \mathbf{y}) \right). \quad (4.93)$$

*Proof.* The theorem was proved in [3] for the case when 1-loop determinants were neglected. Accounting for the determinants is easy. If we split the matrix  $||l_{ij}||$  of  $Q(\mathbf{x}, \mathbf{y})$  into  $\mathbf{x}$  and  $\mathbf{y}$  blocks as

$$||l_{ij}|| = \left( \begin{array}{c|c} A & B \\ \hline B^T & C \end{array} \right), \quad (4.94)$$

then it is easy to see that the corresponding  $y$ -matrix of the exponent of  $\int_{\text{FG}} d\mathbf{x} G(\mathbf{x}, \mathbf{y})$  is  $C - B^T A^{-1} B$  and the determinant part of Fubini's formula follows from the well-known identity  $\det ||l_{ij}|| = \det A \det(C - B^T A^{-1} B)$ .  $\square$

**Remark 4.28** The condition that the  $\mathbf{x}$  —  $\mathbf{x}$  part  $Q_x(\mathbf{x})$  of  $Q(\mathbf{x}, \mathbf{y})$  should be non-degenerate, does not restrict the applicability of Fubini's theorem too much. If needed, one can consider a deformed quadratic form  $Q(\mathbf{x}, \mathbf{y}; t)$  which depends on a parameter  $t$  in such a way that  $Q(\mathbf{x}, \mathbf{y}; 0) = Q(\mathbf{x}, \mathbf{y})$  and  $Q_x(\mathbf{x}; t)$  is non-degenerate for general values of  $t$ . Then one can apply Fubini's theorem for a general value of  $t$ , obtain all necessary results and at the very end set  $t = 0$ .

If  $G(\mathbf{x})$  is an SPND integration measure and  $\xi$  is a  $\mathbf{x}$ -substantial vector field, then  $\nabla_\xi G$ ,  $G \operatorname{div}_\mathbf{x} \xi$  and  $L_\xi G$  are all SPND (the latter also being an integration measure).

**Theorem 4.29 (integration by parts)** *If  $G(\mathbf{x})$  is an SPND integration measure and  $\xi(\mathbf{x})$  is a  $\mathbf{x}$ -substantial vector field, then*

$$\int_{\text{FG}} d\mathbf{x} L_\xi G = 0. \quad (4.95)$$

*Proof.* The claim of this theorem obviously does not depend on the inclusion of the 1-loop determinant factor in eq.(4.84), so the proof of [3] applies without modifications.  $\square$

This theorem has a simple corollary

**Corollary 4.30** *Let  $\Xi(\mathbf{x})$  be an SPND vector field and  $\mu(\mathbf{x})$  be an  $\mathbf{x}$ -substantial integration measure. If  $L_{\Xi}\mu = 0$ , then*

$$\int_{\text{FG}} d\mathbf{x} (\text{div}_{\mathbf{x}} \Xi) \mu = 0. \quad (4.96)$$

If an SPND integrand  $G(\mathbf{x}; t)$  depends on a  $\mathbb{Q}$ -parameter  $t$ , then  $\partial_t G(\mathbf{x}; t)$  is also an SPND integrand.

**Theorem 4.31 (derivative of an integral over a parameter)** *If an SPND integrand  $G(\mathbf{x}; t)$  depends on a  $\mathbb{Q}$ -parameter  $t$ , then*

$$\partial_t \left( \int_{\text{FG}} d\mathbf{x} G(\mathbf{x}; t) \right) = \int_{\text{FG}} d\mathbf{x} \partial_t G(\mathbf{x}; t). \quad (4.97)$$

*Proof.* Let  $G(\mathbf{x}; t) = e^{Q(\mathbf{x}; t)} P(\mathbf{x}; t)$ . Since

$$\partial_t G(\mathbf{x}; t) = e^{Q(\mathbf{x}; t)} \left( \frac{1}{2} P(\mathbf{x}; t) \partial_t Q(\mathbf{x}; t) + \partial_t P(\mathbf{x}; t) \right), \quad (4.98)$$

then proving eq.(4.97) amounts to showing the following:

$$\begin{aligned} & \partial_t \left[ \left( \det \hat{Q} \right)^{-1/2} \left( P \ \flat_{\mathbf{x}, \mathbf{x}; \text{LR}} \ e^{-\frac{1}{2} Q^{-1}} \right) \right] \\ &= \frac{1}{2} \left( \det \hat{Q} \right)^{-1/2} \left[ \left( (P \partial_t Q) \ \flat_{\mathbf{x}, \mathbf{x}; \text{LR}} \ e^{-\frac{1}{2} Q^{-1}} \right) + 2 \left( \partial_t P \ \flat_{\mathbf{x}, \mathbf{x}; \text{LR}} \ e^{-\frac{1}{2} Q^{-1}} \right) \right]. \end{aligned} \quad (4.99)$$

We calculate explicitly the *l.h.s.* of this equation

$$\begin{aligned} & \partial_t \left[ \left( \det \hat{Q} \right)^{-1/2} \left( P \ \flat_{\mathbf{x}, \mathbf{x}; \text{LR}} \ e^{-\frac{1}{2} Q^{-1}} \right) \right] \\ &= \frac{1}{2} \left( \det \hat{Q} \right)^{-1/2} \left[ -\text{Tr}(\hat{Q}^{-1} \partial_t \hat{Q}) \left( P \ \flat_{\mathbf{x}, \mathbf{x}; \text{LR}} \ e^{-\frac{1}{2} Q^{-1}} \right) - P \ \flat_{\mathbf{x}, \mathbf{x}; \text{LR}} \left( (\partial_t Q^{-1}) e^{-\frac{1}{2} Q^{-1}} \right) \right. \\ & \quad \left. + 2 \left( \partial_t P \ \flat_{\mathbf{x}, \mathbf{x}; \text{LR}} \ e^{-\frac{1}{2} Q^{-1}} \right) \right], \end{aligned} \quad (4.100)$$

where  $\partial_t Q^{-1}$  can be determined with the help of the formula

$$\widehat{\partial_t Q^{-1}} = -\hat{Q}^{-1} \times \partial_t \hat{Q} \times \hat{Q}^{-1}. \quad (4.101)$$

The last terms in the *r.h.s.* of eqs.(4.99) and (4.100) coincide, so in order to prove eq.(4.99) we have to show that

$$\begin{aligned} & (P \partial_t Q) \mathbin{\lrcorner}_{\mathbf{x}, \mathbf{x}; \text{LR}} e^{-\frac{1}{2}Q^{-1}} \\ &= -\text{Tr}(\hat{Q}^{-1} \partial_t \hat{Q}) \left( P \mathbin{\lrcorner}_{\mathbf{x}, \mathbf{x}; \text{LR}} e^{-\frac{1}{2}Q^{-1}} \right) - P \mathbin{\lrcorner}_{\mathbf{x}, \mathbf{x}; \text{LR}} \left( (\partial_t Q^{-1}) e^{-\frac{1}{2}Q^{-1}} \right) \end{aligned} \quad (4.102)$$

Indeed, consider the gluing in the *l.h.s.* of this equation. The legs of  $\partial_t Q$  can either be glued to the same strut of  $e^{-\frac{1}{2}Q^{-1}}$  or to the different struts. In the first case, the first term of the *r.h.s.* of eq.(4.102) is reproduced. In the second case, the other two legs of the two struts of  $e^{-\frac{1}{2}Q^{-1}}$  which are glued to the strut  $\partial_t Q$  will be glued to the legs of  $P$  and in view of eq.(4.101) this reproduces exactly the second term of the *r.h.s.* of eq.(4.102).  $\square$

Finally we want to prove the invariance of the integral (4.84) under diffeomorphisms if  $G(x)$  is an SPND integration measure. We have to limit ourselves to such diffeomorphisms  $F$  that  $F^*G$  is again SPND. Recall that we call an element of a graph algebra *narrow* if it can be presented as a linear combination of connected graphs. It is easy to see that narrow diffeomorphisms form a group which we denote as  $\text{Diff}_L^{\text{nrw}}$ .

**Lemma 4.32** *The groups  $\text{Diff}_L$  and  $\text{Diff}_L^{\text{nrw}}$  are connected.*

*Proof.* It is easy to connect a diffeomorphism  $F$  to its linear part: one can use a path

$$F(t) = F_{\text{lin}} + t(F - F_{\text{lin}}), \quad 0 \leq t \leq 1. \quad (4.103)$$

Since  $(F(t))_{\text{lin}} = F_{\text{lin}}$ , then all maps  $\mathbf{y} = F(\mathbf{x}; t)$  are non-degenerate and thus belong to  $\text{Diff}_L$ . If  $F$  is narrow, then all diffeomorphisms  $F(t)$  are also narrow.  $F_{\text{lin}}$ , in its turn, is connected to the identity diffeomorphism  $\text{Id}$ , because the group  $GL$  is connected.  $\square$

**Theorem 4.33 (diffeomorphism invariance of an integral of a measure)** *If  $G(\mathbf{y})$  is an SPND integration measure and  $\mathbf{y} = F(\mathbf{x})$  is narrow, then  $F^*G(\mathbf{x})$  is also SPND and*

$$\int_{\text{FG}} d\mathbf{x} F^*G(\mathbf{x}) = \int_{\text{FG}} d\mathbf{y} G(\mathbf{y}). \quad (4.104)$$

*Proof.* Consider eq.(4.74) for  $F^*G(\mathbf{x})$ . It is easy to check that  $G(F(\mathbf{x}))$  is SPND. Also it follows from eqs.(4.58), (4.80) and (4.45) that if  $\mathbf{y} = F(\mathbf{x})$  is narrow, then the graphs of the determinant  $\det F'(\mathbf{x})$  of eq.(4.74) do not contain connected tree subgraphs, so  $\det F'(\mathbf{x})$  is  $\mathbf{x}$ -substantial and  $F^*G(\mathbf{x})$  is SPND.

Since, according to Lemma 4.32, the group  $\text{Diff}_L^{\text{nrw}}$  is connected, then there exists a 1-parametric family of narrow diffeomorphisms  $F(t)$ , such that  $F(t)|_{t=0} = \text{Id}$  and  $F(t)|_{t=1} = F(\mathbf{x})$ . Then eq.(4.104) follows from the following calculation

$$\partial_t \left( \int_{\text{FG}} F^*(t) G(\mathbf{x}) \right) = \int_{\text{FG}} \partial_t F^*(t) G(\mathbf{x}) = \int_{\text{FG}} L_{(\partial_t F(t)) \circ F^{-1}(t)} \left( F^*(t) G(\mathbf{x}) \right) = 0, \quad (4.105)$$

which is based on eqs.(4.97), (4.24) and (4.95), the latter being applicable since the vector field  $(\partial_t F(t)) \circ F^{-1}(t)$  is narrow and therefore  $\mathbf{x}$ -substantial.  $\square$

**Corollary 4.34** *If  $\mathbf{y} = F(\mathbf{x})$  is a narrow diffeomorphism and  $\mu(\mathbf{x})$  is an  $\mathbf{x}$ -substantial and  $F$ -invariant integration measure, then for any SPND function  $G(\mathbf{y})$*

$$\int_{\text{FG}} d\mathbf{x} G(F(\mathbf{x})) \mu(\mathbf{x}) = \int_{\text{FG}} d\mathbf{y} G(\mathbf{y}) \mu(\mathbf{y}). \quad (4.106)$$

*Proof.* Apply eq.(4.104) to the SPND integration measure  $G(\mathbf{y})\mu(\mathbf{y})$ .  $\square$

Obviously, the definition (4.84) commutes with the first injection  $f_{\text{trv}}$  of (3.63). However, its commutativity with  $f_{\mathcal{QH}}$  is less obvious, since as we have already discussed, the notion of a strut exponent is different in  $\mathcal{B}_{\mathcal{E}}^{\text{C}}$  and  $\mathcal{D}_{\mathcal{E}, \mathcal{E}_c}^{\text{C}}$ . In order to prove it, we have to establish that eq.(4.84) sometimes holds even in the case when  $Q(\mathbf{x})$  is not purely strut.

First, we introduce a few general definitions. A function  $Q(\mathbf{x})$  is called *quadratic* in  $\mathbf{x}$ , if every graph of  $Q(\mathbf{x})$  has exactly two  $\mathbf{x}$  legs ( $Q(\mathbf{x})$  may also depend on other coordinates and parameters). Suppose that a matrix field  ${}^x M$  does not depend on  $\mathbf{x}$ , that is, its graphs do not have  $\mathbf{x}$  legs. Then we can associate to  ${}^x M$  a quadratic (in  $\mathbf{x}$ ) function  $(M)_{\text{qf}}$  by replacing the  $\partial_{\mathbf{x}}$  and  $d\mathbf{x}$  labels in the graphs of  ${}^x M$  with corresponding labels  $\mathbf{x}$ . Conversely, if  $Q(\mathbf{x})$  is a quadratic function of  $\mathbf{x}$ , then we can associate to it a unique matrix field  $\hat{Q}$  which satisfies two properties:  $(\hat{Q})_{\text{qf}} = Q(\mathbf{x})$  and  $\hat{Q}^{\text{T}} = \hat{Q}$  (that is,  $\hat{Q}$  is symmetric). Obviously, our definition is consistent with eq.(4.83).

Let  $Q(\mathbf{x})$  be a quadratic function of  $\mathbf{x}$ . As usual,  $Q_{\text{str}}(\mathbf{x})$  denotes its strut part.

**Theorem 4.35** *Suppose that  $Q_{\text{str}}(\mathbf{x})$  is non-degenerate and that  $P(\mathbf{x})$  is  $\mathbf{x}$ -substantial. Then  $e^{\frac{1}{2} Q(\mathbf{x})} P(\mathbf{x})$  is SPND and eq. (4.84) holds, where  $Q^{-1}(\mathbf{x})$  is defined as  $Q^{-1}(\mathbf{x}) = (\hat{Q}^{-1})_{\text{qf}}$ .*

*Proof.* According to the definition of formal gaussian integral, in our case

$$\begin{aligned} \int_{\text{FG}} d\mathbf{x} e^{\frac{1}{2} Q(\mathbf{x})} P(\mathbf{x}) &= \int_{\text{FG}} d\mathbf{x} e^{\frac{1}{2} Q_{\text{str}}} \left( e^{\frac{1}{2} (Q - Q_{\text{str}})} P \right) \\ &= \left( \det \hat{Q}_{\text{str}} \right)^{-1/2} \left[ \left( e^{\frac{1}{2} (Q - Q_{\text{str}})} P \right) \mathfrak{b}_{\mathbf{x}, \mathbf{x}; \text{LR}} \epsilon^{-\frac{1}{2} Q_{\text{str}}} \right]. \end{aligned} \quad (4.107)$$

We introduce the functions

$$Q(x; t) = Q(x) + t \left( Q_{\text{str}}(x) - Q(x) \right), \quad P(\mathbf{x}; t) = e^{\frac{t}{2} (Q(\mathbf{x}) - Q_{\text{str}}(\mathbf{x}))} P(\mathbf{x}). \quad (4.108)$$

depending on a  $\mathbb{Q}$ -parameter  $t$  and consider an expression

$$A(t) = \left( \det \hat{Q}(t) \right)^{-1/2} \left( P(\mathbf{x}; t) \text{ } \text{ } \text{ } e^{-\frac{1}{2} Q(\mathbf{x}; t)} \right). \quad (4.109)$$

In view of eq.(4.107),  $A(1)$  equals the *l.h.s.* of eq.(4.84), whereas  $A(0)$  obviously equals the *r.h.s.* of that equation. Let us consider  $\partial_t A(t)$ . It is easy to see that the proof of eq.(4.99) also works in the case when  $Q(\mathbf{x})$  is a quadratic function of  $\mathbf{x}$  which is not necessarily strut. Since in our particular case

$$P(\mathbf{x}; t) \partial_t Q(\mathbf{x}; t) + 2 \partial_t P(\mathbf{x}; t) = 0, \quad (4.110)$$

then, according to eq.(4.99),  $\partial_t A(t) = 0$ . Hence  $A(0) = A(1)$  and this proves eq.(4.84) in case when  $Q(\mathbf{x})$  is a quadratic function of  $\mathbf{x}$ .  $\square$

Now we are ready to prove that a gaussian integral commutes with the injection  $f_{\mathcal{QH}}$ .

**Theorem 4.36** *Let  $G(\mathbf{x}; \mathbf{a})$  be a tensor field or an integration measure which depends on Cartan parameters  $\mathbf{a}$ . Suppose that  $\mathbf{a} \in \mathfrak{L}'_c$  and denote  $G'(\mathbf{x}) = f_{\mathcal{QH}}(G(\mathbf{x}; \mathbf{a}))$ . If  $G'$  is SP, then  $G$  is SP. If in addition  $G$  is non-degenerate, then  $G'$  is also non-degenerate and*

$$\int_{\text{FG}} d\mathbf{x} G'(\mathbf{x}) = f_{\mathcal{QH}} \left( \int_{\text{FG}} d\mathbf{x} G(\mathbf{x}; \mathbf{a}) \right). \quad (4.111)$$

*Proof.*  $G'$  being SP and belonging to the image of  $f_{\mathcal{QH}}$  means that  $G' = e^{\frac{1}{2} Q'(\mathbf{x})} P'(\mathbf{x})$ , where

$$Q' = \sum_{i,j=1}^L l_{ij}(\mathbf{a}) x_i \text{ --- } x_j, \quad l_{ij}(\mathbf{a}) \in \mathbb{Q}[[\mathbf{a}]], \quad (4.112)$$

and  $P' \in \mathcal{D}_{\mathfrak{L}}^C \subset \mathcal{QD}_{\mathfrak{L}}^C$  is  $\mathbf{x}$ -substantial. Then  $G(\mathbf{x}; \mathbf{a}) = e^{\frac{1}{2} Q(\mathbf{x})} P(\mathbf{x}; \mathbf{a})$ , where

$$Q(\mathbf{x}) = \sum_{i,j=1}^L l_{ij}(\mathbf{a}) \Big|_{\mathbf{a}=0} x_i \text{ --- } x_j, \quad (4.113)$$

while

$$P(\mathbf{x}; \mathbf{a}) = \exp f_{\mathcal{QH}}^{-1} \left( \frac{1}{2} \sum_{i,j=1}^L (l_{ij}(\mathbf{a}) - l_{ij}(\mathbf{a}) \Big|_{\mathbf{a}=0}) x_i \text{ --- } x_j \right) f_{\mathcal{QH}}^{-1}(P'(\mathbf{x})) \quad (4.114)$$

Since the exponent in eq.(4.114) does not contain  $\mathcal{D}_{\mathfrak{g}}^{\mathbb{C}}$  struts and since  $f_{\mathcal{QH}}^{-1}$  in contrast to  $f_{\mathcal{QH}}$  does not create new struts, then  $P(\mathbf{x}; \mathbf{a})$  is  $\mathbf{x}$ -substantial and hence  $G(\mathbf{x}; \mathbf{a})$  is SP.

If  $G$  is SPND, then  $\det ||l_{ij}(\mathbf{a})|| \Big|_{\mathbf{a}=0} \neq 0$ , hence  $\det ||l_{ij}(\mathbf{a})|| \not\equiv 0$  and  $G'$  is also SPND.

It remains to prove eq.(4.111). Consider its *r.h.s.* :

$$\int_{\text{FG}} d\mathbf{x} G(\mathbf{x}; \mathbf{a}) = \int_{\text{FG}} d\mathbf{x} \exp \left( \frac{1}{2} f_{\mathcal{QH}}^{-1}(Q'(\mathbf{x})) \right) f_{\mathcal{QH}}^{-1}(P'(\mathbf{x})). \quad (4.115)$$

Since  $f_{\mathcal{QH}}^{-1}(Q'(\mathbf{x}))$  is a quadratic function of  $\mathbf{x}$  and its strut part is assumed to be non-degenerate, then Theorem 4.35 says that we can calculate the integral in the *r.h.s.* of eq.(4.115) with the help of eq.(4.84) in which now  $Q = f_{\mathcal{QH}}^{-1}(Q')$  and  $P = f_{\mathcal{QH}}^{-1}(P')$ . The result obviously coincides with the *l.h.s.* of eq.(4.111) (we can use Theorem 4.19 in order to establish the equality of ‘1-loop determinant factors’).  $\square$

**Remark 4.37** Equation (4.112) indicates that

$$\det \hat{Q} = \det ||l_{ij}(\mathbf{a})|| \Big|_{\mathbf{a}=0}. \quad (4.116)$$

Therefore it may happen that  $\det \hat{Q} = 0$ , whereas  $\det \hat{Q}' \neq 0$ , which means that  $G(\mathbf{x}; \mathbf{a})$  is degenerate, while  $G'(\mathbf{x})$  is non-degenerate. This is similar to what we described in Remark 4.20: a definition of a gaussian integral is finer in  $\mathcal{QD}_{\mathfrak{g}}^{\mathbb{C}}$  than in  $\mathcal{B}_{\mathfrak{g}}^{\mathbb{C}}$ . This is important for us, since it will turn out that the  $U(1)$ -RCC invariant is defined through a gaussian integral which is defined only in  $\mathcal{QD}_{\mathfrak{g}}^{\mathbb{C}}$ .

## 5 A formal integration over a coadjoint orbit

### 5.1 Invariant integration measure on a coadjoint orbit

In subsection 2.3 we used  $\mathfrak{G}/\mathfrak{T}$  as a model of a coadjoint orbit, and exponential map turned the root space  $\mathfrak{r} \subset \mathfrak{g}$  into a coordinate chart in the vicinity of  $e \in \mathfrak{G}$ . Now we are going to do the same at the level of graph algebras. Since we consider just a single coadjoint orbit (a case of multiple orbits would be an obvious generalization), we introduce a single root coordinate color  $x$ . Our immediate task is to define a diffeomorphism  $F_y$  ( $y$  being a root parameter), which is an analog of diffeomorphism  $F_{\vec{y}}$  defined by the commutative diagram (2.45), and then find an  $F_y$ -invariant integration measure  $\mu(x)$ .

In order to distinguish between root, Cartan and total labels and coordinates, we assume the following convention: root coordiantes are denoted as  $x, y, \dots$ , Cartan coordinates are

denoted as  $a, b, \dots$  and total coordinates are denoted as  $X, Y, \dots$ . For any map  $Y = F(\cdot)$  we define two related maps  $y = F_r(\cdot)$  and  $a = F_c(\cdot)$  in which the total  $Y$  leg of  $F$  is replaced by a root  $y$  leg or by a Cartan  $a$  leg. Also, if we replace a total label  $X$  in the argument of any object  $A(X)$  by either  $y$  or  $a$ , then this means that the  $X$  legs of  $A$  are declared root or Cartan.

We have to express eq. (2.46) in purely graphical terms. First, we have to learn how to graphically multiply the exponentials with non-commutative exponents. This has been already described in [3], where a BCH ‘forest’ has been introduced. If  $X$  and  $Y$  are two formal non-commutative variables, then according to BCH,

$$e^X e^Y = e^Z$$

$$Z = X + Y + \frac{1}{2} [X, Y] + \frac{1}{12} [X, [X, Y]] - \frac{1}{12} [Y, [X, Y]] + \text{higher commutators.} \quad (5.1)$$

Following [3], we turn every term in the expression for  $Z$  into a tree graph with total edges, placing the color  $Z$  at the ‘root’ and colors  $X$  and  $Y$  at the ends of the ‘branches’ of these trees. The sum of all these trees defines a BCH map which we denote as  $Z = E(X, Y)$ :

$$E = Z \text{ --- } X + Z \text{ --- } Y + \text{trees with multiple edges.} \quad (5.2)$$

We also define a multiple BCH map  $Y = E(X_1, \dots, X_n)$  as a logarithm of the product of  $n$  exponentials. The multiple map can be presented as a composition of binary BCH maps in any order compatible with the associativity. The BCH maps satisfy some obvious properties coming from the basic properties of exponentials. Among these properties are

$$E(t_1 X, t_2 X) = E((t_1 + t_2)X, 0) = \cdot \text{ --- } (t_1 + t_2)X, \quad (5.3)$$

which follows from the commutativity of legs of the same color,

$$E(a, b) = E(a + b, 0) = \cdot \text{ --- } (a + b), \quad (5.4)$$

which follows from CC relations expressing the commutativity of Cartan edges (Lemma 3.1), and

$$E(Y, X, -Y) = \text{Ad}_{\exp(Y)}(X), \quad (5.5)$$

$$\text{Ad}_{\exp(E(X, Y))} = \text{Ad}_{\exp(X)} \circ \text{Ad}_{\exp(Y)}. \quad (5.6)$$

( $\text{Ad}_{\exp(\cdot)}$  map being defined in Example 4.6), which follows from the IHX relation and the combinatorial identity  $e^x e^y e^{-x} = e^{e^{\text{ad}_x} y}$ . Also, due to the presence of struts in (5.2), the map  $E$  can not mix root and Cartan input in order to produce a purely root output:



**Lemma 5.1** *Consider two maps:  $y = H_{(r)}(X)$  with a root output and  $a = H_{(c)}(X)$  with a Cartan output. If a Cartan part  $F_c$  of a map  $Y = F(X)$  defined as*

$$F(X) = E(H_{(r)}(X), H_{(c)}(X)), \quad (5.7)$$

*is zero, then*

$$F(X) = H_{(r)}(X) \quad \text{and} \quad H_{(c)}(X) = 0. \quad (5.8)$$

*Proof.* Consider a graph of  $H_{(c)}$  with the smallest number of edges. Then due to the struts in (5.2) it will produce the graph with the smallest number of edges among the graphs in *r.h.s.* of eq. (5.7) with Cartan output. Hence, this graph can not be cancelled to conform with the fact that the *l.h.s.* of eq. (5.7) has only root output.  $\square$

Next, we have to learn how to split a single exponential into a product of two exponentials: one of a root element and the other of a Cartan element. Consider a label space with two coordinates  $(x, a)$ . The map  $(y, b) = S(x, a)$  that we want is the inverse of the map  $(x, a) = \tilde{E}(y, b)$  defined by the formula

$$\tilde{E}(y, b) = (E_r(y, b), E_c(y, b)). \quad (5.9)$$

The formula for the  $E$  map (5.2) demonstrates that the strut part of  $\tilde{E}$  is the identity map  $\text{Id}$ , so according to Theorem 4.4,  $\tilde{E}$  is invertible and its inverse can be constructed perturbatively as described in its proof. Now we define the map  $z = F_y(x)$  as

$$F_y(x) = S_r(E_r(y, x), E_c(y, x)), \quad (5.10)$$

where  $y = S_r(x, a)$  is a part of the map  $(y, b) = S(x, a)$  which has the color  $y$  at its outgoing leg. The strut part of  $z = F_y(x)$  is

$$(F_y)_{\text{str}} = z - - - x + z - - - y. \quad (5.11)$$

We will also need a similar map

$$\tilde{F}_y(x) = S_c(E_r(y, x), E_c(y, x)), \quad (5.12)$$

so that according to the definition of  $S$ ,

$$E(y, x) = E(F_y(x), \tilde{F}_y(x)). \quad (5.13)$$

**Lemma 5.2** *The diffeomorphism  $z = F_y(x)$  is narrow.*

*Proof.* The BCH formula shows that  $Z = E(X, Y)$  is narrow, therefore  $(x, a) = \tilde{E}(y, b)$  is narrow and so is its inverse  $(y, b) = S(x, a)$ . Hence the composition (5.10) is also narrow.  $\square$

Let us establish a few properties of  $F_y$ . First of all, it is easy to verify that

$$F_{t_2 y}(t_1 y) = (t_1 + t_2)y, \quad \tilde{F}_{t_2 y}(t_1 y) = 0. \quad (5.14)$$

Next, we establish the graph analog of eq. (2.49), but in order to formulate it, we need a slightly modified version of the map  $Z = \text{Ad}_{\exp(Y)}(X)$ . Namely, when the total parameter  $Y$  is replaced by a Cartan parameter  $a$ , and the argument is a root coordinate  $x$ , then the  $\text{CC}_2$  relation says that  $\left(\text{Ad}_{\exp(a)}(x)\right)_{\mathfrak{c}} = 0$  and  $Z = \text{Ad}_{\exp(a)}(x)$  is reduced to  $z = \text{Ad}_{\exp(a)}x$ . It is easy to see that the root-root maps  $\text{Ad}_{\exp(ta)}$  still form a 1-parametric group, which is generated by a (purely root) vector field  $\vec{\text{ad}}_a$ .

**Lemma 5.3** *The diffeomorphisms  $F_y$  satisfy the following composition formula*

$$F_{y_2} \circ F_{y_1} = F_{F_{y_2}(y_1)} \circ \text{Ad}_{\exp(\tilde{F}_{y_2}(y_1))}. \quad (5.15)$$

*Proof.* consider a map from  $(x, y_1, y_2)$  to  $Z$

$$Z = E\left(y_2, y_1, x, -\tilde{F}_{y_1}(x) - \tilde{F}_{\tilde{F}_{y_2}} \circ F_{y_1}(x)\right). \quad (5.16)$$

We will transform the *r.h.s.* by applying eq. (5.13) to pairs of adjacent root coordinates in two different orders. First, we begin by applying eq. (5.13) to  $(y_1, x)$

$$\begin{aligned} Z &= E\left(y_2, F_{y_1}(x), \tilde{F}_{y_1}(x), -\tilde{F}_{y_1}(x) - \tilde{F}_{y_2} \circ \tilde{F}_{y_1}(x)\right) \\ &= E\left(y_2, F_{y_1}(x), -\tilde{F}_{y_2} \circ F_{y_1}(x)\right) \\ &= E\left(F_{y_2} \circ F_{y_1}(x), \tilde{F}_{y_2} \circ F_{y_1}(x), -\tilde{F}_{y_2} \circ F_{y_1}(x)\right) \\ &= F_{y_2} \circ F_{y_1}(x). \end{aligned} \quad (5.17)$$

On the other hand, we may first apply eq. (5.13) to the pair  $(y_2, y_1)$

$$\begin{aligned} Z &= E\left(F_{y_2}(y_1), \tilde{F}_{y_2}(y_1), x, -\tilde{F}_{y_1}(x) - \tilde{F}_{y_2} \circ F_{y_1}(x)\right) \\ &= E\left(F_{y_2}(y_1), \tilde{F}_{y_2}(y_1), x, -\tilde{F}_{y_2}(y_1), \tilde{F}_{y_2}(y_1) - \tilde{F}_{y_1}(x) - \tilde{F}_{y_2} \circ F_{y_1}(x)\right) \\ &= E\left(F_{y_2}(y_1), \text{Ad}_{\exp(\tilde{F}_{y_2}(y_1))}(x), \tilde{F}_{y_2}(y_1) - \tilde{F}_{y_1}(x) - \tilde{F}_{y_2} \circ F_{y_1}(x)\right) \\ &= E\left(F_{F_{y_2}(y_1)} \circ \text{Ad}_{\exp(\tilde{F}_{y_2}(y_1))}(x), \tilde{F}_{F_{y_2}(y_1)} \circ \text{Ad}_{\exp(\tilde{F}_{y_2}(y_1))}(x) + \tilde{F}_{y_2}(y_1) \right. \\ &\quad \left. - \tilde{F}_{y_1}(x) - \tilde{F}_{y_2} \circ F_{y_1}(x)\right). \end{aligned} \quad (5.18)$$

Comparing eqs.(5.17) and (5.18) we find that

$$\begin{aligned} F_{y_2} \circ F_{y_1}(x) = E \Big( & F_{F_{y_2}(y_1)} \circ \text{Ad}_{\exp(\tilde{F}_{y_2}(y_1))}(x), \tilde{F}_{F_{y_2}(y_1)} \circ \text{Ad}_{\exp(\tilde{F}_{y_2}(y_1))}(x) \\ & + \tilde{F}_{y_2}(y_1) - \tilde{F}_{y_1}(x) - \tilde{F}_{y_2} \circ F_{y_1}(x) \Big). \end{aligned} \quad (5.19)$$

According to the first of eqs.(5.8) of Lemma 5.1, eq.(5.19) implies the composition law (5.15).

□

**Corollary 5.4** *The diffeomorphisms  $F_{ty}$  form a 1-parametric group.*

*Proof.* . This corollary follows from a simple calculation based on eqs.(2.71) and (5.14)

$$F_{t_2y} \circ F_{t_1y} = F_{F_{t_2y}(t_1y)} \circ \text{Ad}_{\exp(\tilde{F}_{t_2y}(t_1y))} = F_{(t_1+t_2)y} \circ \text{Ad}_{\exp(0)} = F_{(t_1+t_2)y}. \quad (5.20)$$

□

Let  $\xi_{F,y}$  denote the vector field which generates  $F_{ty}$ . It follows from eq.(5.11) that its strut part is

$$\left( \xi_{F,y}(x) \right)_{\text{str}} = \partial_x - - - y. \quad (5.21)$$

Following eq.(2.47) we choose the measure for a root space to be the inverse determinant of the derivative of  $F_x$

$$\mu(x) = \left( \det F'_x(0) \right)^{-1}. \quad (5.22)$$

Obviously,  $\mu(0) = 1$  and it follows from eq.(4.74) that

$$\mu(x) = F_{-x}^* \mu(0) \quad (5.23)$$

(note that eq.(4.74) defines the action of a diffeomorphism as a pull-back).

**Theorem 5.5** *The measure (5.22) is invariant under the substitution  $y = \text{Ad}_{\exp(a)}(x)$ .*

*Proof.* Since the substitutions  $\text{Ad}_{\exp(ta)}$  form a 1-parametric group generated by  $\vec{\text{ad}}_a$ , then it is sufficient to check that

$$L_{\vec{\text{ad}}_a} \mu = \nabla_{\vec{\text{ad}}_a} \mu + \mu \text{div}_x(\vec{\text{ad}}_a) = 0. \quad (5.24)$$

In fact, both terms in the middle expression are equal to 0. Indeed  $\nabla_{\vec{\text{ad}}_a} \mu = 0$  because of eq.(4.29) of Lemma 4.7, while  $\text{div}_x(\vec{\text{ad}}_a) = 0$ , because the symmetry of the graph  $\left( \begin{smallmatrix} \_ \\ \_ \end{smallmatrix} \right)$  sends an odd power of  $a$  to zero. □

**Theorem 5.6** *The measure (5.22) is invariant under the action of  $F_y$ .*

*Proof.* This invariance is a simple corollary of Lemma 5.3 and Theorem 5.5. Indeed

$$\begin{aligned}
F_y^* \mu(x) &= F_y^* \left( F_{-x}^* \mu(0) \right) = \left( F_{-x} \circ F_y \right)^* \mu(0) = \left( (F_{-y} \circ F_x)^{-1} \right)^* \mu(0) \\
&= \left[ \left( F_{F_{-y}(x)} \circ \text{Ad}_{\exp(\tilde{F}_{-y}(x))} \right)^{-1} \right]^* \mu(0) \\
&= \left( \text{Ad}_{\exp(-\tilde{F}_{-y}(x))} \circ F_{-F_{-y}(x)} \right)^* \mu(0) \\
&= F_{-F_{-y}(x)}^* \left( \text{Ad}_{\exp(-\tilde{F}_{-y}(x))}^* \mu(0) \right) \\
&= F_{-F_{-y}(x)}^* \mu(0) = \mu(F_{-y}(x)) = \mu(F_y^{-1}(x)).
\end{aligned} \tag{5.25}$$

The resulting equation  $F_y^* \mu(x) = \mu(F_y^{-1}(x))$  means that the measure  $\mu(x)$  is indeed invariant under the action of  $F_y$ .  $\square$

Let us describe the structure of  $\mu(x)$  in more details.

**Theorem 5.7** *The measure  $\mu(x)$  is an exponential  $\mu(x) = e^{\omega(x)}$ , where  $\omega(x)$  is a sum of wheel graphs, and  $\mu(x)$  is an even function of  $x$ :  $\mu(-x) = \mu(x)$ .*

*Proof.* For the purpose of calculating  $\mu(x)$  it is better to use the inverse function derivative formula (4.37) in order to rewrite eq.(5.22) as

$$\mu(x) = \det F'_{-x}(x). \tag{5.26}$$

The reason is that  $F'_{-x}$  is much easier to calculate. Indeed,  $F$  is defined by eq.(5.10) as a composition of two maps, while  $E(-x, x) = 0$  and the derivative  $\partial_z S_r(z, a)$  at  $z = 0, a = 0$  is the identity matrix. Therefore,

$$F'_{-x}(x) = \left( \partial_x E_r(y, x) \right) \Big|_{y=-x}. \tag{5.27}$$

The formula for the derivative  $\left( \partial_X E(Y, X) \right) \Big|_{Y=-X}$  of  $E(Y, X)$ , all of whose edges and indices are total, follows from the combinatorial relation which is similar to that of eq.(2.52):

$$e^{-X} e^{X+\Delta X} - 1 \approx \frac{1 - e^{-\text{ad}_X}}{\text{ad}_X} \Delta X = \left( 1 + \sum_{n=1}^{\infty} \frac{(-\text{ad}_X)^n}{(n+1)!} \right) \Delta X. \tag{5.28}$$

Therefore,

$$\left( \partial_X E(Y, X) \right) \Big|_{Y=-X} = I + M_X, \quad M_X = \sum_{n=1}^{\infty} \frac{(-\widehat{\text{ad}_X})^n}{(n+1)!}. \tag{5.29}$$

where  $\widehat{\text{ad}}_X$  is a matrix field with total indices. In order to express the derivative  $\left(\partial_x E(y, x)\right)\Big|_{y=-x}$ , we have to replace the total legs  $dX$  and  $\partial_X$  of  $M_X$  (and  $I$ ) with root legs  $dx$  and  $\partial_x$ , and replace the total  $X$  coordinate label with root label  $x$ . We denote the corresponding operator as  $M_{r,x}$ . Then we find from the definition (4.51) that

$$\begin{aligned}\mu(x) &= \det F'_{-x}(x) = \det \left(\partial_x E_r(y, x)\right)\Big|_{y=-x} = \det(I + M_{r,x}) = \exp \left( \text{Tr} \log_x (I + M_{r,x}) \right) \\ &= \exp \left( - \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{Tr}(M_{r,x}^{\times n}) \right).\end{aligned}\tag{5.30}$$

Since each trace  $\text{Tr}(M_{r,x}^{\times n})$  is obviously represented by a wheel graph, then  $\omega(x) = \log \mu(x)$  is indeed a sum of wheels. Note that these wheels are different from those related to the wheeling map and Duflo isomorphism, because  $n$  internal edges of  $\text{Tr}(M_{r,x}^{\times n})$  are root rather than total due to the fact that we passed from  $M_X$  to  $M_{r,x}$ .

In order to prove that  $\mu(x)$  is even, we observe that  $(\widehat{\text{ad}}_X)^T = -\widehat{\text{ad}}_X = \widehat{\text{ad}}_{-X}$ , hence  $M_X^T = M_{-X}$  and  $M_{r,x}^T = -M_{r,-x}$ . Then, in view of eqs. (4.40),

$$\text{Tr}(M_{r,x}^{\times n}) = \text{Tr}(M_{r,x}^{\times n})^T = \text{Tr}(M_{r,-x}^{\times n})\tag{5.31}$$

and hence  $\mu(x) = \mu(-x)$ .  $\square$

Comparing eqs. (5.30) and (2.57) we see that

$$\mu_{\vec{a}}(\vec{x}) = (-2\pi)^{-|\Delta_+|} d_{\mathfrak{g}}(\vec{a}) \mathcal{T}_{\mathfrak{g}} \mu(\vec{x}).\tag{5.32}$$

For  $L$  root coordinates  $\mathbf{x}$  we denote

$$\boldsymbol{\mu}(\mathbf{x}) = \prod_{j=1}^L \mu(x_j).\tag{5.33}$$

Theorem 5.7 has an obvious

**Corollary 5.8** *The integration measure  $\boldsymbol{\mu}(\mathbf{x})$  is  $\mathbf{x}$ -substantial.*

## 5.2 Properties of the coadjoint orbit integral

So far, we have described a graph analog of a coadjoint orbit defined as  $\mathfrak{G}/\mathfrak{I}$ . That quotient space is related to an actual coadjoint orbit  $\mathcal{O}_{\vec{a}}$  by the isomorphism (2.41). Its graph analog is a map  $X = R_a(x)$  (where  $x$  is a root argument and  $a$  is a Cartan parameter) defined as

$$R_a(x) = \text{Ad}_{\exp(x)}(a).\tag{5.34}$$

**Lemma 5.9** *The map  $R_a$  is narrow and it satisfies the following properties*

$$R_a \circ \text{Ad}_{\exp(b)} = \text{Ad}_{\exp(b)} \circ R_a, \quad (5.35)$$

$$R_a \circ F_y = \text{Ad}_{\exp(y)} \circ R_a. \quad (5.36)$$

*Proof.* The narrowness of  $Z = R_a(x)$  is obvious. To prove eq. (5.35), we calculate its *l.h.s.* starting with the definition (5.34) and then applying eq. (4.28) and the relation

$$\text{Ad}_{\exp(b)}(a) = a, \quad (5.37)$$

which follows from CC relations. Thus

$$\begin{aligned} R_a \circ \text{Ad}_{\exp(b)}(x) &= \text{Ad}_{\exp(\text{Ad}_{\exp(b)}(x))}(a) = \text{Ad}_{\exp(b)} \circ \text{Ad}_{\exp(x)} \circ \text{Ad}_{\exp(-b)}(a) \\ &= \text{Ad}_{\exp(b)} \circ \text{Ad}_{\exp(x)}(a) = \text{Ad}_{\exp(b)} \circ R_a. \end{aligned} \quad (5.38)$$

In order to prove eq. (5.36), we start with its *r.h.s.*

$$\begin{aligned} \text{Ad}_{\exp(y)} \circ R_a(x) &= \text{Ad}_{\exp(y)} \circ \text{Ad}_{\exp(x)}(a) = \text{Ad}_{\exp(E(y,x))}(a) \\ &= \text{Ad}_{\exp(E(F_y(x), \tilde{F}_y(x)))}(a) = \text{Ad}_{\exp(F_y(x))} \circ \text{Ad}_{\exp(\tilde{F}_y(x))}(a) \\ &= \text{Ad}_{\exp(F_y(x))}(a) = R_a \circ F_y. \end{aligned} \quad (5.39)$$

□

The substitution (5.34) converts a function  $G(\mathbf{X})$  of total coordinates  $\mathbf{X}$  into a function  $G^R(\mathbf{x}; \mathbf{a})$  of root coordinates  $\mathbf{x}$  and Cartan coordinates  $\mathbf{a}$  defined as

$$G^R(\mathbf{x}; \mathbf{a}) = G(R_{a_1}(x_1), \dots, R_{a_L}(x_L)). \quad (5.40)$$

Since we will be interested in the integrals of functions  $G^R$ , then following Remark 4.37, we go further and apply the map  $f_{\mathcal{QH}}$  to  $G^R$  assuming that labels  $\mathbf{a}$  belong to  $\mathcal{L}'_c$ :

$$G^{\mathcal{QR}}(\mathbf{x}) = f_{\mathcal{QH}}(G^R(\mathbf{x}; \mathbf{a})). \quad (5.41)$$

In other words,  $\mathbf{a}$  are no longer parameters in  $G^{\mathcal{QR}}(\mathbf{x})$ , rather their legs are hidden in symmetric algebras of cohomologies of graphs.

**Theorem 5.10** *Let  $G(Z)$  be a function such that  $G^{\mathcal{QR}}(x) = f_{\mathcal{QH}}(G(R_a(x)))$  is SPND as a function of  $x$ . Then*

$$\int_{\text{FG}} dx G(\text{Ad}_{\exp(b)} \circ R_a(x)) \mu(x) = \int_{\text{FG}} dx G(x) \mu(x), \quad (5.42)$$

$$\int_{\text{FG}} dx G(\text{Ad}_{\exp(y)} \circ R_a(x)) \mu(x) = \int_{\text{FG}} dx G(x) \mu(x). \quad (5.43)$$

*Proof.* We will prove the first formula, since both proofs are essentially the same. According to eq.(5.35),

$$\int_{\text{FG}} dx G\left(\text{Ad}_{\exp(b)} \circ R_a(x)\right) \mu(x) = \int_{\text{FG}} dx G\left(R_a \circ \text{Ad}_{\exp(b)}(x)\right) \mu(x). \quad (5.44)$$

Now eq.(5.42) follows from Corollary 4.34, where  $G(y)$  is  $G(R_a(y))$  (this corollary is applicable, since  $y = \text{Ad}_{\exp(b)}(x)$  is narrow), and from Theorem 5.5 which states the  $\text{Ad}_{\exp(b)}$ -invariance of the measure  $\mu(x)$ .  $\square$

In what follows we will be interested in functions  $G(\mathbf{X})$  such that  $G^{\mathcal{Q}\mathcal{R}}(\mathbf{x}; \mathbf{a})$  considered as functions of  $\mathbf{x}$  are SPND, so that we can define an integral  $I(\mathbf{a}) = \int_{\text{FG}} d\mathbf{x} G^{\mathcal{Q}\mathcal{R}}(\mathbf{x}; \mathbf{a}) \mu(\mathbf{x})$ .

**Example 5.11** For  $L = 1$  let  $G(X; b) = \exp(X \cdots b)$ , where  $b$  is a Cartan parameter. We assume that  $a, b \in \mathfrak{L}'_c$ , so  $G^{\mathcal{Q}\mathcal{R}}(x; a, b) = \exp(ab \circ + \frac{1}{2}ab x - - x + \mathcal{O}(x^3))$ . Obviously,  $G^{\mathcal{Q}\mathcal{R}}(x; a, b)$  is SPND in  $x$  and  $Q(x) = -ab x - - x$ .

**Conjecture 5.12 (Duistermaat-Heckmann for graphs)** *The stationary phase integral  $\int_{\text{FG}} dx G^{\mathcal{Q}\mathcal{R}}(x; a, b) \mu(x)$  is ‘1-loop exact’, that is*

$$\int_{\text{FG}} dx \exp\left(R_a(x) \cdots b\right) \mu(x) = \exp(ab \circ) \exp\left(-\frac{1}{2} \log(ab) \left(\begin{smallmatrix} \square \\ \square \end{smallmatrix}\right)\right). \quad (5.45)$$

In other words, this conjecture says that for this particular integral, the term

$$P(x) \vdash_{x,x;\text{LR}} e^{-\frac{1}{2}Q^{-1}(x)}$$

of eq.(4.84) is equal to  $P(0)$ . We hope to prove this conjecture in [15] by modifying the standard Duistermaat-Heckman proof for graph algebra case with the help of grassmanian variables.

We are particularly interested in integrals which come from  $\text{Ad}_{\exp(y)}$ -invariant functions  $G(\mathbf{X})$ :

$$G(\text{Ad}_{\exp(y)}(X_1), \dots, \text{Ad}_{\exp(y)}(X_L)) = G(X_1, \dots, X_L). \quad (5.46)$$

Suppose that  $G(\mathbf{X})$  is  $\text{Ad}_{\exp(y)}$ -invariant and that  $G^{\mathcal{Q}\mathcal{R}}(\mathbf{x})$  is SP. Denote the (twice) corresponding strut exponent of  $G^{\mathcal{Q}\mathcal{R}}(\mathbf{x})$  as

$$Q(\mathbf{x}) = \sum_{i,j=1}^L l_{ij}(\mathbf{a}) x_i - - x_j. \quad (5.47)$$

We denote

$$Q_k(\mathbf{x}_{(k)}) = Q(\mathbf{x})|_{x_k=0}. \quad (5.48)$$

The matrix of the ‘quadratic form’  $Q_k$  is  $||l_{ij}(\mathbf{a})||_k$  which is the  $(k, k)$ -minor of the full matrix  $||l_{ij}(\mathbf{a})||$ .

**Lemma 5.13** *If  $G(\mathbf{X})$  is  $\text{Ad}_{\exp(y)}$ -invariant, then*

$$\det ||l_{ij}(\mathbf{a})|| = 0 \quad (5.49)$$

*and the determinant  $\det ||l_{ij}(\mathbf{a})||_k$  does not depend on  $k$ .*

*Proof.* The claim of this lemma follows easily from the following relation

$$\sum_{j=1}^L l_{ij}(\mathbf{a}) = 0, \quad 1 \leq i \leq L. \quad (5.50)$$

In order to prove it, we observe that since  $G(\mathbf{X})$  is  $\text{Ad}_{\exp(y)}$ -invariant, then, in view of eq.(5.36),  $G^{\mathcal{Q}\text{R}}(\mathbf{x})$  is  $F_y$ -invariant, which means that

$$L_{\xi_{F,y}} G^{\mathcal{Q}\text{R}}(\mathbf{x}) = 0, \quad (5.51)$$

where  $\xi_{F,y}(\mathbf{x}) = \sum_{j=1}^L \xi_{F,y}(x_j)$ . Therefore

$$L_{(\xi_{F,y})_{\text{str}}} Q(\mathbf{x}) = 0, \quad (5.52)$$

where  $(\xi_{F,y})_{\text{str}}$  is determined by eq.(5.21). Equation (5.52) implies eq.(5.50).  $\square$

Since  $\det ||l_{ij}(\mathbf{a})||_k$  does not depend on  $k$ , then neither does  $\det \hat{Q}_k$ ,  $\hat{Q}$  being defined by eq.(4.83). Thus we denote these determinants as

$$\det' ||l_{ij}(\mathbf{a})|| = \det ||l_{ij}(\mathbf{a})||_k, \quad \det' \hat{Q} = \det \hat{Q}_k. \quad (5.53)$$

The following theorem will imply the independence of the universal  $U(1)$ -RCC invariant of the choice of the broken link component.

**Theorem 5.14** *Let  $G(\mathbf{X})$  be an  $\text{Ad}_{\exp(y)}$ -invariant function of  $L$  total coordinates  $\mathbf{X}$*

$$G(\text{Ad}_{\exp(y)}(X_1), \dots, \text{Ad}_{\exp(y)}(X_L)) = G(X_1, \dots, X_L). \quad (5.54)$$

*Suppose that for some  $k$  ( $1 \leq k \leq L$ ) the function  $G^{\mathcal{Q}\text{R}}(\mathbf{x})|_{x_k=0}$  is non-degenerate SP. Then the same is true for any other  $j$ ,  $1 \leq j \leq L$  and*

$$\int_{\text{FG}} d\mathbf{x}_{(j)} \left( G^{\mathcal{Q}\text{R}}(\mathbf{x}) \boldsymbol{\mu}(\mathbf{x}) \right) \Big|_{x_j=0} = \int_{\text{FG}} d\mathbf{x}_{(k)} \left( G^{\mathcal{Q}\text{R}}(\mathbf{x}) \boldsymbol{\mu}(\mathbf{x}) \right) \Big|_{x_k=0}. \quad (5.55)$$



*Proof.* In order to simplify our notations, let us assume that  $j = 1$  and  $k = 2$ . Then assuming that the individual integral over  $x_1$  is non-degenerate, we can rewrite the *r.h.s.* of eq.(5.43) as

$$\begin{aligned}
& \int_{\text{FG}} dx_3 \cdots dx_L \mu(x_3) \cdots \mu(x_L) \left( \int_{\text{FG}} dx_1 \mu(x_1) G\left(R_{a_1}(x_1), a_2, R_{a_3}(x_3), \dots, R_{a_L}(x_L)\right) \right) \\
&= \int_{\text{FG}} dx_3 \cdots dx_L \mu(x_3) \cdots \mu(x_L) \left( \int_{\text{FG}} dx_1 \mu(x_1) \right. \\
&\quad \times G\left(\text{Ad}_{\exp(-x_1)} \circ R_{a_1}(x_1), \text{Ad}_{\exp(-x_1)}(a_2), \text{Ad}_{\exp(-x_1)} \circ R_{a_3}(x_3), \dots, \text{Ad}_{\exp(-x_1)} \circ R_{a_L}(x_L)\right) \Big) \\
&= \int_{\text{FG}} dx_3 \cdots dx_L \mu(x_3) \cdots \mu(x_L) \left( \int_{\text{FG}} dx_2 \mu(x_2) G\left(a_1, R_{a_2}(-x_2), R_{a_3}(x_3), \dots, R_{a_L}(x_L)\right) \right) \\
&= \int_{\text{FG}} dx_3 \cdots dx_L \mu(x_3) \cdots \mu(x_L) \left( \int_{\text{FG}} dx_2 \mu(x_2) G\left(a_1, R_{a_2}(x_2), R_{a_3}(x_3), \dots, R_{a_L}(x_L)\right) \right).
\end{aligned} \tag{5.56}$$

Here the first equality is due to the  $\text{Ad}_{\exp(y)}$ -invariance of  $G(\mathbf{X})$ , the second equality is due to eq.(5.43) of Theorem 5.10 and the definition of  $R_a$  and the third equality is due to parity-invariance of the gaussian integral and integration measure  $\mu(x)$ .  $\square$

**Remark 5.15** Our assumption that the integral over  $x_1$  is non-degenerate is not restrictive. If the integral is degenerate, then we can add an  $\text{Ad}$ -invariant ‘regularizing’ term (*e.g.*  $\epsilon X_1 \longrightarrow X_2$ ) to the strut exponent so that the integral becomes non-degenerate. After proving eq.(5.55) in the regularized form we remove the regularization by setting  $\epsilon = 0$ .

**Lemma 5.16** *Let  $T(X)$  be a tensor field such that  $T(R_a(x))$  is SPND. Consider another tensor field*

$$\check{T}(X; Y) = \nabla_{\vec{\text{ad}}_Y(X)} T(X), \tag{5.57}$$

*which depends on  $X$  and on a total parameter  $Y$ . Then*

$$\int_{\text{FG}} dx \check{T}(R_a(x); Y) \mu(x) = 0. \tag{5.58}$$

*Proof.* Since a total leg is a sum of a root leg and a Cartan leg, then it is sufficient to prove eq.(5.58) for two cases: when  $Y$  is replaced either by a Cartan parameter  $b$  or by a root parameter  $y$ . Since both proofs are essentially the same, we will prove the  $b$  case. It is easy to see that

$$\check{T}(X; b) = \partial_t T(\text{Ad}_{\exp(tb)}(X)) \Big|_{t=0}. \tag{5.59}$$

Then according to eq.(5.42),

$$\begin{aligned} \int_{\text{FG}} dx \tilde{T}(\mathbf{R}_a(x); b) \mu(x) &= \partial_t \int_{\text{FG}} dx T\left(\text{Ad}_{\exp(tb)} \circ \mathbf{R}_a(x)\right) \mu(x) \Big|_{t=0} \\ &= \partial_t \int_{\text{FG}} dx T(\mathbf{R}_a(x)) \mu(x) \Big|_{t=0} = 0. \end{aligned} \quad (5.60)$$

□

For a total valued vector field  $\Xi(X)$  denote

$$\text{dvd}_X \Xi(X) = \flat_{\partial_X, Y; 1} \left( \nabla_{\vec{\text{ad}}_Y(X)} \Xi(X) \right) \quad (5.61)$$

(the name dvd is a combination of div and ad).




Figure 8: The definition of  $\text{dvd}_X \Xi(X)$

**Remark 5.17** It is easy to see that the definition (5.61) can be modified:

$$\text{dvd}_X \Xi(X) = \text{div}_X \left( \widehat{\text{ad}}_X \times \Xi(X) \right). \quad (5.62)$$

(note that  $\text{div}_X \vec{\text{ad}}_Y(X) = 0$ , since the resulting graph is zero because of the AS relation).

The next theorem is essential for the proof of the topological invariance of the universal  $U(1)$ -RCC invariant.

**Theorem 5.18** *Let  $\Xi(X)$  be a total valued vector field such that  $\Xi(\mathbf{R}_a(x))$  is SPND. Then*

$$\int_{\text{FG}} dx \text{dvd}_X \Xi(\mathbf{R}_a(x)) \mu(x) = 0. \quad (5.63)$$

*Proof.* We modify the l.h.s. of eq.(5.63) by using the definition (5.61), commuting the gluing  $\flat_{\partial_X, Y; 1}$  with the formal gaussian integral and then applying (5.58) with  $A = \Xi$

$$\begin{aligned} \int_{\text{FG}} dx \text{dvd}_X \Xi(\mathbf{R}_a(x)) \mu(x) &= \int_{\text{FG}} dx \flat_{\partial_X, Y; 1} \left( \nabla_{\vec{\text{ad}}_Y(X)} \Xi(X) \right) \mu(x) \\ &= \flat_{\partial_X, Y; 1} \left( \int_{\text{FG}} dx \nabla_{\vec{\text{ad}}_Y(X)} \Xi(X) \mu(x) \right) \\ &= 0. \end{aligned} \quad (5.64)$$

□

## 6 A universal $U(1)$ -RCC invariant

### 6.1 Definition and invariance

Let  $\mathcal{L} \in M$  be an  $L$ -component oriented marked Morse link in a rational homology sphere  $M$ . We denote by  $Z_{\mathcal{B}}(\mathcal{L})$  its Kontsevich integral in  $\mathcal{B}_{\mathcal{L}}$ . By replacing the  $\mathcal{L}$  labels  $1, \dots, L$  at the legs of  $Z_{\mathcal{B}}(\mathcal{L})$  with the corresponding variables  $\mathbf{X} = (X_1, \dots, X_L)$ , we turn Kontsevich integral into a function  $Z_{\mathcal{B}}(\mathcal{L}; \mathbf{X})$ . In subsection 3.2 we described an injection  $f_{\text{trv}}$  of the algebra  $\mathcal{B}_{\mathcal{L}}$  into  $\mathcal{B}_{\mathcal{L}}^{\text{C}}$ , which transforms a graph of  $\mathcal{B}_{\mathcal{L}}$  into the same graph in  $\mathcal{B}_{\mathcal{L}}^{\text{C}}$ , all of whose edges are total. Let us denote as  $Z_{\mathcal{B}^{\text{C}}}(\mathcal{L}; \mathbf{X})$  the Kontsevich integral of  $\mathcal{L} \in M$  in  $\mathcal{B}_{\mathcal{L}}^{\text{C}}$ :

$$Z_{\mathcal{B}^{\text{C}}}(\mathcal{L}; \mathbf{X}) = f_{\text{trv}}(Z_{\mathcal{B}}(\mathcal{L}; \mathbf{X})). \quad (6.1)$$

Obviously,  $Z_{\mathcal{B}^{\text{C}}}(\mathcal{L}; \mathbf{X})$  is also a function of  $L$  total variables  $\mathbf{X}$ .

**Lemma 6.1** *Kontsevich integral  $Z_{\mathcal{B}^{\text{C}}}(\mathcal{L}; \mathbf{X})$  is both  $\text{Ad}_{\exp(y)}$ - and  $\text{Ad}_{\exp(b)}$ -invariant as a function of coordinates  $\mathbf{X}$ .*

*Proof.* Let us prove the invariance under  $\text{Ad}_{\exp(y)}$ , the proof for  $\text{Ad}_{\exp(b)}$  is similar. Since  $\text{Ad}_{\exp(ty)}$  forms a 1-parametric group, then it is enough to establish that

$$L_{\vec{\text{ad}}_y(\mathbf{X})} Z_{\mathcal{B}^{\text{C}}}(\mathcal{L}; \mathbf{X}) = \nabla_{\vec{\text{ad}}_y(\mathbf{X})} Z_{\mathcal{B}^{\text{C}}}(\mathcal{L}; \mathbf{X}) = 0, \quad (6.2)$$

where  $\vec{\text{ad}}_y(\mathbf{X}) = \sum_{j=1}^L \vec{\text{ad}}_y(X_j)$ , so that  $\nabla_{\vec{\text{ad}}_y(\mathbf{X})}$  acts on all coordinates  $\mathbf{X}$ . Equation (6.2) follows easily from Lemma 4.7 and from the definition (6.1), which establishes that all edges of  $Z_{\mathcal{B}^{\text{C}}}(\mathcal{L}; \mathbf{X})$  are total.  $\square$

Our next step is to perform a substitution (5.40) and then convert the legs carrying Cartan coordinates  $\mathbf{a}$  into the elements of graph cohomology algebras through the application of the map  $f_{\mathcal{QH}}$  as in eq. (5.41). We define  $\mathcal{D}_{\mathcal{L}}^{\text{C}}$  as an algebra  $\mathcal{D}_{\mathfrak{L}', \mathfrak{L}_c}^{\text{C}}$ , where  $\mathfrak{L}'_c = \{\mathbf{a}\}$  is a set of  $L$  Cartan labels and  $X \setminus \mathfrak{L}'_c = \{\mathbf{x}\}$  is a set of  $L$  root labels. We also define  $\mathcal{QD}_{\mathcal{L}}^{\text{C}}$  as the corresponding algebra  $\mathcal{QD}_{\mathfrak{L}', \mathfrak{L}_c}^{\text{C}}$ . Now we convert Kontsevich integral into an element of  $\mathcal{D}_{\mathcal{L}}^{\text{C}}$  by the formula

$$Z_{\mathcal{D}^{\text{C}}}^{\text{QR}}(\mathcal{L}; \mathbf{x}) = f_{\mathcal{QH}}\left(Z_{\mathcal{B}^{\text{C}}}(\mathcal{L}; R_{a_1}(x_1), \dots, R_{a_L}(x_L))\right). \quad (6.3)$$

**Lemma 6.2** *The function  $Z_{\mathcal{D}^{\text{C}}}^{\text{QR}}(\mathcal{L}; \mathbf{x})$  is SP.*

*Proof.* In view of eq.(2.6), this function is an exponential

$$Z_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x}) = \exp \left( W_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x}) \right), \quad (6.4)$$

where

$$W_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x}) = f_{\mathcal{QH}}(W_{\mathcal{B}^C}(\mathcal{L}; R_{a_1}(x_1), \dots, R_{a_L}(x_L))) \quad (6.5)$$

and  $W_{\mathcal{B}^C}(\mathcal{L}; \mathbf{X}) = f_{\text{trv}}(W(\mathcal{L}; \mathbf{X}))$ . Since  $W(\mathcal{L})$  and  $X = R_a(x)$  are both narrow, then so is  $W_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x})$ . In particular, this means that the dependence of  $Z_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x})$  on struts is purely exponential.  $\square$

Following eq.(4.79), let  $Q(\mathcal{L}; \mathbf{x})$  denote (twice) the strut part of  $W_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x})$

$$Q(\mathcal{L}; \mathbf{x}) = 2(W_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x}))_{\text{str}} \quad (6.6)$$

We present  $Q(\mathcal{L}; \mathbf{x})$  as

$$Q(\mathcal{L}; \mathbf{x}) = \sum_{i,j=1}^L l_{ij}(\mathbf{a}) \ x_i - \dots - x_j, \quad (6.7)$$

$$l_{ij}(\mathbf{a}) \in \mathbb{Q}[[\mathbf{a}]] \subset \mathcal{Q}(\mathbb{Q}[[\mathbf{a}]]), \quad l_{ji}(\mathbf{a}) = l_{ij}(-\mathbf{a})$$

(the origin of the coefficients  $l_{ij}(\mathbf{a})$  from the haircomb graphs of  $W(\mathcal{L})$  will be explained in details in subsection 6.2).  $Z_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x})|_{x_k=0}$  is also SP, and its quadratic exponent part is given by

$$Q_k(\mathcal{L}; \mathbf{x}_{(k)}) = Q(\mathcal{L}; \mathbf{x})|_{x_k=0}. \quad (6.8)$$

Since Lemma 6.1 claims that  $Z_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x})$  is  $\text{Ad}_{\exp(y)}$ -invariant, then we are going to study the integral

$$Z_{\mathcal{QD}^C}^{\text{f}}(\mathcal{L}) = \int_{\text{FG}} d\mathbf{x}_{(k)} \left( Z_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x}) \boldsymbol{\mu}(\mathbf{x}) \right) \Big|_{x_k=0}, \quad 1 \leq k \leq L \quad (6.9)$$

(cf. eq.(5.44)), which is an invariant of the oriented marked Morse link  $\mathcal{L} \subset M$ .

**Theorem 6.3** *For  $1 \leq k \leq L$ , the function  $Z_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x})|_{x_k=0}$  is SPND (that is,  $||l_{ij}(\mathbf{a})||_k$  is non-degenerate) iff*

$$\nabla_A(\mathcal{L}; \mathbf{t}) \not\equiv 0. \quad (6.10)$$

Moreover,

$$\frac{\det' ||l_{ij}(\mathbf{a})||}{\nabla_A(\mathcal{L}; e^{\mathbf{a}})} = \left( \prod_{j=1}^L a_j \right) (1 + \mathcal{O}(\mathbf{a})). \quad (6.11)$$

*Proof.* It is sufficient to prove eq.(6.11). Our proof is based on the results of [13]. It is easy to see that

$$\mathcal{T}_{\mathfrak{g}} \hat{Q}_k(\mathcal{L}) = Q_{\mathfrak{g},k}(\mathcal{L}; \vec{\mathbf{a}}), \quad (6.12)$$

the operator  $Q_{\mathfrak{g},k}(\mathcal{L}; \vec{\mathbf{a}})$  being defined by eq.(2.67). Therefore

$$\mathcal{T}_{\mathfrak{g}} \det \hat{Q}_k(\mathcal{L}) = \det Q_{\mathfrak{g},k}(\mathcal{L}; \vec{\mathbf{a}}), \quad (6.13)$$

or, more explicitly, according to eq.(3.77),

$$\left( \prod_{\vec{\lambda} \in \Delta_+} \det ||l_{ij}((\vec{\mathbf{a}}, \vec{\lambda}))||_k \right)^2 = \det Q_{\mathfrak{g},k}(\mathcal{L}; \vec{\mathbf{a}}). \quad (6.14)$$

If  $\mathfrak{g} = su(2)$ , then  $\mathfrak{h}$  is 1-dimensional and there is only one positive root  $\vec{\lambda}$ , so relation (6.14) becomes

$$\det Q_{su(2),k}(\mathcal{L}; \mathbf{a}) = \left( \det ||l_{ij}(\mathbf{a})||_k \right)^2, \quad (6.15)$$

where we use an  $su(2)$   $\mathfrak{h}^*$  coordinate  $a = (\vec{a}, \vec{\lambda})$ . On the other hand, it was established in [13] that

$$\frac{\left( \det Q_{su(2),k}(\mathcal{L}; \mathbf{a}) \right)^{1/2}}{\nabla_A(\mathcal{L}; e^{\mathbf{a}})} = \left( \prod_{j=1}^L a_j \right) (1 + \mathcal{O}(\mathbf{a})). \quad (6.16)$$

Equation (6.11) follows from eqs.(6.15) and (6.16).  $\square$

**Remark 6.4** Equation (6.11) indicates that  $\det' ||l_{ij}(\mathbf{a})|| \Big|_{\mathbf{a}=0} = 0$ , which means that although  $Z_{\mathcal{D}^C}^{\mathcal{Q}^R}(\mathcal{L}; \mathbf{x})|_{x_k=0}$  may be non-degenerate, its predecessor  $Z_{\mathcal{B}^C}(\mathcal{L}; R_{a_1}(x_1), \dots, R_{a_L}(x_L))$  is always degenerate and therefore the integral (6.9) would not be well-defined in  $\mathcal{B}_{\mathcal{L}}^C$  (cf. Remark 4.37).

**Theorem 6.5** *The integral (6.9) does not depend on the choice of  $k$  ( $1 \leq j \leq L$ ) in eq.(6.9).*

*Proof.* This theorem is a particular case of Theorem 5.14.  $\square$

**Theorem 6.6** *The integral (6.9) is an invariant of an oriented link  $\mathcal{L} \in S^3$ , that is, it does not depend on a presentation of  $\mathcal{L}$  as a marked Morse link.*

Before we prove this theorem, we have to describe how Kontsevich integral  $Z_{\mathcal{B}}(\mathcal{L}; \mathbf{X})$  depends on the choice of marked points on the components of  $\mathcal{L}$ . Let us introduce a cyclic smooth parameter  $t$  on  $\mathcal{L}_1$ . Then  $Z_{\mathcal{B}}(\mathcal{L}; \mathbf{X}; t)$  depends explicitly on a position  $t$  of the marked point on  $\mathcal{L}_1$ .

**Lemma 6.7** *The derivative  $\partial_t Z_{\mathcal{B}}(\mathcal{L}; \mathbf{X}; t)$  has a form*

$$\partial_t Z_{\mathcal{B}}(\mathcal{L}; \mathbf{X}; t) = \text{dvd}_{X_1} \left( \xi(\mathbf{X}) Z_{\mathcal{B}}(\mathcal{L}; \mathbf{X}; t) \right), \quad (6.17)$$

where  $\xi(\mathbf{X})$  is a vector field with  $\partial_{X_1}$  component. The vector field  $\xi(\mathbf{X})$  is narrow and therefore has a polynomial (actually, at most linear) dependence on the struts.

*Proof of Theorem 6.6.* This theorem follows immediately from eq.(6.17) and Theorem 5.18.

□

*Proof of Lemma 6.7.* Consider graph algebras  $\mathcal{A}(\uparrow_X, \uparrow_Y)$  and  $\mathcal{A}(\uparrow_Z)$ . There is an  $\mathcal{A}$ -multiplication map  $m_{X,Y}^Z: \mathcal{A}(\uparrow_X, \uparrow_Y) \rightarrow \mathcal{A}(\uparrow_Z)$ . With a slight abuse of notations we will use the same notation for an  $\mathcal{A}$ -multiplication map  $m_{X,Y}^Z: B_{\{X,Y\}} \rightarrow B_{\{Z\}}$  induced on  $\mathcal{B}$  algebras by the isomorphisms  $\hat{\chi}$  (see the diagram (2.19)). We will also define  $m_{[X,Y]}^Z = m_{X,Y}^Z - m_{Y,X}^Z$ . Now consider a vector field  $\Xi(x)$ . Recall that its graph has a few  $X$  legs and exactly one  $\partial_X$  leg. The relation

$$m_{[\partial_X, X]}^X \Xi(X) = \text{dvd}_X \Xi(X) \quad (6.18)$$

follows easily from a combinatorial identity

$$Y X^n - X^n Y = \sum_{m=0}^{n-1} X^m [Y, X] X^{n-m-1}. \quad (6.19)$$

Consider two marked points  $t$  and  $t + \Delta t$  on a link component  $\mathcal{L}_1$ . They split it into two pieces (one big and one small). Thus we get a modified Morse ‘link’  $\mathcal{L}'$  in which a closed component  $\mathcal{L}$  is replaced by two segments. Consider a Kontsevich integral  $Z_{\mathcal{A}}(\mathcal{L}') \in \mathcal{A}(\uparrow_{\Delta X_1}, \uparrow_{X_1}, \dots, \uparrow_{X_L})$  and its PBW-symmetrized version  $Z_{\mathcal{B}}(\mathcal{L}'; \Delta X, \mathbf{X}) \in B_{\{\Delta X_1, X_1, \dots, X_L\}}$ . Since

$$Z_{\mathcal{A}}(\mathcal{L}'; t) = m_{\Delta X_1, X_1}^{X_1} Z_{\mathcal{A}}(\mathcal{L}'), \quad Z_{\mathcal{A}}(\mathcal{L}'; t + \Delta t) = m_{X_1, \Delta X_1}^{X_1} Z_{\mathcal{A}}(\mathcal{L}'), \quad (6.20)$$

then

$$\Delta Z_{\mathcal{B}}(\mathcal{L}; \mathbf{X}; t) = Z_{\mathcal{B}}(\mathcal{L}; \mathbf{X}; t + \Delta t) - Z_{\mathcal{B}}(\mathcal{L}; \mathbf{X}; t) = -m_{[\Delta X_1, X_1]}^{X_1} Z_{\mathcal{B}}(\mathcal{L}'; \Delta X, \mathbf{X}). \quad (6.21)$$

As usual,  $Z_{\mathcal{B}}(\mathcal{L}'; \Delta X, \mathbf{X}) = \exp \left( W(\mathcal{L}'; \Delta X; \mathbf{X}) \right)$ , where  $W(\mathcal{L}'; \Delta X; \mathbf{X})$  is narrow. We are interested only in the terms which are at most linear in  $\Delta t$ . It is easy to see from the

definition of Kontsevich integral that every  $\Delta X$  leg in the graphs of  $W(\mathcal{L}'; \Delta X; \mathbf{X})$  carries with it at least a factor of  $\Delta t$ . Therefore  $W(\mathcal{L}'; \Delta X; \mathbf{X})|_{\Delta t=0} = W(\mathcal{L}; \mathbf{X}; t)$  and the graphs whose coefficients are linear in  $\Delta t$  must have exactly one  $\Delta X_1$  leg. Thus if we take a sum of all graphs of  $W(\mathcal{L}'; \Delta X; \mathbf{X})$  whose coefficients are proportional to  $\Delta t$  and replace there  $\Delta X_1$  with  $\partial_{X_1}$ , then the result can be declared a vector field  $\xi(\mathbf{X}) \Delta t$ . Now

$$\Delta Z_B(\mathcal{L}; \mathbf{X}; t) \approx -m_{[\partial_{X_1}, X_1]}^{X_1} \left( \xi(\mathbf{X}) Z_B(\mathcal{L}; \mathbf{X}; t) \right) \Delta t \quad (6.22)$$

and eq.(6.17) follows in view of eq.(6.18) in which  $\Xi(X)$  is replaced by  $\xi(\mathbf{X}) Z_B(\mathcal{L}; \mathbf{X}; t)$ .  $\square$

Theorems 6.5 and 6.6 allow us to formulate the following

**Definition 6.8** *For a link  $\mathcal{L}$  in a rational homology sphere  $M$  such that  $\nabla_A(\mathcal{L}; \mathbf{t}) \neq 0$ , the unwheeled graph  $U(1)$ -RCC invariant  $Z_{\mathcal{QD}^c}^r(\mathcal{L})$  is defined by the integral (6.9).*

If a link has a single component ( $L = 1$ ), then there is no integration in (6.9), hence  $Z_{\mathcal{QD}^c}^r(\mathcal{L})$  is obtained from  $Z_{\mathcal{D}^c}(\mathcal{L}; \mathbf{X})$  by changing its legs from total to Cartan. In other words, for a knot  $\mathcal{K}$ ,

$$Z_{\mathcal{QD}^c}^r(\mathcal{K}) = f_{\mathcal{QH}} \circ f_{\text{Crtn}}(Z_B(\mathcal{K})), \quad (6.23)$$

or, less formally,  $Z_{\mathcal{QD}^c}^r(\mathcal{K}) = Z_B(\mathcal{K}; a)$ , since according to our conventions, replacing the total labels  $X$  of  $Z_B(\mathcal{K}; X)$  by Cartan labels  $a$  makes its total legs Cartan. Theorem 3.2,  $f_{\text{Crtn}}$  is an injection in a knot case, so no information about a knot is lost in passing from the traditional Kontsevich integral  $Z_B(\mathcal{K}; X)$  to its Cartan-restricted version  $Z_{\mathcal{QD}^c}^r(\mathcal{K})$ .

The dependence of  $Z_{\mathcal{QD}^c}^r(\mathcal{L})$  on the orientation of  $\mathcal{L}$  is quite transparent. For an oriented link  $\mathcal{L}$  let  $\mathcal{L}_{\{j\}}$  denote the same link in which we switched the orientation of the  $j$ -th component  $\mathcal{L}_j$  and let  $\bar{\mathcal{L}}$  denote the link  $\mathcal{L}$  in which we switched the orientation of all components.

**Theorem 6.9** *For an oriented link  $\mathcal{L}$  with  $\nabla_A(\mathcal{L}; \mathbf{t}) \neq 0$*

$$Z_{\mathcal{QD}^c}^r(\mathcal{L}_{\{j\}}) = P_j \left( Z_{\mathcal{QD}^c}^r(\mathcal{L}) \right), \quad Z_{\mathcal{QD}^c}^r(\bar{\mathcal{L}}) = Z_{\mathcal{QD}^c}^r(\mathcal{L}), \quad (6.24)$$

where the parity operators  $P_j$  are defined by eqs. (3.21) and (3.54).

*Proof.* It is well-known (and easy to see from its definition) how Kontsevich integral changes when orientation of link components is switched:

$$Z_B(\mathcal{L}_{\{j\}}; \mathbf{X}) = Z_B(\mathcal{L}; X_1, \dots, -X_j, \dots, X_{|\mathcal{L}|}), \quad Z_B(\bar{\mathcal{L}}; \mathbf{X}) = Z_B(\mathcal{L}; \mathbf{X}). \quad (6.25)$$

Therefore, since  $R_{-a}(x) = -R_a(x)$ , then

$$Z_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}_{\{j\}}; \mathbf{x}) = P_j \left( Z_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x}) \right), \quad Z_{\mathcal{D}^C}^{\mathcal{Q}R}(\bar{\mathcal{L}}; \mathbf{x}) = Z_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x}), \quad (6.26)$$

and eqs.(6.24) follow by applying the integral of eq.(6.9) to these equations.  $\square$

In order to make connection with  $\mathfrak{g}$ -based  $U(1)$ -RCC invariant of subsection 2.4, we have to construct a graph algebra  $U(1)$ -RCC invariant from the wheeled Kontsevich integral (2.24). We follow the same steps that led us to eq.(6.9) and introduce

$$Z_{\mathcal{QD}^C}^{\Omega, r}(\mathcal{L}) = \int_{\text{FG}} d\mathbf{x}_{(k)} \left( Z_{\mathcal{D}^C}^{\Omega, \mathcal{Q}R}(\mathcal{L}; \mathbf{x}) \boldsymbol{\mu}(\mathbf{x}) \right) \Big|_{x_k=0}, \quad 1 \leq j \leq L, \quad (6.27)$$

where again

$$Z_{\mathcal{D}^C}^{\Omega, \mathcal{Q}R}(\mathcal{L}; \mathbf{x}) = Z_{\mathcal{B}^C}^{\Omega}(\mathcal{L}; R_{a_1}(x_1), \dots, R_{a_L}(x_L)). \quad (6.28)$$

**Theorem 6.10**  $Z_{\mathcal{D}^C}^{\Omega, \mathcal{Q}R}(\mathcal{L}; \mathbf{x})$  depends neither on the choice of  $k$  ( $1 \leq k \leq L$ ) nor on the presentation of  $\mathcal{L}$  as a marked Morse link.

*Proof.* The proof of the first claim is exactly the same as the proof of Theorem 6.5. In order to prove the second claim, note that according to Lemma 4.7, the wheeling operator  $\hat{\Omega}_{\mathcal{L}}$  (which is a particular case of gluing  $\vec{\text{ad}}$ -invariant graphs) commutes with  $\text{ad}$ -divergence  $\text{dvd}_{X_1}$ . Therefore eq.(6.17) implies a similar relation for  $Z_{\mathcal{B}}^{\Omega}(\mathcal{L}; \mathbf{x}; t)$ :

$$\partial_t Z_{\mathcal{B}}^{\Omega}(\mathcal{L}; \mathbf{x}; t) = \hat{\Omega}_{\mathcal{L}} \text{dvd}_{x_1} \left( \xi(x) Z_{\mathcal{B}}(\mathcal{L}; \mathbf{X}; t) \right) = \text{dvd}_{x_1} \hat{\Omega}_{\mathcal{L}} \left( \xi(x) Z_{\mathcal{B}}(\mathcal{L}; \mathbf{X}; t) \right), \quad (6.29)$$

The rest of the proof is similar to that of Theorem 6.6.  $\square$

**Definition 6.11** For a link  $\mathcal{L}$  in a rational homology sphere  $M$  such that  $\nabla_A(\mathcal{L}; \mathbf{t}) \neq 0$ , the wheeled graph  $U(1)$ -RCC invariant is defined by the integral (6.27).

**Remark 6.12** Since graphs of wheeling operators have even numbers of legs, then wheeled Kontsevich integral still satisfies eqs.(6.25). Hence the wheeled graph  $U(1)$ -RCC invariant satisfies eqs.(6.24) of Theorem 6.9.

The wheeled graph  $U(1)$ -RCC invariant determines the Lie algebra based  $U(1)$ -RCC invariant (2.59). In order to see this, we introduce a weight system  $\mathcal{T}_{\mathfrak{g}}^{2, \hbar}$  which acts in the same way as  $\mathcal{T}_{\mathfrak{g}}^{1, \hbar}$ , except that it multiplies the graphs by  $\hbar^{\deg_2 D}$  rather than by  $\hbar^{\deg_1 D}$ . Since the exponent of eq.(2.35) comes from the *r.h.s.* of eq.(2.31) (which is the exponent of



eq. (2.32)) through the substitution  $\vec{\chi} = \hbar^{-1} \vec{\mathbf{X}}$ , and since according to eq. (2.3),  $\deg_2 D = \deg_1 D - \#(1\text{-vertices})$ , then

$$\sum_{m,n=0}^{\infty} L_{m,n}(\mathcal{L}; \vec{\mathbf{X}}) \hbar^{n-1} = \mathcal{T}_{\mathfrak{g}}^{2,\hbar} W^{\Omega}(\mathcal{L}). \quad (6.30)$$

where  $\vec{\mathbf{X}}$  are the elements of  $\mathfrak{g}^*$  which are placed at the legs of the graphs of  $W^{\Omega}(\mathcal{L})$ . Since, according to Theorem (5.7),  $\mu(x)$  is an exponential of the sum of wheel graphs, then

$$\deg_2 \mu(x) = 0. \quad (6.31)$$

Therefore we can replace  $\mathcal{T}_{\mathfrak{g}}$  by  $\mathcal{T}_{\mathfrak{g}}^{2,\hbar}$  in eq. (5.32):

$$\mu_{\vec{a}}(\vec{x}) = (-2\pi)^{-|\Delta_+|} d_{\mathfrak{g}}(\vec{a}) \mathcal{T}_{\mathfrak{g}} \mu(\vec{x}). \quad (6.32)$$

A combination of eqs. (6.30) and (6.32) indicates that eq. (2.59) can be rewritten as

$$I_{\mathfrak{g}}^r(\mathcal{L}; \vec{\mathbf{a}}; \hbar, k) = \hbar^{-|\Delta_+|} (-2\pi\hbar)^{(1-L)|\Delta_+|} \frac{d_{\mathfrak{g}}(\vec{\mathbf{a}})}{d_{\mathfrak{g}}(\vec{\rho})} \int_{[\vec{\mathbf{x}}_{(k)}=0]} d\vec{\mathbf{x}}_{(k)} \mathcal{T}_{\mathfrak{g}}^{2,\hbar} \left( Z_{\mathcal{D}^c}^{\Omega, \mathcal{Q}R}(\mathcal{L}; \mathbf{x}) \mu(\mathbf{x}) \right) \Big|_{\vec{x}_k=0}. \quad (6.33)$$

Applying eq. (4.90) to this equation we come to the following

**Theorem 6.13** *The graph algebra  $U(1)$ -RCC invariant  $Z_{\mathcal{D}^c}^{\Omega, r}(\mathcal{L})$  is ‘universal’ in the sense that it determines Lie algebra based  $U(1)$ -RCC invariants through the application of the  $\mathcal{T}_{\mathfrak{g}}^{2,\hbar}$  weight system*

$$I_{\mathfrak{g}}^r(\mathcal{L}; \vec{\mathbf{a}}; \hbar, k) = \hbar^{-|\Delta_+|} \frac{d_{\mathfrak{g}}(\vec{\mathbf{a}})}{d_{\mathfrak{g}}(\vec{\rho})} \mathcal{T}_{\mathfrak{g}}^{2,\hbar} \left( Z_{\mathcal{D}^c}^{\Omega, r}(\mathcal{L}) \right). \quad (6.34)$$

Although Lie algebra based  $U(1)$ -RCC invariants are completely determined by the universal invariant, they are still interesting in their own right: at least, in case of  $\mathfrak{g} = su(2)$  they can be derived from  $R$ -matrix type calculation rather than from Kontsevich integral, and they exhibit interesting integrality properties [12].

## 6.2 Haircomb graphs, wheels and Kontsevich integral formula for the Alexander polynomial

The Alexander polynomial of a link is hidden inside its  $U(1)$ -RCC invariant. In order to see this, first, we have to express the Alexander polynomial in terms of Kontsevich integral

of the link. Let us concentrate on the strut and circle parts of the logarithm of Kontsevich integral  $W(\mathcal{L}; \mathbf{X})$  after the substitution (6.5):

$$W_{\mathcal{D}^{\text{CR}}}(\mathcal{L}; \mathbf{x}) = \frac{1}{2}Q(\mathcal{L}; \mathbf{x}) + \frac{1}{2}F_0(\mathcal{L}; \mathbf{a}) \quad (6.35)$$

where  $Q(\mathcal{L}; \mathbf{x})$  is the strut expression (6.7) and  $F_0(\mathcal{L}; \mathbf{a}) \in \left[ \mathbb{Q}[[\mathbf{a}]] \right]^{\text{even}}$  is a new formal power series. Since series  $l_{ij}(\mathbf{a})$  and  $F_0(\mathcal{L}; \mathbf{a})$  will play an important role in our calculations, we take a short digression in order to explain explicitly how they originate from the graphs of  $W(\mathcal{L})$ .

We begin with the strut coefficients  $l_{ij}(\mathbf{a})$ . The strut exponent  $Q(\mathcal{L}; \mathbf{x})$  comes from the haircomb graphs of  $W(\mathcal{L}; \mathbf{X})$ . A *haircomb* graph is a graph which consists of a single chain connecting two 1-valent vertices and legs attached to that chain (*cf.* Fig. 1). The chain is called a *spine* of the graph and the 1-valent vertices are called *endpoints* of the haircomb. The same graph may have up to four different spines depending on the choice of end-points.

A haircomb graph of  $W(\mathcal{L}; \mathbf{X})$  may contribute to  $Q(\mathbf{x})$  in two different ways. The first way is to make a substitution  $X_i = \frac{1}{2} \text{ad}_{x_i}^2(a_i)$  at one of the legs and  $X_i = a_i$  at all other legs. Only the struts of  $W(\mathcal{L}; \mathbf{X})$  may contribute in this way (struts are also haircombs). If the strut part of  $W(\mathcal{L}; \mathbf{X})$  is

$$W_{\text{str}}(\mathcal{L}; \mathbf{X}) = \frac{1}{2} \sum_{i,j=1}^L \ell_{ij} \quad X_i \text{ --- } X_j, \quad (6.36)$$

then its first-way contribution to  $Q(\mathbf{x})$  is

$$\sum_{i,j=1}^L \ell_{ij} \quad a_i a_j \quad x_i \text{ --- } x_j. \quad (6.37)$$

The second way, in which a haircomb graph of  $W(\mathcal{L}; \mathbf{X})$  may contribute to  $Q(\mathbf{x})$ , is to perform the substitutions  $X_i = \text{ad}_{x_i}(a_i)$  at its endpoints and the substitutions  $X_i = a_i$  at all other 1-vertices. Let us quantify this contribution. For a set of non-negative integers  $\mathbf{n}$  let  $\mathbf{D}_{i,j;\mathbf{n}}$  be a set of haircomb graphs with selected spines such that their endpoints have colors  $X_i, X_j$  and they have  $n_k$  legs of color  $X_k$  attached to the spine for all  $k$ ,  $1 \leq k \leq L$ . Let  $c(D)$  be the coefficients of these haircomb graphs in an expression of  $W(\mathcal{L}; \mathbf{X})$  as an element of  $\tilde{\mathcal{B}}_{\mathcal{L}}$  (we fix the signs of  $c(D)$  by assuming that legs are attached to the right of the spine, if it is oriented from  $X_j$  to  $X_i$ ). Now we define the power series

$$l'_{ij}(\mathbf{a}) = - \sum_{\mathbf{n} \geq 0} \left( \sum_{D \in \mathbf{D}_{i,j;\mathbf{n}}} c(D) \right) a_1^{n_1} \cdots a_L^{n_L}. \quad (6.38)$$

Note that  $l'_{ji}(\mathbf{a}) = l'_{ij}(-\mathbf{a})$ . Now the second contribution of haircomb graphs to  $Q(\mathbf{x})$  is

$$\sum_{i,j=1}^L l'_{ij}(\mathbf{a}) \quad x_i - - - x_j. \quad (6.39)$$

Since a combination of contributions (6.37) and (6.39) is equal to the *r.h.s.* of eq.(5.47), then we conclude that

$$l_{ij}(\mathbf{a}) = a_i \left( a_j l'_{ij}(\mathbf{a}) + \delta_{ij} \sum_{k=1}^L a_k \ell_{ik} \right). \quad (6.40)$$

**Remark 6.14** The contribution of diagonal struts  $\ell_{ii} X_i \text{ --- } X_j$  of  $W(\mathcal{L}; \mathbf{X})$  to both expressions (6.37) and (6.39) are equal up to a sign and thus cancel each other in the final expression (6.40). Therefore, diagonal coefficients  $\ell_{ii}$  do not participate in the final expression (6.40).

The circle graph contribution  $\frac{1}{2} F_0(\mathcal{L}; \mathbf{a}) \quad \langle \text{---} \rangle$  to  $W_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x})$  comes from the wheel graphs of  $W(\mathcal{L}; \mathbf{X})$ , if we substitute  $X_i = a_i$  at all their legs. Thus let  $\mathbf{D}_{\mathbf{n}}$  be the set of wheel graphs of  $\mathcal{B}_{\mathcal{L}}$  which have  $n_i$  legs of color  $X_i$  for all  $i$ ,  $1 \leq i \leq L$ . Then

$$F_0(\mathcal{L}; \mathbf{a}) = 2 \sum_{\substack{\mathbf{n} \geq 0 \\ n_1 + \dots + n_L \in 2\mathbb{Z}}} \sum_{D \in \mathbf{D}_{\mathbf{n}}} c(D), \quad (6.41)$$

where  $c(D)$  are again the coefficients at graphs  $D$  with which they appear in  $W(\mathcal{L}; \mathbf{X})$ .

Since the tree parts of  $W(\mathcal{L}; \mathbf{X})$  and  $W^{\Omega}(\mathcal{L}; \mathbf{X})$  are the same, then similarly to eq.(6.35) we can write

$$W_{\mathcal{D}^C}^{\Omega, \mathcal{Q}R}(\mathcal{L}; \mathbf{x}) = \frac{1}{2} Q(\mathcal{L}; \mathbf{x}) + \frac{1}{2} F_0^{\Omega}(\mathcal{L}; \mathbf{a}) \quad \langle \text{---} \rangle + \text{other graphs}, \quad (6.42)$$

The circle parts of  $W_{\mathcal{D}^C}^{\Omega, \mathcal{Q}R}(\mathcal{L}; \mathbf{x})$  and  $W_{\mathcal{D}^C}^{\mathcal{Q}R}(\mathcal{L}; \mathbf{x})$  are different, because the wheeling of  $W(\mathcal{L}; \mathbf{X})$  produces new wheel graphs by gluing the struts of  $W(\mathcal{L}; \mathbf{X})$  to the legs of the wheels of  $\Omega$  defined by eq.(2.16). Therefore it follows from eqs.(2.15) and (2.7) that

$$F_0^{\Omega}(\mathcal{L}; \mathbf{a}) = F_0(\mathcal{L}; \mathbf{a}) - \log \left( \Omega_{\nabla}(\mathcal{L}; \mathbf{a}) \right), \quad (6.43)$$

where we defined a function  $\Omega_{\nabla}(\mathcal{L}; \mathbf{a})$  as

$$\Omega_{\nabla}(\mathcal{L}; \mathbf{a}) = \prod_{i=1}^L \left( \frac{\sinh \left( \frac{1}{2} \sum_{\substack{1 \leq j \leq L \\ j \neq i}} \ell_{ij} a_j \right)}{\frac{1}{2} \sum_{\substack{1 \leq j \leq L \\ j \neq i}} \ell_{ij} a_j} \right). \quad (6.44)$$

Now we can prove a theorem that we neglected to formulate in [13]:

**Theorem 6.15** *The Alexander polynomial  $\nabla_A(\mathcal{L}; \mathbf{t})$  is determined by the haircomb tree and ‘1-loop’ parts of the (logarithm of) Kontsevich integral  $W^\Omega(\mathcal{L})$ :*

$$\begin{aligned}\nabla_A(\mathcal{L}; e^{\mathbf{a}}) &= \frac{\det' ||l_{ij}(\mathbf{a})||}{\prod_{j=1}^L a_j} \exp \left( -F_0^\Omega(\mathcal{L}; \mathbf{a}) \right) \\ &= \frac{\det' ||l_{ij}(\mathbf{a})||}{\prod_{j=1}^L a_j} \Omega_\nabla(\mathcal{L}; \mathbf{a}) \exp \left( -F_0(\mathcal{L}; \mathbf{a}) \right).\end{aligned}\tag{6.45}$$

*Proof.* In view of eq. (6.43) it is sufficient to prove only the first equation of (6.45).

Since an individual polynomial  $L_{m,n}(\mathcal{L}; \vec{\mathbf{X}})$  of the *l.h.s.* of eq. (6.30) comes from the graphs  $D$  of  $W^\Omega(\mathcal{L})$  which have  $\chi(D) = n - 1$  and  $m$  legs, then eq. (6.30) implies that

$$\sum_{m=0}^{\infty} L_{m,1}(\mathcal{L}; \vec{\mathbf{a}}) = \sum_{\vec{\lambda} \in \Delta_+} F_0^\Omega(\mathcal{L}; (\vec{\mathbf{a}}, \vec{\lambda})).\tag{6.46}$$

Consider now a case of  $\mathfrak{g} = su(2)$ . Since  $su(2)$  has only one positive root, then (in terms of the  $su(2)$   $\mathfrak{h}^*$  coordinate  $a = (\vec{a}, \vec{\lambda})$ )

$$\sum_{m=0}^{\infty} L_{m,0}(\mathcal{L}; \mathbf{a}) = F_0^\Omega(\mathcal{L}; \mathbf{a}).\tag{6.47}$$

Then eq. (6.45) follows easily from the combination of equations (2.75), (6.15) and (6.46).  $\square$

We give a more direct proof of Theorem 6.15 in Appendix A. It relies neither on Cartan and root edges, nor on the properties of the Jones polynomial, but rather uses a relation between the Alexander polynomial and the supergroup  $U(1|1)$ .

Note that neither  $\det' ||l_{ij}(\mathbf{a})||$ , nor  $F_0(\mathcal{L}; \mathbf{a})$  is the invariant of an ‘unmarked’ link  $\mathcal{L}$ . Only their combination (6.45) is independent of the ‘Morse marking’.

According to eq. (6.40), the series  $l_{ij}(\mathbf{a})$  is proportional to both  $a_i$  and  $a_j$  and, as a result, the determinant  $\det' ||l_{ij}(\mathbf{a})||$  of eq. (6.45) is divisible by  $\left(\prod_{j=1}^L a_j\right)^{-1}$  in  $\mathbb{Q}[[\mathbf{a}]]$  (if  $L > 1$ ):

$$\left(\prod_{j=1}^L a_j\right)^{-1} \det' ||l_{ij}(\mathbf{a})|| \in \mathbb{Q}[[\mathbf{a}]].\tag{6.48}$$

In view of the relation between the tree part of the Kontsevich integral and Milnor’s linking numbers of the link established by N. Habegger and G. Masbaum [5], it is easy to see that the factorization (6.45) of the Alexander polynomial  $\nabla_A(\mathcal{L}; e^{\mathbf{a}})$  into the determinant (6.48) and the exponential of  $-F_0^\Omega(\mathcal{L}; \mathbf{a})$  corresponds to that of Theorem 2 of J. Levine’s paper [9].

### 6.3 Alexander polynomial and the structure of the universal $U(1)$ -RCC invariant

Let us consider the structure of the graph invariant  $Z_{\mathcal{QD}^C}^r(\mathcal{L})$  defined by eq. (6.9) in more details. Let  $\mathbf{D}'_{\mathcal{L}} \subset \mathbf{D}_{\mathcal{L}}$  be the set of connected graphs of  $\mathbf{D}_{\mathcal{L}}$  which have no legs and such that their  $\chi$  is strictly positive. Also, we introduce a notation

$$\tilde{\nabla}_A(\mathcal{L}; \mathbf{a}) = \left( \prod_{j=1}^L a_j \right) \nabla_A(\mathcal{L}; e^{\mathbf{a}}). \quad (6.49)$$

**Theorem 6.16** *The  $U(1)$ -RCC invariant  $Z_{\mathcal{QD}^C}^r(\mathcal{L})$  can be presented in the form*

$$Z_{\mathcal{QD}^C}^r(\mathcal{L}) = \exp \left( W_{\mathcal{QD}^C}^r(\mathcal{L}) \right). \quad (6.50)$$

where

$$\begin{aligned} W_{\mathcal{QD}^C}^r(\mathcal{L}) = & \frac{1}{2} \sum_{i,j=1}^L \ell_{ij} a_i a_j \circ - \frac{1}{2} \log \left( \tilde{\nabla}_A(\mathcal{L}; \mathbf{a}) \right) \langle \bigcirc \rangle \\ & + \sum_{D \in \mathbf{D}'_{\mathcal{L}}} \frac{F_D(\mathcal{L})}{\prod_{e \in \mathbf{E}(\tilde{D})} \tilde{\nabla}_A(\mathcal{L}; \hat{e})} \end{aligned} \quad (6.51)$$

and

$$F_D(\mathcal{L}) \in \mathcal{H}(D). \quad (6.52)$$

*Proof.* Consider the logarithm of the integrand of eq. (6.9)

$$\log \left( Z_{\mathcal{D}^C}^{\mathcal{QR}}(\mathcal{L}; \mathbf{x}) \mu(\mathbf{x}) \right) = W_{\mathcal{D}^C}^{\mathcal{QR}}(\mathcal{L}; \mathbf{x}) + \omega(\mathbf{x}), \quad (6.53)$$

where  $\omega(\mathbf{x}) = \sum_{j=1}^L \omega(x_j)$ . Let us define  $W_{\mathcal{D}^C}^{\text{s,FD}}(\mathcal{L}; \mathbf{x}) \in \mathcal{D}_{\mathcal{L}}^C$  by the equation

$$\begin{aligned} & W_{\mathcal{D}^C}^{\mathcal{QR}}(\mathcal{L}; \mathbf{x}) + \omega(\mathbf{x}) \\ & = \frac{1}{2} \sum_{i,j=1}^L \ell_{ij} a_i a_j \circ + \frac{1}{2} F_0(\mathcal{L}; \mathbf{a}) \langle \bigcirc \rangle + \frac{1}{2} Q(\mathcal{L}; \mathbf{x}) + W_{\mathcal{D}^C}^{\text{s,FD}}(\mathcal{L}; \mathbf{x}) \end{aligned} \quad (6.54)$$

as the ‘least trivial’ part of this logarithm. Since  $Z_{\mathcal{D}^C}(\mathcal{L}; \mathbf{a})$  does not contain the graph  $\langle \bigcirc \rangle$ , then  $W_{\mathcal{D}^C}^{\text{s,FD}}(\mathcal{L}; \mathbf{x})$  contains only the graphs  $D$  such that either  $D$  has at least three legs, or  $\chi(D) = 0$  and  $D$  has at least one leg, or  $\chi(D) \geq 1$ .

The definition (4.84) applied to the integral (6.9) yields

$$Z_{\mathcal{QD}^C}^r(\mathcal{L}) = \exp \frac{1}{2} \left( \sum_{i,j=1}^L \ell_{ij} a_i a_j \circ + F_0(\mathcal{L}; \mathbf{a}) \right) \left( \det' \hat{Q}(\mathcal{L}) \right)^{-1/2} \text{FD}(\mathcal{L}), \quad (6.55)$$

where by definition

$$\left( \det' \hat{Q}(\mathcal{L}) \right)^{-1/2} = \exp \left( -\frac{1}{2} \log \det' \|l_{ij}(\mathbf{a})\| \right), \quad (6.56)$$

while

$$\text{FD}(\mathcal{L}) = \exp \left( W_{\mathcal{D}^C}^{\text{s,FD}}(\mathcal{L}; \mathbf{x}) \Big|_{x_k=0} \right) \mathfrak{b}_{\mathbf{x}_{(k)}, \mathbf{x}_{(k)}; \text{LR}} \exp \left( -\frac{1}{2} Q_k^{-1}(\mathcal{L}; \mathbf{x}_{(k)}) \right). \quad (6.57)$$

In view of eq.(6.45),

$$\tilde{\nabla}_A(\mathcal{L}; \mathbf{a}) = \det' \|l_{ij}(\mathbf{a})\| \Omega_{\nabla}(\mathcal{L}; \mathbf{a}) \exp \left( -F_0(\mathcal{L}; \mathbf{a}) \right), \quad (6.58)$$

so we can rewrite eq.(6.55) as

$$Z_{\mathcal{QD}^C}^r(\mathcal{L}) = \exp \frac{1}{2} \left( \sum_{i,j=1}^L \ell_{ij} a_i a_j \circ - \log \left( \frac{\tilde{\nabla}_A(\mathcal{L}; \mathbf{a})}{\Omega_{\nabla}(\mathcal{L}; \mathbf{a})} \right) \right) \text{FD}(\mathcal{L}). \quad (6.59)$$

Now it remains to determine the structure of  $\text{FD}(\mathcal{L})$ . Both exponents in the *r.h.s.* of eq. (6.57) are narrow, therefore (see, *e.g.* [3])  $\text{FD}(\mathcal{L})$  is also an exponential of a narrow exponent, whose terms are constructed by gluing the struts of  $Q(\mathcal{L}; \mathbf{x})$  to *all* legs of the graphs of  $W_{\mathcal{D}^C}^{\text{s,FD}}(\mathcal{L}; \mathbf{x})$  (possibly, to more than one graph simultaneously, but in such a way that a resulting graph is always connected).

$$\text{FD}(\mathcal{L}) = \exp \left( \text{FD}_{\text{nrw}}(\mathcal{L}) \right). \quad (6.60)$$

Therefore  $\text{FD}_{\text{nrw}}(\mathcal{L})$  is a sum of contributions of connected legless graphs and the conditions on the graphs of  $W_{\mathcal{D}^C}^{\text{s,FD}}(\mathcal{L}; \mathbf{x})$  imply that  $\chi$  of the graphs of  $\text{FD}_{\text{nrw}}(\mathcal{L})$  must be not less than 1. Thus  $\text{FD}_{\text{nrw}}(\mathcal{L})$  contains only the graphs of  $\mathbf{D}'_{\mathcal{L}}$ , and it remains to check the denominators of their contributions.

Since  $Z_{\mathcal{D}^C}^{\mathcal{QR}}(\mathcal{L}; \mathbf{x})$ ,  $\boldsymbol{\mu}(\mathbf{x}) \in \mathcal{D}_{\mathcal{L}}^C \subset \mathcal{QD}_{\mathcal{L}}^C$ , then  $W_{\mathcal{D}^C}^{\text{s,FD}}(\mathcal{L}; \mathbf{x}) \in \mathcal{D}_{\mathcal{L}}^C \subset \mathcal{QD}_{\mathcal{L}}^C$  and the struts of  $Q_k^{-1}(\mathcal{L}; \mathbf{x}_{(k)})$  are the only source of denominators in  $\text{FD}_{\text{nrw}}(\mathcal{L})$ . These denominators are  $\det' \|l_{ij}(\mathbf{a})\|$  which appear in the expression for the elements of the inverse matrix  $\|l_{ij}(\mathbf{a})\|_k^{-1}$ . Thus, if a root edge  $e$  of a graph  $D$  in the expression for  $\text{FD}_{\text{nrw}}(\mathcal{L})$  is a glued strut of  $Q_k^{-1}(\mathcal{L}; \mathbf{x}_{(k)})$ , then it carries the denominator  $\det' \|l_{ij}(\hat{\mathbf{e}})\|$ , and the expression for  $\text{FD}_{\text{nrw}}(\mathcal{L})$  has a form

$$\text{FD}_{\text{nrw}}(\mathcal{L}) = \sum_{D \in \mathbf{D}'_{\mathcal{L}}} \frac{\tilde{F}_D(\mathcal{L})}{\prod_{e \in \mathbf{E}(\tilde{D})} \det' \|l_{ij}(\hat{\mathbf{e}})\|}. \quad (6.61)$$

If we multiply the numerators and denominators of the summands in the *r.h.s.* of this equation by  $\prod_{e \in \mathbf{E}(\tilde{D})} \exp\left(-F_0^\Omega(\mathcal{L}; \hat{\mathbf{e}})\right)$ , then in view of eq.(6.58) they take the form of the summands of eq.(6.51) if we set

$$F_D(\mathcal{L}) = \tilde{F}_D(\mathcal{L}) \prod_{e \in \mathbf{E}(\tilde{D})} \exp\left(-F_0^\Omega(\mathcal{L}; \hat{\mathbf{e}})\right). \quad (6.62)$$

□

Since  $W^\Omega(\mathcal{L}; \mathbf{X})$  has the same tree part as  $W(\mathcal{L}; \mathbf{x})$ , while their circle parts are related by eq. (6.43), then it is easy to work out a formula for  $Z_{\mathcal{D}^C}^{\Omega, r}(\mathcal{L})$  which is similar to eqs.(6.50), (6.51):

$$Z_{\mathcal{D}^C}^{\Omega, r}(\mathcal{L}) = \exp\left(W_{\mathcal{D}^C}^{\Omega, r}(\mathcal{L})\right). \quad (6.63)$$

where

$$\begin{aligned} W_{\mathcal{D}^C}^{\Omega, r}(\mathcal{L}) &= \frac{1}{2} \sum_{i,j=1}^L \ell_{ij} a_i a_j \circ - \frac{1}{2} \log\left(\tilde{\nabla}_A(\mathcal{L}; \mathbf{a})\right) \left(\begin{array}{c} \circ \\ \circ \end{array}\right) \\ &\quad + \sum_{D \in \mathbf{D}'_{\mathcal{L}}} \frac{F_D^\Omega(\mathcal{L})}{\prod_{e \in \mathbf{E}(\tilde{D})} \tilde{\nabla}_A(\mathcal{L}; \hat{\mathbf{e}})} \end{aligned} \quad (6.64)$$

and

$$F_D^\Omega(\mathcal{L}) \in \mathcal{H}(D). \quad (6.65)$$

*Proof of Theorem 2.8.* Equation (2.77) is derived from (the exponent of) eq.(6.63) with the help of eq.(6.34), which relates the  $\mathfrak{g}$ -based and universal  $U(1)$ -RCC invariants. Indeed, it is easy to verify that

$$\mathcal{T}_{\mathfrak{g}}^{2, \hbar} \exp\left(\frac{1}{2} \sum_{i,j=1}^L \ell_{ij} a_i a_j \circ\right) = e^{\frac{1}{\hbar} \text{lk}(\mathcal{L}, \vec{\mathbf{a}})}, \quad (6.66)$$

$$\mathcal{T}_{\mathfrak{g}}^{2, \hbar} \exp\left(\frac{1}{2} \log \tilde{\nabla}_A(\mathcal{L}; \mathbf{a}) \left(\begin{array}{c} \circ \\ \circ \end{array}\right)\right) = F_{\nabla, \mathfrak{g}}(\mathcal{L}, \vec{\mathbf{a}}). \quad (6.67)$$

Also

$$\log \text{FD}(\mathcal{L}; \vec{\mathbf{a}}) = \sum_{n=1}^{\infty} \hbar^n \sum_{\substack{D \in \mathbf{D}'_{\mathcal{L}} \\ \deg_2(D)=n}} \mathcal{T}_{\mathfrak{g}}\left(\frac{F_D^\Omega(\mathcal{L})}{\prod_{e \in \mathbf{E}(\tilde{D})} \tilde{\nabla}_A(\mathcal{L}; \hat{\mathbf{e}})}\right). \quad (6.68)$$

The calculation of  $\mathcal{T}_{\mathfrak{g}}\left(\frac{F_D^\Omega(\mathcal{L})}{\prod_{e \in \mathbf{E}(\tilde{D})} \tilde{\nabla}_A(\mathcal{L}; \hat{\mathbf{e}})}\right)$  is performed according to the definition (3.75). Each assignment of roots  $c$  converts a denominator  $\prod_{e \in \mathbf{E}(\tilde{D})} \tilde{\nabla}_A(\mathcal{L}; \hat{\mathbf{e}})$  into

$\prod_{e \in \mathbf{E}(\tilde{D})} \tilde{\nabla}_A(\mathcal{L}; (\vec{\mathbf{a}}, c(e)))$ . Since a graph  $D \in \mathbf{D}'_{\mathcal{L}}$  has  $3\chi(D)$  edges, then  $[d_{\mathfrak{g}}(\vec{\mathbf{a}})F_{\nabla, \mathfrak{g}}(\mathcal{L}, \vec{\mathbf{a}})]^{3n}$  may serve as a common denominator for all graphs with  $\chi(D) = n$ . Hence we can present the summand of eq.(6.68) in the form

$$\sum_{\substack{D \in \mathbf{D}'_{\mathcal{L}} \\ \deg_2(D) = n}} \tau_{\mathfrak{g}} \left( \frac{F_D^{\Omega}(\mathcal{L})}{\prod_{e \in \mathbf{E}(\tilde{D})} \tilde{\nabla}_A(\mathcal{L}; \hat{\mathbf{e}})} \right) = \frac{p_n(\mathcal{L}; \vec{\mathbf{a}})}{[d_{\mathfrak{g}}(\vec{\mathbf{a}})F_{\nabla, \mathfrak{g}}(\mathcal{L}, \vec{\mathbf{a}})]^{3n}}, \quad p_n(\mathcal{L}; \vec{\mathbf{a}}) \in \mathbb{Q}[[\vec{\mathbf{a}}]], \quad (6.69)$$

and this proves eq.(2.77).  $\square$

**Remark 6.17** The power of the Alexander polynomial in denominators of eq.(1.4) is  $2n$  rather than  $3n$  as in eq.(2.80), because  $su(2)$  has only two roots. Therefore, for a graph  $D \in \mathbf{D}'_{\mathcal{L}}$  to produce a non-zero contribution to the  $su(2)$ -based  $U(1)$ -RCC invariant, at least one edge at every 3-vertex must be Cartan. As a result, out of  $3\chi D$  edges only  $2\chi D$  edges are root and  $[d_{\mathfrak{g}}(\vec{\mathbf{a}})F_{\nabla, \mathfrak{g}}(\mathcal{L}, \vec{\mathbf{a}})]^{2n}$  may serve as common denominator.

## 6.4 The rationality conjecture

The rational structure of the summands in eq.(1.4) for the  $su(2)$   $U(1)$ -RCC invariant suggests that the wheeled universal  $U(1)$ -RCC invariant also has a more restrictive form than (6.64). In order to describe it, we have to define the graph analogues of the variables  $\mathbf{t} = q^{\alpha}$  which appear as arguments of the polynomials  $P_n$  of eq.(1.4). Therefore, we come back to the definition of the spaces  $\mathcal{QH}(D)$  for graphs  $D \in \mathbf{D}_{\mathfrak{L}, \mathfrak{L}'_c}$ .

Let  $\mathbf{D}'_{\mathfrak{L}, \mathfrak{L}'_c} \subset \mathbf{D}_{\mathfrak{L}, \mathfrak{L}'_c}$  be the set of ‘normal’ (1,3)-trivalent graphs (we exclude dots and circles). This time, for a graph  $D \in \mathbf{D}'_{\mathfrak{L}, \mathfrak{L}'_c}$  we consider a group algebra

$$\tilde{H}_{\text{exp}}^1(D) = \mathbb{Z}(H^1(\tilde{D}, \partial\tilde{D}; \mathbb{Z})) \quad (6.70)$$

instead of the symmetric algebra  $S^* \tilde{H}^1(D)$ . Similarly to eq.(3.16) we define

$$\mathcal{H}_{\mathfrak{L}'_c}^{\text{exp}}(D) = ((\tilde{H}_{\text{exp}}^1(D))^{\otimes |\mathfrak{L}'_c|})^{G_D}, \quad (6.71)$$

where the action of the symmetry group  $G_D$  on  $(\tilde{H}_{\text{exp}}^1(D))^{\otimes |\mathfrak{L}'_c|}$  is defined in an obvious way (it includes a sign coming from a possible change of cyclic order at 3-vertices). Then we define the algebra  $\mathcal{QH}_{\mathfrak{L}'_c}^{\text{exp}}(D)$  as an extension of this algebra, which permits denominators of the form

$$\prod_{e \in \mathbf{E}(\tilde{D})} p_e((\hat{e}_a)_{a \in \mathfrak{L}'_c}), \quad p_e(x_1, \dots, x_{|\mathfrak{L}'_c|}) \in \mathbb{Q}[x_1^{\pm 1}, \dots, x_{|\mathfrak{L}'_c|}^{\pm 1}], \quad (6.72)$$



and the graph algebra

$$\mathcal{QE}_{\mathfrak{L}, \mathfrak{L}'_c}^C = \bigoplus_{D \in \mathbf{D}'_{\mathfrak{L}, \mathfrak{L}'_c}} \mathcal{QH}_{\mathfrak{L}'_c}^{\exp}(D). \quad (6.73)$$

As usual, when  $\mathfrak{L}'_c$  is well-known, we may drop it from our notations. Also, when working with a link  $\mathcal{L}$ , we assume that  $\mathfrak{L}'_c = \{\mathbf{a}\}$ .

Next, we define the IHX ideal  $\mathcal{QE}_{\mathfrak{L}, \text{IHx}}^C$  similarly to the definition of  $\tilde{\mathcal{D}}_{\mathfrak{L}, \text{IHx}}^C$  and  $\mathcal{QD}_{\mathfrak{L}, \text{IHx}}^C$  and finally we introduce the algebra

$$\mathcal{QE}_{\mathfrak{L}}^C = \mathcal{QE}_{\mathfrak{L}}^C / \mathcal{QE}_{\mathfrak{L}, \text{IHx}}^C. \quad (6.74)$$

We expect the latter algebra to appear in the theory of properly defined loop-filtered finite type invariants of links.

The algebras  $\mathcal{QD}_{\mathfrak{L}}^C$  and  $\mathcal{QE}_{\mathfrak{L}}^C$  are related. For an integer number  $o$  we define an injection

$$\tilde{H}_{\exp}^1(D) \xrightarrow{F_o} S^* \tilde{H}^1(D) \quad (6.75)$$

by its action on the edges  $\hat{e} \in \tilde{H}_{\exp}^1(D)$ :  $F_o(\hat{e}) = \exp(o\hat{e})$ . For a set of integers  $\mathbf{o} = (o_1, \dots, o_{|\mathfrak{L}'_c|})$  we define an injection

$$\left( \tilde{H}_{\exp}^1(D) \right)^{\otimes |\mathfrak{L}'_c|} \xrightarrow{F_{\mathbf{o}}} (S^* \tilde{H}^1(D))^{\otimes |\mathfrak{L}'_c|}, \quad F_{\mathbf{o}} = \bigotimes_{j=1}^{|\mathfrak{L}'_c|} F_{o_j} \quad (6.76)$$

and project it to an injection

$$\mathcal{H}_{\mathfrak{L}'_c}^{\exp}(D) \xrightarrow{F_{\mathbf{o}}} \mathcal{H}_{\mathfrak{L}'_c}(D). \quad (6.77)$$

This injection can be further extended to

$$\mathcal{QH}_{\mathfrak{L}'_c}^{\exp}(D) \xrightarrow{F_{\mathbf{o}}} \mathcal{QH}_{\mathfrak{L}'_c}(D) \quad \text{and} \quad \mathcal{QE}_{\mathfrak{L}, \mathfrak{L}'_c}^C \xrightarrow{F_{\mathbf{o}}} \mathcal{QD}_{\mathfrak{L}, \mathfrak{L}'_c}^C. \quad (6.78)$$

Obviously,  $F_{\mathbf{o}}(\mathcal{QE}_{\mathfrak{L}, \text{IHx}}^C) \subset \mathcal{QD}_{\mathfrak{L}, \text{IHx}}^C$ , so there is a map

$$\mathcal{QE}_{\mathfrak{L}, \mathfrak{L}'_c}^C \xrightarrow{F_{\mathbf{o}}} \mathcal{QD}_{\mathfrak{L}, \mathfrak{L}'_c}^C. \quad (6.79)$$

**Conjecture 6.18** *We conjecture the relation*

$$F_{\mathbf{o}}(\mathcal{QE}_{\mathfrak{L}}^C) \cap \mathcal{QD}_{\mathfrak{L}, \text{IHx}}^C = F_{\mathbf{o}}(\mathcal{QE}_{\mathfrak{L}, \text{IHx}}^C), \quad (6.80)$$

*which implies that the map (6.77) is an injection.*

**Conjecture 6.19 (rationality)** *Let  $\mathcal{L}$  be a link with at least 2 components in a rational homology sphere such that  $\nabla_A(\mathcal{L}; \mathbf{t}) \neq 0$ . Let  $\mathbf{o}$  denote the orders of link components as elements of  $H_1(M; \mathbb{Z})$ . Then for every graph  $D \in \mathbf{D}'_{\mathcal{L}}$  there exists an element  $P_D(\mathcal{L}) \in \mathcal{H}_{\mathcal{L}}^{\text{exp}}(D)$  such that*

$$W_{\mathcal{QD}^c}^{\Omega, r}(\mathcal{L}) = \frac{1}{2} \sum_{i,j=1}^L \ell_{ij} a_i a_j \circ - \frac{1}{2} \log \left( \tilde{\nabla}_A(\mathcal{L}; \mathbf{a}) \right) \left( \cdot \right) + F_{\mathbf{o}} \left( \sum_{D \in \mathbf{D}'_{\mathcal{L}}} \frac{P_D(\mathcal{L})}{\prod_{e \in \mathbf{E}(\tilde{D})} \nabla_A(\mathcal{L}; \hat{\mathbf{e}})} \right). \quad (6.81)$$

**Remark 6.20** We expect that the same conjecture holds also for  $W_{\mathcal{QD}^c}^r(\mathcal{L})$ .

**Remark 6.21** If the map (6.77) is an injection, as implied by Conjecture 6.18, then the element

$$\sum_{D \in \mathbf{D}'_{\mathcal{L}}} \frac{P_D(\mathcal{L})}{\prod_{e \in \mathbf{E}(\tilde{D})} \nabla_A(\mathcal{L}; \hat{\mathbf{e}})} \in \mathcal{QE}_{\mathcal{L}}^c \quad (6.82)$$

is an invariant of the oriented link  $\mathcal{L}$  (actually, a similar statement for knots was proved in [4] without proving the injectivity of (6.77)).

*Proof of Conjecture 2.9.* Equation (2.80) is derived from eq.(6.81) in exactly the same way as eq.(2.77) is derived from eq.(6.64).  $\square$

**Acknowledgements.** I am very thankful to S. Garoufalidis, R. Lawrence, D. Thurston and A. Vaintrob for discussing this work. I am especially thankful to D. Thurston for helping me to improve the exposition. This work was supported by NSF Grants DMS-0196235 and DMS-0196131.

## A A superalgebra $u(1|1)$ weight system and the Alexander polynomial of a link

The proof of the formula (6.45) which expresses the Alexander polynomial of a link in terms of its Kontsevich integral, is too convoluted. It uses the ‘rational’ expansion of the colored Jones polynomial described in [12] plus the equivalence of this expansion to the  $su(2)$ -based  $U(1)$ -RCC invariant established in [13]. One would certainly expect that a simple formula (6.45)

has a short direct proof. Indeed, such proofs exist and they are based on calculations with the weight system which produces the Alexander polynomial of a link from its Kontsevich integral. One possible approach is to describe this weight system in terms of spanning trees of 3-valent graphs. This calculation is being carried out by G. Masbaum and A. Vaintrob [10]. We suggest an alternative (and, perhaps, simpler) approach which is based on a relation between the Alexander polynomial and the  $U(1|1)$  Lie supergroup. Kauffman and Saleur [6] were the first to notice that if the  $U_q U(1|1)$  supergroup  $R$ -matrix is used instead of the  $SU_q(2)$   $R$ -matrix in the formula for the Jones polynomial, then one obtains the Alexander polynomial. Following this idea, A. Vaintrob [16] showed that an application of the  $u(1|1)$  weight system to Kontsevich integral of a link also produces the Alexander polynomial. Our formula (6.45) may be thought of as a result of calculating a fermionic (Berezin) integral in the Reshetikhin formula for  $u(1|1)$ . However, since we are not aware of Kirillov's integral formula for superalgebra characters, which would have immediately proved eq. (6.45), then we will do the corresponding weight calculations explicitly.

The superalgebra  $u(1|1)$  has four generators: two bosonic ones  $E, F$  and two fermionic ones  $\Psi^+, \Psi^-$ . The non-zero (super-)commutation relations are

$$[F, \Psi^\pm] = \pm \Psi^\pm, \quad \{\Psi^+, \Psi^-\} = E, \quad (\text{A.1})$$

where  $\{\Psi^+, \Psi^-\} = \Psi^+ \Psi^- + \Psi^- \Psi^+$ . The non-degenerate Killing form  $(\cdot, \cdot)$  is symmetric on bosonic generators, but anti-symmetric on fermionic ones:

$$(F, E) = (E, F) = 1, \quad (\Psi^+, \Psi^-) = -(\Psi^-, \Psi^+) = 1. \quad (\text{A.2})$$

Thus the non-zero components of the inverse metric tensor  $h^{-1} \in S^2 u(1|1)$  and structure constant tensor  $f \in \Lambda^3 \left( u(1|1) \right)^*$  are

$$(h^{-1})^{EF} = (h^{-1})^{FE} = (h^{-1})^{-+} = 1, \quad (h^{-1})^{+-} = -1, \quad (\text{A.3})$$

$$f_{F+-} = f_{F-+} = f_{+-F} = f_{-+F} = 1, \quad f_{+F-} = f_{-F+} = -1. \quad (\text{A.4})$$

The definition of the maps  $\mathcal{T}_{\mathfrak{g}}$  of (2.10) for a superalgebra  $\mathfrak{g}$  is slightly more complicated than for an ordinary Lie algebra. For a graph  $D \in \mathcal{A}(\uparrow_{\mathcal{L}})$  choose a ‘fermionic’ subgraph  $D_F$ , which consists of mutually nonintersecting chains. These chains may be either closed (cycles) or end at 1-vertices of  $D$ . We call such 1-vertices fermionic. Let  $n_F$  be the number of edges in  $D_F$  (we call them fermionic) and let  $n_B$  be the number of edges in  $D \setminus D_F$  (we call them bosonic).

For a pair  $D, D_F$  we define an element  $\mathcal{T}_{\mathfrak{g}}(D, D_F) \in ((U\mathfrak{g})^{\otimes |\mathcal{L}|})^{\mathfrak{g}}$ . First of all, we choose an orientation on all chains of  $D_F$  (ultimately  $\mathcal{T}_{\mathfrak{g}}(D, D_F)$  will not depend on it). As a

linear space  $\mathfrak{g}$  splits into a direct sum of its bosonic and fermionic subspaces  $\mathfrak{g} = \mathfrak{g}_B \oplus \mathfrak{g}_F$ . Correspondingly, the inverse Killing form tensor  $h_{\mathfrak{g}}^{-1}$  also splits:  $h_{\mathfrak{g}}^{-1} = h_{\mathfrak{g},B}^{-1} + h_{\mathfrak{g},F}^{-1}$ , where  $h_{\mathfrak{g},B}^{-1} \in S^2 \mathfrak{g}_B$  and  $h_{\mathfrak{g},F}^{-1} \in \bigwedge^2 \mathfrak{g}_F$ . Thus we assign  $h_{\mathfrak{g},B}^{-1}$  to the bosonic edges and  $h_{\mathfrak{g},F}^{-1}$  to the oriented fermionic edges of  $D$  in order to contract them with the  $\mathfrak{g}$  structure tensors assigns to 3-vertices, as prescribed by the index contraction map (2.11). This contraction produces an element  $\mathcal{C}_D \left( f_{\mathfrak{g}}^{\otimes n_3(D)} \otimes (h_{\mathfrak{g},B}^{-1})^{\otimes n_B} \otimes (h_{\mathfrak{g},F}^{-1})^{\otimes n_F} \right) \in ((U\mathfrak{g})^{\otimes |\mathcal{L}|})^{\mathfrak{g}}$ , which we still have to multiply by sign factors in order to get  $\mathcal{T}_{\mathfrak{g}}(D, D_F)$ . The first sign factor is  $(-1)^{\#D_F}$ , where  $\#D_F$  is the number of connected components of  $D_F$ . When passing through 3-vertices, fermionic chains induce cyclic order on incident edges. Let  $n_{\text{cycl}}$  be the number of 3-vertices on fermionic chains on which the induces order is opposite to that of the original order of  $D$ . Then the second sign factor is  $(-1)^{n_{\text{cycl}}}$ . In order to define the third ordering we pick a linear order on the set of open fermionic chains. Since these chains are oriented, this order induces a linear order on the set of fermionic 1-vertices (the final vertex of an open chain is considered to be immediately following the initial vertex of that chain). The same set of fermionic 1-vertices has an alternative linear order coming from the linear ordering of the elements of the set  $\mathcal{L}$  and from the orientation of the segments of  $\uparrow_{\mathcal{L}}$ . Let  $n_{\text{lin}}$  be the number of pairs of fermionic 1-vertices, on which both orders differ. Then the third sign factor is  $(-1)^{n_{\text{lin}}}$ . Thus we define

$$\mathcal{T}_{\mathfrak{g}}(D, D_F) = (-1)^{\#D_F + n_{\text{cycl}} + n_{\text{lin}}} \mathcal{C}_D \left( f_{\mathfrak{g}}^{\otimes n_3(D)} \otimes (h_{\mathfrak{g},B}^{-1})^{\otimes n_B} \otimes (h_{\mathfrak{g},F}^{-1})^{\otimes n_F} \right), \quad (\text{A.5})$$

$$\mathcal{T}_{\mathfrak{g}}(D) = \sum_{D_F \subset D} \mathcal{T}_{\mathfrak{g}}(D, D_F), \quad (\text{A.6})$$

where the sum in the latter formula goes over all possible fermionic subgraphs of  $D$ .

For an element  $x \in ((U\mathfrak{g})^{\otimes |\mathcal{L}|})^{\mathfrak{g}}$  consider its representation in a  $\mathfrak{g}$ -module  $V_{\vec{\alpha}} = V_{\vec{\alpha}_1} \otimes \cdots \otimes V_{\vec{\alpha}_{|\mathcal{L}|}}$ , where  $V_{\vec{\alpha}_j}$  are irreducible  $\mathfrak{g}$ -modules with (shifted) highest weights  $\vec{\alpha}_j$ . A supertrace over a linear superspace  $V = V_B \oplus V_F$  is a difference between the traces over its bosonic and fermionic subspaces:  $\text{STr}_V = \text{Tr}_{V_B} - \text{Tr}_{V_F}$ . For  $1 \leq k \leq |\mathcal{L}|$ , let  $\text{STr}_{V_{\vec{\alpha}};k} x \in \text{End}(V_{\vec{\alpha}_k})$  denote the supertrace of  $x$  over all the spaces  $V_{\vec{\alpha}_j}$  except the space  $V_{\vec{\alpha}_k}$ . Since  $x$  is  $\mathfrak{g}$ -invariant, then so is  $\text{STr}_{V_{\vec{\alpha}};k} x$ . Since we assumed that  $V_{\vec{\alpha}_k}$  is irreducible, this means that  $\text{STr}_{V_{\vec{\alpha}};k} x = CI$ , where  $I$  is the identity operator acting on  $V_{\vec{\alpha}_k}$ , while  $C$  is a constant. We will denote this constant as  $\text{STr}_{\vec{\alpha};k} x$ . It can be presented as a complete trace over  $V_{\vec{\alpha}}$ : if we choose  $O \in \text{End}(V_{\vec{\alpha}_k})$  such that  $\text{STr}_{V_{\vec{\alpha}};k} O = 1$ , and denote  $O_k = I^{\otimes (k-1)} \otimes O \otimes I^{\otimes (|\mathcal{L}|-k-1)} \in \text{End } V_{\vec{\alpha}}$ , then

$$\text{STr}_{\vec{\alpha};k} x = \text{STr}_{V_{\vec{\alpha}}} O_k x. \quad (\text{A.7})$$

Now we turn back to the case of  $\mathfrak{g} = u(1|1)$ . Consider a family of 2-dimensional  $u(1|1)$ -modules  $V_a$  parametrized by a formal parameter  $a$ . Their basis is formed by two vectors:

$|a, +\frac{1}{2}\rangle$  (boson) and  $|a, -\frac{1}{2}\rangle$  (fermion), the action of the algebra generators being

$$\begin{aligned} E|a, \pm\frac{1}{2}\rangle &= a|a, \pm\frac{1}{2}\rangle, & F|a, \pm\frac{1}{2}\rangle &= \pm\frac{1}{2}|a, \pm\frac{1}{2}\rangle, \\ \Psi^-|a, +\frac{1}{2}\rangle &= |a, -\frac{1}{2}\rangle, & \Psi^-|a, -\frac{1}{2}\rangle &= 0, \\ \Psi^+|a, -\frac{1}{2}\rangle &= a|a, +\frac{1}{2}\rangle, & \Psi^+|a, +\frac{1}{2}\rangle &= 0. \end{aligned} \quad (\text{A.8})$$

**Theorem A.1 (A. Vaintrob)** *For an  $L$ -component link  $\mathcal{L} \in S^3$  and for a tensor product of  $u(1|1)$ -modules  $V_{\mathbf{a}} = V_{a_1} \otimes \cdots \otimes V_{a_L}$ ,*

$$\nabla_{\mathbf{A}}(\mathcal{L}; e^{\mathbf{a}}) = a_k^{-1} \text{STr}_{\mathbf{a};k} \mathcal{T}_{u(1|1)}(Z_{\mathcal{A}}(\mathcal{L})) \quad (\text{A.9})$$

for any  $k$ ,  $1 \leq k \leq L$ .

If we define  $O \in \text{End } V_a$  by its action on the basis vectors as

$$O|a, +\frac{1}{2}\rangle = |a, +\frac{1}{2}\rangle, \quad O|a, -\frac{1}{2}\rangle = 0, \quad (\text{A.10})$$

then  $\text{STr}_{V_a} O = 1$  and, according to eq. (A.7), equation (A.9) is equivalent to

$$\nabla_{\mathbf{A}}(\mathcal{L}; e^{\mathbf{a}}) = a_k^{-1} \text{STr}_{V_{\mathbf{a}}} \left[ O_k \mathcal{T}_{u(1|1)}(Z_{\mathcal{A}}(\mathcal{L})) \right]. \quad (\text{A.11})$$

*Proof of Theorem 6.15.* We will prove the second line of eq. (6.45) by modifying the *r.h.s.* of eq. (A.11). We will simplify  $\mathcal{T}_{u(1|1)}(Z_{\mathcal{A}}(\mathcal{L}))$  and find a way to calculate the supertrace  $\text{STr}_{V_{\mathbf{a}}}$  in a way, which is similar to Kirillov's integral formula. Actually, we will work through the symmetric algebra  $Su(1|1)$ , that is, we will describe  $\mathcal{T}_{u(1|1)}(Z_{\mathcal{B}}(\mathcal{L})) \in Su(1|1)$  and then describe a trick to calculate a composition of maps

$$Su(1|1) \xrightarrow{\beta_{u(1|1)}} Uu(1|1) \xrightarrow{\text{STr}_{V_a}} \mathbb{C}, \quad (\text{A.12})$$

where  $\beta_{u(1|1)}$  is the PBW isomorphism (*cf.* eq. (2.14)).

Let  $D$  be a graph of  $\mathcal{B}_{\mathcal{L}}$  and  $D_F$  – one of its possible fermionic subgraphs. Suppose that  $D$  has a bosonic edge which is incident to two 3-vertices. According to eq. (A.3), the tensor  $h_B^{-1}$  has only two non-zero matrix elements:  $EF$  and  $FE$ , while the only bosonic index of the non-zero structure tensor elements (A.4) is  $F$ . Therefore, the  $E$  index of  $h_9^{-1}B$  can not be matched by structure tensors at 3-vertices, and  $\mathcal{T}_{u(1|1)}(D, D_F) = 0$ . Hence, if  $\mathcal{T}_{u(1|1)}(D, D_F) \neq 0$ , then every bosonic edge of  $D$  must be a leg. This means that there are only three types of pairs  $D, D_F$ ,  $D$  being connected, with non-zero  $u(1|1)$  weights: a fermionic strut, a haircomb with fermionic spine and a wheel with fermionic cycle and bosonic legs.

Let us find the  $u(1|1)$  weights of these graphs. We present  $(Su(1|1))^{\otimes L}$  as a polynomial algebra of  $4L$  variables  $E_i, F_i, \Psi_i^+, \Psi_i^-, 1 \leq i \leq L$ . Now if  $D$  is a bosonic strut  $X_i \text{ --- } X_j$ , then  $\mathcal{T}_{u(1|1)}(D, D_F) = E_i F_j + F_i E_j$ . If  $D$  is a haircomb graph of Fig. 1 with fermionic spine, then  $\mathcal{T}_{u(1|1)}(D, D_F) = (\Psi_i^- \Psi_j^+ - (-1)^{n_1 + \dots + n_L} \Psi_i^+ \Psi_j^-) E_1^{n_1} \dots E_L^{n_L}$ , where  $n_i$  ( $1 \leq i \leq L$ ) are the numbers of bosonic legs of colors  $\mathcal{L}_i$  attached to the fermionic spine. If  $D$  is a wheel with fermionic circle, then  $\mathcal{T}_{u(1|1)}(D, D_F) = -(1 - (-1)^{n_1 + \dots + n_L}) E_1^{n_1} \dots E_L^{n_L}$ .

We are going to convert  $\mathcal{T}_{u(1|1)}(D)$  into an element of  $Uu(1|1)$  and then evaluate its supertrace in the tensor product of 2-dimensional representation (A.8). Since  $E = aI$  in  $V_a$ , then for the purpose of our calculations we can replace  $E_i$  by  $a_i$  in our formulas from  $\mathcal{T}_{u(1|1)}(D, D_F)$ . Also, we know that the coefficients at the struts in  $W(\mathcal{L}; \mathbf{X})$  are linking numbers  $\ell_{ij}$  of  $\mathcal{L}$ , while the coefficients at its haircomb graphs form generating series  $l'_{ij}(\mathbf{a})$  of eq.(6.38) and the coefficients at wheels form the generating series  $F_0(\mathcal{L}; \mathbf{a})$  (see the discussion at the beginning of subsection 6.2). Therefore, for the purpose of evaluating the *r.h.s.* of eq.(A.11), we can replace  $\mathcal{T}_{u(1|1)}(Z_B(\mathcal{L})) \in (Su(1|1))^{\otimes L}$  by a rather simple exponential

$$\nabla_A(\mathcal{L}; e^{\mathbf{a}}) = a_k^{-1} e^{-F_0(\mathcal{L}; \mathbf{a})} \text{STr}_{V_{\mathbf{a}}} O_k \beta_{u(1|1)} \left[ \exp \left( \sum_{i,j=1}^L \left( \ell_{ij} a_i F_j - l'_{ij}(\mathbf{a}) \Psi_i^- \Psi_j^+ \right) \right) \right]. \quad (\text{A.13})$$

Now we describe a useful trick for calculating the composition of maps (A.12). This trick works for any Lie (super-)algebra  $\mathfrak{g}$  and for any  $\mathfrak{g}$ -module  $V$ . There is a canonical isomorphism  $\gamma_{\mathfrak{g}}: S\mathfrak{g} \rightarrow \mathfrak{D}_{\text{const}}(\mathfrak{g})$  between the symmetric algebra  $S\mathfrak{g}$  and the algebra of differential operators on  $\mathfrak{g}$  with constant coefficients  $\mathfrak{D}_{\text{const}}(\mathfrak{g})$ . Namely, for  $\vec{X} \in \mathfrak{g} \subset S\mathfrak{g}$ ,  $\gamma(\vec{X})$  is the Lie derivative along the constant vector field on  $\mathfrak{g}$ , which is equal to  $\vec{X}$  at every point. Then it is well-known that for any  $x \in S\mathfrak{g}$ ,

$$\text{STr}_V \beta_{\mathfrak{g}}(x) = \gamma_{\mathfrak{g}}(x) \left( \text{STr}_V e^{\vec{Y}} \right) \Big|_{\vec{Y}=0}, \quad (\text{A.14})$$

where in the *r.h.s.* the differential operator  $\gamma_{\mathfrak{g}}(x)$  is acting on the function  $\text{STr}_V e^{\vec{Y}}$  of  $\vec{Y} \in \mathfrak{g}$  and the resulting function is restricted at  $\vec{Y} = 0$ . We will also need a slight generalization of this equation: for any  $O \in \text{End}(V)$ ,

$$\text{STr}_V \left( O \beta_{\mathfrak{g}}(x) \right) = \gamma_{\mathfrak{g}}(x) \left( \text{STr}_V O e^{\vec{Y}} \right) \Big|_{\vec{Y}=0}. \quad (\text{A.15})$$

We introduce one bosonic coordinate  $\phi$  and two fermionic coordinates  $\eta^+, \eta^-$  on  $u(1|1)$

$$\vec{Y} = \phi F + \eta^+ \Psi^+ + \eta^- \Psi^- \quad (\text{A.16})$$

(we did not introduce a coordinate for  $E$ , because all operators  $E_j$  are already replaced by the corresponding variables  $a_j$ ). Then eqs. (A.14) and (A.15) allow us to rewrite eq. (A.13) as

$$\begin{aligned} \nabla_A(\mathcal{L}; e^{\mathbf{a}}) &= a_k^{-1} e^{-F_0(\mathcal{L}; \mathbf{a})} \\ &\times \exp \left( \sum_{i,j=1}^L \left( \ell_{ij} a_i \partial_{\phi_j} - l'_{ij}(\mathbf{a}) \partial_{\eta_i^-} \partial_{\eta_j^+} \right) \right) \text{STr}_{V_{a_k}} \left( O e^{\vec{Y}_k} \right) \prod_{\substack{1 \leq i \leq L \\ i \neq k}} \text{STr}_{V_{a_i}} e^{\vec{Y}_i} \Bigg|_{\vec{Y}=0}. \end{aligned} \quad (\text{A.17})$$

Now we calculate the traces by using a QFT trick. We introduce two (fermionic creation and annihilation) operators  $\hat{\psi}_+, \hat{\psi}_-$ , which satisfy the anti-commutation relations

$$\{\hat{\psi}_+, \hat{\psi}_+\} = \{\hat{\psi}_-, \hat{\psi}_-\} = 0, \quad \{\hat{\psi}_+, \hat{\psi}_-\} = I \quad (\text{A.18})$$

and act on  $V_a$  as

$$\begin{aligned} \hat{\psi}_+ |a, +\tfrac{1}{2}\rangle &= 0, & \hat{\psi}_+ |a, -\tfrac{1}{2}\rangle &= |a, +\tfrac{1}{2}\rangle, \\ \hat{\psi}_- |a, +\tfrac{1}{2}\rangle &= |a, -\tfrac{1}{2}\rangle, & \hat{\psi}_- |a, -\tfrac{1}{2}\rangle &= 0. \end{aligned} \quad (\text{A.19})$$

Comparing the matrix elements (A.8) and (A.19) we conclude that in  $V_a$

$$\Psi^- = \hat{\psi}_-, \quad \Psi^+ = a\hat{\psi}_+, \quad F = \frac{1}{2} I - \hat{\psi}_- \hat{\psi}_+. \quad (\text{A.20})$$

and

$$e^{\vec{Y}} = e^{\phi/2} \exp(-\phi \hat{\psi}_- \hat{\psi}_+ + a \eta^+ \hat{\psi}_+ + \eta^- \hat{\psi}_-). \quad (\text{A.21})$$

Let us define new operators

$$\hat{\psi}'_+ = \hat{\psi}_+ + \frac{1}{\phi} \eta^-, \quad \hat{\psi}'_- = \hat{\psi}_- - \frac{a}{\phi} \eta^+, \quad (\text{A.22})$$

so that eq. (A.21) becomes

$$\epsilon^{\vec{Y}} = e^{\phi/2} e^{\frac{a}{\phi} \eta^- \eta^+} \exp(-\phi \hat{\psi}'_- \hat{\psi}'_+). \quad (\text{A.23})$$

Since the operators  $\hat{\psi}'_+, \hat{\psi}'_-$  satisfy the same anticommutation relations (A.18) as  $\hat{\psi}_+, \hat{\psi}_-$ , then they have the same matrix elements (A.19) in an appropriate basis of  $V_a$ . Those basis vectors are eigenvectors of  $\exp(-\phi \hat{\psi}'_- \hat{\psi}'_+)$  with eigenvalues 1 and  $e^{-\phi}$ , so

$$\text{STr}_{V_a} \exp(-\phi \hat{\psi}'_- \hat{\psi}'_+) = 1 - e^{-\phi} \quad (\text{A.24})$$

and

$$\text{STr}_{V_a} e^{\vec{Y}} = (e^{\phi/2} - e^{-\phi/2}) e^{\frac{a}{\phi} \eta^- \eta^+}. \quad (\text{A.25})$$

Since for two fermionic variables  $\psi_+, \psi_-$

$$\int \exp(\phi \psi_+ \psi_- + a \eta^+ \psi_+ + \eta^- \psi_-) d\psi_- d\psi_+ = \phi e^{\frac{a}{\phi} \eta^- \eta^+}, \quad (\text{A.26})$$

then eq.(A.25) can be rewritten as

$$\text{STr}_{V_a} e^{\vec{Y}} = \frac{\sinh(\phi/2)}{(\phi/2)} \int \exp(\phi \psi_+ \psi_- + a \eta^+ \psi_+ + \eta^- \psi_-) d\psi_- d\psi_+. \quad (\text{A.27})$$

If we interpret  $\psi_+, \psi_-$  as coordinates on the coadjoint orbit of  $u(1|1)$ , then this equation represents Kirillov's integral formula for the character of this superalgebra.

Comparing the matrix elements (A.10) and (A.19), we find that in  $V_a$

$$O = \hat{\psi}_+ \hat{\psi}_- = (\hat{\psi}'_+ - \frac{1}{\phi} \eta^-)(\hat{\psi}'_- + \frac{a}{\phi} \eta^+), \quad (\text{A.28})$$

so that

$$\begin{aligned} \text{STr}_{V_a} O e^{\vec{Y}} &= e^{\phi/2} e^{\frac{a}{\phi} \eta^- \eta^+} \text{STr}_{V_a} \left[ (\hat{\psi}'_+ - \frac{1}{\phi} \eta^-)(\hat{\psi}'_- + \frac{a}{\phi} \eta^+) \exp(-\phi \hat{\psi}'_- \hat{\psi}'_+) \right] \\ &= \frac{\sinh(\phi/2)}{(\phi/2)} - \left( \frac{\sinh(\phi/2)}{(\phi/2)} - e^{\phi/2} \right) e^{\frac{a}{\phi} \eta^- \eta^+}, \end{aligned} \quad (\text{A.29})$$

Since

$$\int \psi_+ \psi_- \exp(\phi \psi_+ \psi_- + a \eta^+ \psi_+ + \eta^- \psi_-) d\psi_- d\psi_+ = 1, \quad (\text{A.30})$$

then we can rewrite eq.(A.29) in terms of convenient fermionic integrals

$$\begin{aligned} \text{STr}_{V_a} O e^{\vec{Y}} &= \frac{\sinh(\phi/2)}{(\phi/2)} \left( \int \psi_+ \psi_- \exp(\phi \psi_+ \psi_- + a \eta^+ \psi_+ + \eta^- \psi_-) d\psi_- d\psi_+ \right. \\ &\quad \left. - f_B(\phi) \int \exp(\phi \psi_+ \psi_- + a \eta^+ \psi_+ + \eta^- \psi_-) d\psi_- d\psi_+ \right), \end{aligned} \quad (\text{A.31})$$

where we defined a shortcut function

$$f_B(x) = \frac{1}{x} \left( 1 - \frac{x}{1 - e^{-x}} \right). \quad (\text{A.32})$$

Let us substitute eqs.(A.27) and (A.31) into eq.(A.17). Since for any analytic function  $f(\phi)$  and for any constant  $c$  we have a Taylor series formula

$$\exp(c \partial_\phi) f(\phi)|_{\phi=0} = f(c) \quad (\text{A.33})$$



and since

$$\begin{aligned} \exp \left( - \sum_{i,j=1}^L l'_{ij}(\mathbf{a}) \partial_{\eta_i^-} \partial_{\eta_j^+} \right) \exp \left( \sum_{i=1}^L (\eta_i^+ \psi_{+,i} + \eta_i^- \psi_{-,i}) \right) \Big|_{\boldsymbol{\eta}=0} \\ = \exp \left( \sum_{i,j=1}^L l'_{ij}(\mathbf{a}) \psi_{+,j} \psi_{-,i} \right), \end{aligned} \quad (\text{A.34})$$

then in view of eq.(6.40)

$$\begin{aligned} \nabla_{\mathbf{A}}(\mathcal{L}; e^{\mathbf{a}}) = a_k^{-1} e^{-F_0(\mathcal{L}; \mathbf{a})} \Omega_{\nabla}(\mathcal{L}; \mathbf{a}) \left[ \int \psi_{+,k} \psi_{-,k} \exp \left( \sum_{i,j=1}^L a_i^{-1} l_{ij}(\mathbf{a}) \psi_{+,j} \psi_{-,i} \right) d\boldsymbol{\psi}_- d\boldsymbol{\psi}_+ \right. \\ \left. - f_{\mathbf{B}} \left( \sum_{i=1}^L \ell_{ik} a_i \right) \int \exp \left( \sum_{i,j=1}^L a_i^{-1} l_{ij}(\mathbf{a}) \psi_{+,j} \psi_{-,i} \right) d\boldsymbol{\psi}_- d\boldsymbol{\psi}_+ \right], \end{aligned} \quad (\text{A.35})$$

where  $d\boldsymbol{\psi}_- d\boldsymbol{\psi}_+ = d\psi_{-,1} d\psi_{+,1} \cdots d\psi_{-,L} d\psi_{+,L}$ . Since for any matrix  $\|A_{ij}\|$

$$\int \exp \left( \sum_{i,j=1}^L A_{ij} \psi_{+,j} \psi_{-,i} \right) d\boldsymbol{\psi}_- d\boldsymbol{\psi}_+ = \det \|A_{ij}\|, \quad (\text{A.36})$$

$$\int \psi_{+,k} \psi_{-,k} \exp \left( \sum_{i,j=1}^L A_{ij} \psi_{+,j} \psi_{-,i} \right) d\boldsymbol{\psi}_- d\boldsymbol{\psi}_+ = \det \|A_{ij}\|_k, \quad (\text{A.37})$$

where  $\|A_{ij}\|_k$  is the  $(k, k)$ -minor of  $\|A_{ij}\|$ , then eq.(A.35) can be rewritten as

$$\begin{aligned} \nabla_{\mathbf{A}}(\mathcal{L}; e^{\mathbf{a}}) = a_k^{-1} e^{-F_0(\mathcal{L}; \mathbf{a})} \Omega_{\nabla}(\mathcal{L}; \mathbf{a}) \\ \times \left[ \det \|a_i^{-1} l_{ij}(\mathbf{a})\|_k - f_{\mathbf{B}} \left( \sum_{i=1}^L \ell_{ik} a_i \right) \det \|a_i^{-1} l_{ij}(\mathbf{a})\| \right] \end{aligned} \quad (\text{A.38})$$

Then eq.(6.45) follows, since

$$\det \|a_i^{-1} l_{ij}(\mathbf{a})\|_k = \frac{\det \|l_{ij}(\mathbf{a})\|_k}{\prod_{\substack{1 \leq j \leq L \\ j \neq k}} a_j}, \quad \det \|a_i^{-1} l_{ij}(\mathbf{a})\| = \frac{\det \|l_{ij}(\mathbf{a})\|}{\prod_{j=1}^L a_j} \quad (\text{A.39})$$

and the latter expression is zero in view of eq.(5.49).  $\square$

**Remark A.2** Equation (5.49) can be proved without using the machinery of Cartan and root edges. Indeed, consider the *r.h.s.* of eq.(A.11) if we choose  $O = I$ . Since  $\text{STr}_{V_a} I = \text{sdim } V_a = 0$ , then

$$\text{STr}_{V_{\mathbf{a}}} \left[ \mathcal{T}_{u(1|1)} \left( Z_{\mathcal{A}}(\mathcal{L}) \right) \right] = 0. \quad (\text{A.40})$$

If we calculate the *l.h.s.* of this equation in the same way that we did it for the *r.h.s.* of eq.(A.11), then we find that similarly to eq.(A.38)

$$\text{STr}_{V_{\mathbf{a}}} \left[ \mathcal{T}_{u(1|1)} \left( Z_{\mathcal{A}}(\mathcal{L}) \right) \right] = e^{-F_0(\mathcal{L}; \mathbf{a})} \Omega_{\nabla}(\mathcal{L}; \mathbf{a}) \det ||a_i^{-1} l_{ij}(\mathbf{a})||. \quad (\text{A.41})$$

Now a combination of eqs.(A.40) and (A.41) indicates that

$$\det ||a_i^{-1} l_{ij}(\mathbf{a})|| = 0, \quad (\text{A.42})$$

which is what we need to derive eq.(6.45) from eq.(A.38).

## References

- [1] D. Bar-Natan, *On The Vassiliev knot invariants*, Topology **34** (1995) 423-472.
- [2] D. Bar-Natan, S. Garoufalidis, L. Rozansky, D. Thurston, *Wheels, wheeling and the Kontsevich integral of the unknot*, Israel J. Math. **119** (2000) 217-238.
- [3] D. Bar-Natan, S. Garoufalidis, L. Rozansky, D. Thurston, *The Århus integral of rational homology 3-spheres II: invariance and universality*, Selecta Math., in press.
- [4] S. Garoufalidis, A. Kricker, *A rational noncommutative invariant of boundary links*, preprint math.GT/0105028.
- [5] N. Habegger, G. Masbaum, *The Kontsevich Integral and Milnor's Invariants*, Topology **39** (2000) 1253-1289.
- [6] L. Kauffman, H. Saleur, *Free fermions and the Alexander-Conway polynomial*, Commun. Math. Phys. **141** (1991) 473-545.
- [7] A. Kricker, *The lines of the Kontsevich integral and Rozansky's rationality conjecture*, preprint math.GT/0005284.
- [8] T.T.Q. Le, J. Murakami, T. Ohtsuki, *A universal quantum invariant of 3-manifolds*, Topology **37** (1998) 539-574.
- [9] J. Levine, *A factorization of the Conway polynomial*, Comment. Math. Helveticii **74** (1999) 1-27.
- [10] G. Masbaum, A. Vaintrob, *Milnor numbers, Spanning Trees, and the Alexander-Conway Polynomial*, preprint math.GT/0111102.

- [11] L. Rozansky, The universal  $R$ -matrix, Burau representation and the Melvin-Morton Expansion of the colored Jones polynomial, Adv. in Math. **134** (1998) 1-31.
- [12] L. Rozansky, *A contribution of a  $U(1)$ -reducible connection to quantum invariants of links I:  $R$ -matrix and Burau representation*, preprint math.QA/9806004.
- [13] L. Rozansky, *A contribution of a  $U(1)$ -reducible connection to quantum invariants of links II: Links in rational homology spheres*, preprint math.QA/9806066.
- [14] L. Rozansky, *A rationality conjecture about Kontsevich integral of knots and its implications to the structure of the colored Jones polynomial*, preprint math.GT/0106097.
- [15] L. Rozansky, in preparation.
- [16] A. Vaintrob, preprint.